79.2%) or subcutaneous (n=11; 20.8%). The initial dose was 8mg/kg/IV/4 weeks or 162mg/SC/week, respectively. TCZ was used in monotherapy or combined with immunosuppressants (n=32; 60.4%): methotrexate (n=27), azathioprine (n=2), cyclosporine (n=3). Main clinical features at TCZ onset were: malaise (n=30), limb claudication (n=22), headache (n=18), fever (n=14), abdominal pain (n=10), and chest pain (n=9). Most of the patients experienced a rapid and maintained clinical, analytical improvement (TABLE). After a median follow-up of 18.0 [7.0-45.0] months, TCZ was discontinued in 20 patients due to: sustained remission (n=6), relapse (n=6), adverse event (n=5), gestation (n=3). Most relevant adverse side effects were serious infections: pneumonia (n=2), herpes zoster (n=1), abdominal sepsis (n=1).

Table.

	Basal (N=53)	Month 1 (N=53)	Month 3 (N=46)	Month 6 (N=44)	Month 12 (N=34)
Clinical improvement, n/N(%)					
Complete		17/53 (32.1)	19/46 (41.3)	23/44 (52.3)	26/34 (76.5)
Partial		30/53 (54.6)	26/46 (56.5)	18/44 (40.9)	8/34 (23.5)
No improvement		6/53 (11.3)	1/46 (2.2)	3/44 (6.8)	0/34 (0.0)
Analytical markers,					
ESR (mm/1 st h), median [IQR]	35.0 [16.0-52.0]	7.5 [3.0-14.0] *	3.5 [2.0-8.0] *	5.0 [2.0-6.0] *	5.0 [2.0-8.5] *
CRP (mg/dL), median [IQR]	1.7 [0.6-3.5]	0.21 [0.05-0.6] *	0.14 [0.05-0.5] *	0.14 [0.04-0.4] *	0.10 [0.03-0.30] *
Hb (g/dL), mean ± SD	12.3±1.5	12.8±1.2 *	12.9±1.3 *	12.9±1.4 *	12.9±1.4 *
Prednisone dose (mg/day), median [IQR]	30.0 [15.0-50.0]	20.0 [10.0-37.5] *	10.0 [5.0-20.0] *	5.0 [5.0-12.5] *	5.0 [0.0-7.5] *

*Wilcoxon test p < 0.001.

Conclusion: TCZ appears to be effective and safe in patients with refractory TAK in clinical practice.

References:

- [1] Nakaoka Y et al. Ann Rheum Dis. 2018;77:348-354
- [2] Loricera J et al. Clin Exp Rheumatol. 2016; 34: S44-53.

Disclosure of Interests: D. Prieto-Peña: None declared, Monica Calderón-Goercke: None declared, Pilar Bernabéu: None declared, Paloma Vela-Casasempere: None declared, J. Narváez: None declared, Carlos Fernández-López: None declared, Mercedes Freire González: None declared, Beatriz González-Alvarez: None declared, Roser Solans-Laqué: None declared, Jose Luis Callejas-Rubio: None declared, Norberto Ortego: None declared, Carlos Fernández-Díaz Speakers bureau: Bristol Meyers Squibb, Esteban Rubio Romero: None declared, SALVADOR GARCÍA MORILLO: None declared, Mauricio Minguez: None declared, Cristina Fernández-Carballido Consultant of: Yes, I have received fees for scientific advice (Abbvie, Celgene, Janssen, Lilly and Novartis), Speakers bureau: Yes, I have received fees as a speaker (Abbvie, Celgene, Janssen, Lilly, MSD, Novartis), Eugenio de Miguel Grant/research support from: Yes (Abbvie, Novartis, Pfizer), Consultant of: Yes (Abbvie, Novartis, Pfizer), Paid instructor for: yes (AbbVie, Novartis, Pfizer, MSD, BMS, UCB, Roche, Grunental, Janssen, Sanofi), Speakers bureau: yes (AbbVie, Novartis, Pfizer, MSD, BMS, UCB, Roche, Grunental, Janssen, Sanofi), Sheila Melchor: None declared, Eva Salgado-Pérez: None declared, Beatriz Bravo: None declared, Susana Romero-Yuste: None declared, J Salvatierra: None declared, Cristina Hidalgo: None declared, Sara Manrique Arja: None declared, C. Romero-Gómez: None declared, Patricia Moya: None declared, Noelia Alvarez-Rivas: None declared, Javier Mendizabal: None declared, Francisco Miguel Ortiz Sanjuan: None declared, I. Pérez de Pedro: None declared, Javier Loricera: None declared, Santos Castañeda: None declared, Miguel A González-Gay Grant/research support from: Pfizer, Abbvie, MSD, Speakers bureau: Pfizer, Abbvie, MSD, Ricardo Blanco Grant/research support from: AbbVie, MSD, Roche, Consultant of: Abbvie, Eli Lilly, Pfizer, Roche, Bristol-Myers, Janssen, UCB Pharma and MSD, Speakers bureau: Abbvie, Eli Lilly, Pfizer, Roche, Bristol-Myers, Janssen, UCB Pharma. MSD

DOI: 10.1136/annrheumdis-2020-eular.2445

SAT0271

THE "GOLD STANDARD" DIAGNOSTIC TEST FOR GCA IS THE WHOLE GCA FAST TRACK PATHWAY COMBINED

V. Quick¹, M. Hughes¹, C. H. Chan². ¹Luton and Dunstable Hospital, Rheumatology, Luton, United Kingdom; ²Luton and Dunstable Hospital, Maxillofacial Surgery, Luton, United Kingdom

Background: We have been developing a rheumatologist-led ultrasound driven giant cell arteritis (GCA) fast-track pathway (FTP), which in year 3 had the following structure:

1. **Rapid access to rheumatology assessment (RAS)** to establish clinical probability of GCA (CP-GCA). No referral criteria required
2. **Temporal and axillary artery ultrasound (TAUS)** if GCA not excluded with RAS. TAUS considered positive if bilateral (halo score $\geq 2/8^{1,2}$ at >1 temporal artery)
3. **Second test in selected patients:** GCA diagnosed in those with mod-high CP-GCA and +ve TAUS and excluded in those with low CP-GCA and -ve TAUS. All others had biopsy (TAB) or large vessel imaging (LVI), presentation depending
4. **Protocolised withdrawal of prednisolone:** Patients only treated for GCA if ≥ 1 of: high CP-GCA, +ve TAUS, TAB or LVI
5. **Rapid access** if symptoms recurred on steroid withdrawal for RAS + TAUS

Objectives:

- To compare security of GCA diagnosis, sight loss rate and TAB rate in Year 3 to previous years
- To assess sensitivity and specificity of all components of the Yr 3 FTP for diagnosis of GCA

Methods: As in Yr 2, TAUS was performed by VQ with an Esaote Mylab7, 6-15MHz probe for axillaries, 22MHz for temporal arteries (TAs). In Yr 1 VQ used a GES8 with ML6-15 for axillaries, 18MHz probe for TAs. Year 3 audit data were compared to previous audits

Results:

Year 3 Luton GCA FTP compared to previous years

	Traditional Model	Year 1	Year 2	Year 3
Audit period	2015 12 months	1/1/16-31/3/17 15 months	1/4/17-31/3/18 12 months	1/4/18-31/3/19 12 months
Pathway structure	No rapid access to RAS TAB, no TAUS service	Patients seen ad-hoc for RAS + TAUS TAB requested in all appropriate cases to compare TAUS to TAB	Rapid access for RAS + TAUS, 2 slots/wk 2nd test in selected patients	Rapid access for RAS + TAUS, 3 slots /wk 2nd test in selected patients Protocolised withdrawal prednisolone
No. referrals with suspected GCA/yr	NK	50.4	70	90
Cases GCA /yr	9	24	18	25
Cases GCA excluded /yr	NK	26.4	52	65
% GCA patients with imaging/TAB +ve GCA	44.4	63.3	83.3	92
% GCA patients with clinical GCA (high CP-GCA, no +ve test)	55.6	36.7	16.6	8
% referred where GCA excluded on clinical grounds alone	NK	11.1	5.7	5.6
% referred where TAUS performed and TAB avoided	NA	NA	38.6	50.0
% referred who had TAB	NK (total 27)	84.1	42.8	31.1
% referred where TAUS, then LVI performed instead of TAB	NA	4.8	12.8	13.3
Mean days on prednisolone for GCA before TAUS (median, range)	NA	10.6 (7, 0-81)	4.4 (2, 0-54)	5.9 (3, 0-57)
% GCA patients with permanent sight loss due to GCA	33	23.3	11.2	8

NA: Not applicable NK: not known

Performance of components of Year 3 Luton GCA FTP compared to final diagnosis at >6 months*

Diagnostic Tool	No. pts**	Sensitivity %	Specificity %	PPV %	NPV %
Rheumatologist CP GCA (High)***	90	60	98.3	93.8	86.1
Rheumatologist CP GCA (Moderate & High)***	90	88	63.5	48.9	93
Any halo on TAUS	85	81.5	91.4	81.5	91.4
Bilateral positive TAUS	85	59.3	98.3	94.1	83.8
TAB	28	69.2	100	100	78.9
LVI (CT or CTPET)	12	80	100	100	87.5
Combined diagnostic tests in the whole FTP (bilateral positive TAUS +/- or positive TAB +/- or positive LVI)	88	92	98.3	95.8	96.7

*using patient record to Oct 2019 **2 patients passed through twice ***before imaging/TAB

Conclusion: Unlike Yr 2, the higher secure diagnosis rate in Yr 3 could not be attributed to shorter time on prednisolone or better equipment. The increase was likely due to several factors including further improved sonographer skill and increased confidence to withdraw steroids in insecure cases with low/moderate CP-GCA. This approach did not increase sight loss. Further reduction in TAB rate financially justified a 3rd FTP slot/wk created in Yr 3.

Each component of the FTP was an inadequate diagnostic tool. Combinations of diagnostic tools are needed to obtain the highest sensitivity and specificity for GCA diagnosis. FTPs limit tests to the minimum required for secure diagnosis. The "gold standard" diagnostic test for GCA is the whole FTP combined.

References:

- [1] Schäfer et al *Rheum (Ox)* 2017;56(9):1479-83
- [2] van der Geest et al *ARD* doi: 10.1136/annrheumdis-2019-216343

Disclosure of Interests: Vanessa Quick Consultant of: Roche, Speakers bureau: Roche, Mark Hughes: None declared, Chi-Hwa Chan: None declared

DOI: 10.1136/annrheumdis-2020-eular.2038

SAT0272

THE VALUE OF CLINICAL AND LABORATORY FEATURES TO PREDICT EXTENT OF LARGE VESSEL VASCULITIS ON PET CT

G. Reynolds¹, C. Holland¹, G. Petrides¹, A. Lorenzi¹, B. Thompson¹, J. Heaney¹, T. Ali¹. ¹Newcastle Freeman Hospital, High Heaton, United Kingdom

Background: Giant cell arteritis (GCA) is an idiopathic vasculitis affecting large and medium-sized vessels. The pattern of arterial involvement is heterogeneous with two overlapping categories recognised: classical cranial GCA and extra-cranial GCA (or large vessel vasculitis – LVV) that predominantly affects the aorta and its proximal branches. Although LVV is present in around 80% of patients with cranial GCA, and around one third will develop large vessel complications, there are no guidelines for which patients should be screened for it (1). We sought to investigate whether clinical and laboratory features were a useful guide to the severity of LVV on FDG PET-CT.

Objectives: To retrospectively analyse whether baseline patient characteristics are able to predict the extent of large vessel vasculitis on PET-CT.

Methods: Clinical data for 65 patients referred for a PET-CT scan by Rheumatology at the Freeman Hospital, Newcastle between January 2015 and May 2018 were retrospectively analysed. The most recent full blood count and inflammatory markers prior to the scan were used. Scans were reviewed by a consultant radiologist and trainee. The arterial network was split in to ten potentially involved territories (aortic arch, thoracic aorta, abdominal aorta, iliac vessels, axillary, brachiocephalic, subclavian, carotid, vertebral and femoral arteries. Both the value of highest standardised uptake value (SUV max) and the territory affected was recorded for each positive scan.

Results: In the period analysed 65 PET-CT scans were requested, mostly (77%) as baseline investigations for symptoms with LVV in the differential diagnosis. Of these 22 (34%) were positive for LVV and in that group the majority of patients (64%) were female. In those with a negative scan, 47.5% were on concurrent steroid treatment compared to 9% with a positive scan. Regression analysis suggested that the number of systemic features (weight loss, pyrexia, polymyalgia) was weakly correlated with the number of affected territories ($p=0.04$). In contrast there was no correlation between laboratory tests ((CRP ($p=0.91$), ESR ($p=0.46$), Hb ($p=0.44$), platelets ($p=0.74$)) and the number of territories affected. The aortic arch (47%) was most commonly the territory with the highest degree of FDG

uptake (SUV max) followed by the abdominal aorta (21%) and thoracic (10%) and femoral arteries (10%). There was no correlation between SUV max and laboratory tests ((CRP ($p=0.55$), ESR ($p=0.89$), Hb ($p=0.82$), platelets ($p=0.17$)) or the number of systemic features ($p=0.7$). There was no significant difference in the number of territories affected between those on steroid treatment at the time of the scan and steroid-naïve patients, albeit the number of positive scans in those on steroid treatment was low ($n=5$).

Conclusion: These results suggest that clinical and laboratory features are a poor guide to predicting the maximal severity and extent of disease on FDG PET-CT.

References: [1]Koster MJ, Matteson EL, Warrington KJ. Large-vessel giant cell arteritis: diagnosis, monitoring and management. *Rheumatology (Oxford)*. 2018;57(suppl_2):ii32-ii42.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.2452

SAT0273

PROGNOSTIC FACTORS FOR POOR OUTCOME IN BEHÇET'S SYNDROME WITH NEUROLOGICAL INVOLVEMENT: RESULTS FROM A CLINICAL LONG-TERM FOLLOW-UP OF A SINGLE CENTRE

D. Sahin Eroglu¹, M. Torgutalp¹, S. Sezer¹, A. Boyvat², C. Yucesan³, A. Ates¹. ¹Ankara University Faculty of Medicine, Department of Rheumatology, Ankara, Turkey; ²Ankara University Faculty of Medicine, Department of Dermatology, Ankara, Turkey; ³Ankara University Faculty of Medicine, Department of Neurology, Ankara, Turkey

Background: Behçet's Syndrome (BS) is a vasculitis of unknown cause. Neurological disease is a type of its serious involvement and causes morbidity and mortality.

Objectives: In this study, we aimed to describe the clinical characteristics of neuro-Behçet Syndrome (NBS) and to define prognostic factors that were associated with poor outcome.

Methods: Among 2033 patients with BS, we performed a retrospective analysis of 94 patients (52.1% male; mean age 36.1 (11.9) years), who fulfilled the ISG-1990 for BS. We divided patients into two subgroups, either parenchymal (p-NBS) or non-parenchymal (np-NBS). Clinical, laboratory, and imaging characteristics of the patients were defined. We described the poor outcome as a modified Rankin score (mRS) ≥ 3 at last follow-up and/or death, and assessed the predictor factors associated such kind of outcome.

Results: In total, 52 (55.3%) patients presented with p-NBS, of whom 15 (28.8%) had progressive course, and 42 (44.7%) presented with np-NBS (Table.1). Ocular involvement was more common in p-NBS than np-NBS (59.6% vs. 39.1% respectively, $p=0.04$), whereas vascular involvement excluding sinus vein thrombosis was more frequent in patients with np-NBS (15.4% vs. 50.0%, respectively, $p=0.001$). Headache and papilledema were frequently observed in np-NBS ($p<0.001$), whereas paresis, sensory symptoms, pyramidal and cerebellar signs were significantly common in p-NBS (<0.001). Twenty-five patients (26.6%) experienced at least one relapse. After a median of 54 (IQR 79.4) months, 11 patients (11.7%) had disabling

Table 1. Clinical characteristics of the patients in each subgroup

	All n= 94	Parenchymal NBS n= 52	Non parenchymal NBS n= 42	p value
Gender (male, %)	49 (52.1)	29 (55.8)	20 (47.6)	0.43
Age at Behçet diagnosis	29.2 $\pm$ 9.52	29.93 $\pm$ 9.49	28.26 $\pm$ 9.60	0.40
Age at NBS diagnosis	36.1 $\pm$ 11.9	37.02 $\pm$ 11.43	35.00 $\pm$ 12.61	0.42
Initial mRS ≥ 3	33 (35.1)	28 (53.8)	5 (11.9)	<0.001
Oral aphthae	94 (100.0)	52 (100.0)	42 (100.0)	-
Genital ulcer	71 (75.5)	40 (76.9)	31 (73.8)	0.73
Skin involvement (PPE+EN)	72 (76.6)	35 (67.3)	37 (88.1)	0.018
Musculoskeletal involvement	46 (48.9)	28 (53.8)	18 (42.8)	0.29
Ocular involvement	47 (50)	31 (59.6)	16 (38.1)	0.038
Vascular involvement (except SVT)	29 (30.9)	8 (15.4)	21 (50)	<0.001
Gastrointestinal involvement	5 (5.3)	2 (3.8)	3 (7.1)	0.65
HLA B51 positivity (n=32)	14 (43.8)	6/12 (50)	8/20 (40)	0.58
Pathergy positivity (n=51)	26 (51.0)	13 (54.2)	13 (48.1)	0.67
Clinical characteristics				
Headache	73 (77.7)	32 (61.5)	41 (97.6)	<0.001
Fever	6 (6.4)	5 (9.6)	1 (2.4)	0.22
Paresis	26 (27.7)	24 (46.2)	2 (4.8)	<0.001
Pyramidal signs	34 (36.2)	31 (59.6)	3 (7.1)	<0.001
Sensory symptoms	34 (36.2)	30 (57.7)	4 (9.5)	<0.001
Cerebellar signs	19 (20.2)	18 (34.6)	1 (2.4)	<0.001
Cranial nerve involvement	13 (13.8)	9 (7.3)	4 (9.5)	0.28
Papilledema	24 (25.5)	3 (5.8)	21 (50)	<0.001
Relapse	25 (26.6)	16 (30.8)	9 (21.4)	0.31
Total follow-up time (mo)	54.01 (IQR: 79.4)	64.8 (IQR: 90.2)	44.0 (IQR: 68.1)	0.12
Last mRS ≥ 3	16 (16.8)	15 (28.8)	1 (2.4)	0.001
Death	5 (5.3)	5 (9.6)	0 (0)	0.06

mRS: Modified Rankin Score, PPE: Papulopustular eruption, EN: Erythema nodosum, SVT: Sinus Vein Thrombosis