department from Genesis Pharma (2018), Speakers bureau: Received honoraria from Genesis Pharma and Janssen(2017) and from Roche and Pharmaserve Lilly(2018), Efterpi Zafiriou Speakers bureau: Received honoraria from Genesis Pharma, Abbvie, Novartis, Roche, Jansses(2017) and Novartis, Abbvie(2018), Dimitrios Bogdanos: None declared, Lazaros Sakkas Grant/research support from: Received a grant to support the educational and research activities of the department from Bristol-Myers Squibb, Speakers bureau: Received honoraria from Actellion(2018), Janssen(2017), Novartis(2017), Sanofi-Aventis(2018), Abbvie(2017) and Roche(2017)

DOI: 10.1136/annrheumdis-2020-eular.4147

SATURDAY, 06 JUNE 2020

Spondyloarthritis - clinical aspects (other than treatment)

SAT0364

DATA TO BE COLLECTED FOR AN OPTIMAL MANAGEMENT OF AXIAL SPONDYLOARTHRITIS IN DAILY PRACTICE: PROPOSAL FROM AN EVIDENCE BASED AND CONSENSUAL APPROACHES

A. Baillet¹, X. Romand², A. Pfimlin³, M. Dalecky¹, M. Dougados⁴ on behalf of the "Rencontre d'Expert en Rhumatologie" Task Force. ¹Univ. Grenoble Alpes, GREPI TMC CNRS UMR 5525, Grenoble, France; ²Univ. Grenoble Alpes, GREPI-TMC CNRS UMR 5525, Grenoble, France; ³Rheumatology, R Salengro Hospital, University of Lille, Lille, France; ⁴Department of Rheumatology - Hôpital Cochin. Assistance Publique - Hôpitaux de Paris, Paris, France

Background: Standardization of clinical practice has been proven to be effective in management of chronic diseases. This is particularly true at the time where the concept of treat to target is becoming more and more important in the field of axial spondyloarthritis (ax-SpA).

Objectives: To propose a list of variables to be collected at the time of the diagnosis and over the follow-up of patients with axial spondyloarthritis (ax-SpA) for an optimal management in daily practice.

Methods: The process comprised (1) the evaluation of the interest of 51 variables proposed for the assessment of axSpA via a systematic literature research, (2) a consensus process involving 78 hospital-based or office-based rheumatologists, considering the collection of the variable in a 4 grade scale from "potentially useful" to "mandatory", (3) a consensus on optimal timeline for periodic

assessment of the selected variables on a 5 grade scale from "at each visit" to "never to be re-collected".

Results: The systematic literature research retrieved a total of 14,133 abstracts, of which 213 were included in the final qualitative synthesis. Concerning the data to be collected at the time of the diagnosis and during follow-up, we proposed to differentiate the results based on a) the way of collection of the variables (e.g. questionnaires by the patient, interview by the physician, physical examination, investigations) b) the usefulness these variables in daily practice based on the opinion of the rheumatologists " c) the optimal timeline between 2 evaluations of the variable based on the opinion of the rheumatologists. In the initial systematic review, symptoms of heart failure history of inflammatory bowel disease, psoriasis or uveitis, patient global visual analogic scale, spine radiographs, modified Schöber test, coxo-femoral rotations, swollen joint count, urine strip test, BASDAI and ASDAS global scores were considered very useful and nocturnal back pain/ morning stiffness, sacro-iliac joints radiographs and CRP were considered mandatory (Figure 1). Timeline between 2 evaluations of variables to collect in the periodic review are summarized in Figure 2.

Figure 2. Periodic review timeline of variables to collect ASDAS=Ankylosing Spondylitis Disease Activity Score, BASDAI=Bath Ankylosing Spondylitis Disease Activity Index, Spondylitis Metrology Index, CRP=C Reactive Protein, IBD = inflammatory bowel diseases, PRO = Patient Reported Outcomes

Conclusion: Using an evidence-based and an expert consensus approaches, this initiative defined a core set of variables to be collected and reported at the time of the diagnosis and during follow-up of patients with ax-SpA in daily practice.

Acknowledgments: this study has been conducted in two parts: the first one (evidence-based) was conducted thanks to a support from Abbvie France. AbbVie did not review the content or have influence on this manuscript. The second part of this initiative (consensus) has been conducted thanks to a support from the scientific non-profit organization: Association de Recherche Clinique en Rhumatologie

Disclosure of Interests: Athan Baillet Consultant of: Athan BAILLET has received honorarium fees from Abbvie for his participation as the coordinator of the systematic literature review, Xavier Romand Consultant of: Xavier ROMAND has received honorarium fees from Abbvie, Arnaud Pfimlin Consultant of: Arnaud PFIMLIN has received honorarium fees from Abbvie, Mickael Dalecky Consultant of: Mickael DALECKY has received honorarium fees from Abbvie, Maxime Dougados Grant/ research support from: AbbVie, Eli Lilly, Merck, Novartis, Pfizer and UCB Pharma, Consultant of: AbbVie, Eli Lilly, Merck, Novartis, Pfizer and UCB Pharma, Speakers bureau: AbbVie, Eli Lilly, Merck, Novartis, Pfizer and UCB Pharma

DOI: 10.1136/annrheumdis-2020-eular.3588

SAT0365

EFFECTS OF ANTI-TNF-THERAPY ON OSTEOBLASTIC ACTIVITY IN ANKYLOSING SPONDYLITIS – RESULTS FROM A PROSPECTIVE STUDY USING PET-MRI OF SIJ AND SPINE

X. Baraliakos¹, S. Tsiemi¹, C. Rischpler², N. M. Bruckmann³, W. Fendler², J. Kirchner³, K. Hermann³, L. Sawicki³, J. Braun¹. ¹Rheumazentrum Ruhrgebiet Herne, Ruhr-University Bochum, Herne, Germany; ²Dpt. of Nuclear Medicine, University Essen-Duisburg, Essen, Germany; ³Dpt. of Radiology, University Duesseldorf, Duesseldorf, Germany

Background: The clinical efficacy of tumor necrosis factor inhibitors (TNFi) in patients with axial spondyloarthritis (axSpA) is well established but its effect on new bone formation is still unclear (1). Positron emission tomography (PET) using bone-seeking ¹⁸F-Fluoride [¹⁸F]F in combination with magnetic resonance imaging ([¹⁸F]F /MRI) has been shown to depict not only bone marrow edema (BME) but also shows the quantity of tracer uptake in the late phase of perfusion suggestive of remodeling and osteoblastic activity, not only in radiographic axSpA (r-axSpA) (2).

Objectives: Assess the effect of TNFi on bone remodeling processes in the axial skeleton of r-axSpA patients using [¹⁸F]F/MRI prior (baseline, BL) and 4 months after (follow-up, FU) treatment.

Figure 1. Core sets of items to collect and report in the systematic review in axial spondyloarthritis management in daily practice ASDAS=Ankylosing Spondylitis Disease Activity Score, BASDAI=Bath Ankylosing Spondylitis Disease Activity Index, BASFI=Bath Ankylosing Spondylitis Functional Index, BASMI=Bath Ankylosing Spondylitis Metrology Index, CRP=C Reactive Protein, CT=computerized tomography, FIRST=Fibromyalgia Rapid Screening Tool, HLA=Human Leukocyte Antigen, MRI=Magnetic resonance imaging, PET=positron emission tomography.