

Table 2. Safety Profile: Number of participants (%) experiencing any event

	PBO	BIIB059				OVERALL
	N=33	50 mg N=26	150 mg N=25	450 mg N=48	Pooled N=99	N=132
Any Event, n(%)	18 (54.5)	17 (65.4)	12 (48)	33 (68.8)	62 (62.6)	80 (60.6)
Severity						
Mild	11 (33.3)	11 (42.3)	8 (32.0)	19 (39.6)	38 (38.4)	49 (37.1)
Moderate	4 (12.1)	6 (23.1)	3 (12.0)	12 (25.0)	21 (21.2)	25 (18.9)
Severe	3 (9.1)	0	1 (4.0)	2 (4.2)	3 (3.0)	6 (4.5)
Related events	6 (18.2)	9 (34.6)	4 (16.0)	16 (33.3)	29 (29.3)	35 (26.5)
Serious events	2 (6.1)	0	3 (12.0)	2 (4.2)	5 (5.1)	7 (5.3)
Related serious events	1 (3.0)	0	1 (4.0)	1 (2.1)	2 (2.0)	3 (2.3)
Events leading to drug withdrawal	0	1 (3.8)	1 (4.0)	1 (2.1)	3 (3.0)	3 (2.3)
Events leading to study withdrawal	0	0	0	1 (2.1)	1 (1.0)	1 (0.8)
Fatal events	0	0	0	0	0	0

Alliance, Sanford Consortium, Consultant of: Genentech, Nektar, BMS, Janssen, AstraZeneca, Biogen, Vielabio, Equillum, Eli Lilly, ILTOO, Abbvie, Amgen, Roche, Gilead, Ronald van Vollenhoven Grant/research support from: AbbVie, Arthrogen, Bristol-Myers Squibb, GlaxoSmithKline, Lilly, Pfizer, and UCB, Consultant of: AbbVie, AstraZeneca, Biotest, Bristol-Myers Squibb, Celgene, GSK, Janssen, Lilly, Medac, Merck, Novartis, Pfizer, Roche, and UCB, Filippa Nyberg Consultant of: Biogen, Benjamin Kaffenberger Grant/research support from: Amgen, Biogen, InflaRx, Veloce Biopharmaceuticals, Dermatology Foundation, Saira Sheikh: None declared, Goran Radunovic: None declared, XIAOBI HUANG Shareholder of: Biogen, Employee of: Biogen, HUA CARROLL Shareholder of: Biogen, Employee of: Biogen, Francois Gaudreault Shareholder of: Biogen, Employee of: Biogen, Adam Meyers Shareholder of: Biogen, Employee of: Biogen, Catherine Barbey Shareholder of: Biogen, Employee of: Biogen, Cristina Musselli Shareholder of: Biogen, Employee of: Biogen, NATHALIE FRANCHI-MONT Shareholder of: Biogen, Employee of: Biogen
DOI: 10.1136/annrheumdis-2020-eular.5743

OP0194 IMMUNOSUPPRESSIVE EFFECT OF TOLERGENIC DENDRITIC CELLS ON AUTOLOGOUS T-CELLS IN PATIENTS WITH RHEUMATOID ARTHRITIS

Y. Kurochkina¹, T. Tyrinova², O. Leplina², M. Tikhonova², A. Sizikov², O. Chumasova², A. Ostanin², E. Chernykh². ¹Research Institute of Clinical and Experimental Lymphology - Branch of the Institute of Cytology and Genetics, Siberian Branch of the Russian Academy of Science, Novosibirsk, Russian Federation; ²Federal State Budgetary Scientific Institution Research Institute of Fundamental and Clinical Immunology, Novosibirsk, Russian Federation

Background: Dendritic cells (DCs) are known to contribute to the pathogenesis of rheumatoid arthritis (RA) through presentation of cartilage glycoprotein, production of proinflammatory cytokines and activation of Th1/Th17 responses. Along with stimulating activity, DCs may exhibit suppressive functions via capacity to induce T cell apoptosis/anergy and to generate regulatory T cells. Since these DCs have potential to control autoreactive T-lymphocytes, the enhancing of tolerogenic properties of DCs seems to be a new important strategy in treatment of RA. However, it remains unclear whether autoreactive T lymphocytes are sensitive to the immunosuppressive effects of DC.

Objectives: The aim of our study is to investigate the mechanisms of the inhibitory effect of dexamethasone-modified DCs (dexDCs) in patients with RA on autologous T cells.

Methods: Twenty patients with RA with high and moderate activity of disease were recruited in this study. All patients fullfield ACR/EULAR criteria (2010). All studies were performed after receiving informed consent. DCs were generated from blood monocytes culturing for 5 days with GM-CSF and IFN- α in the presence dexamethasone, applied on third day. LPS as maturation stimuli was added on fourth day. The study evaluated the ability of dexDCs to induce T cell apoptosis, inhibit the production of Th1 and Th17 cytokines in auto-mixed leukocytes culture (MLC), induce the activation of T-regulatory cells, and inhibit the purified protein derivative (PPD)-specific immune response.

Results: We revealed that dexDCs markedly suppressed the production of IFN- γ and IL-17, while the decreasing of IL-4 and IL-13 was less pronounced. Thus, in comparison with Th2 cells, Th1 and Th17 T lymphocytes were more sensitive to suppressor effects of dexDCs. We also revealed a significant increasing of cells in the stage of late apoptosis (An+Pi+) during the cultivation of autologous T cells with dexDCs. We have also proved the suppressive effect of dexDCs on culture T cells that are co-cultivated with mature DCs (reduction of proliferation from 2700 cpm to 1500 cpm, $p=0.0015$). In addition, we have shown the ability of dexDCs to induce T-regulatory cells with a phenotype CD4+IL10+Tr1. In conclusion, we have shown the ability of PPD-loaded dexDCs to inhibit the proliferative response of

both mature DCS and mature dexDCs-loaded PPD, that indicates the ability of dexDCs to possess antigen-specific suppression.

Conclusion: The data obtained indicate that, dexDCs from RA patients have an immunosuppressive effect on autologous T cells through the induction of apoptosis, anergia and activation of T regulatory cells that authorise their application as a DCs-vaccine.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.1291

Creating in vitro patients - how to best model disease

OP0195 VASCULARIZED THREE-DIMENSIONAL MODELS OF HUMAN SKIN FIBROSIS

A.E. Matei¹, C.W. Chen¹, L. Kiesewetter², A.H. Györfi¹, Y.N. Li¹, T. Trinh-Minh¹, T. Van Kuppevelt³, J. Hansmann^{2,4}, A. Juengel⁵, G. Schett¹, F. Groeber-Becker^{2,6}, J. Distler¹. ¹Department of Internal Medicine 3 – Rheumatology and Immunology, Friedrich-Alexander-University Erlangen-Nürnberg (FAU) and University Hospital Erlangen, Erlangen, Germany; ²Translational Center Würzburg, Fraunhofer Translational Center Regenerative Therapies, Fraunhofer Institute for Silicate Research (ISC), Würzburg, Germany; ³Radboud Institute for Molecular Life Sciences, Department of Biochemistry, Radboud University Medical Center, Nijmegen, Netherlands; ⁴University for Applied Sciences Würzburg-Schweinfurt, Würzburg, Germany; ⁵Center of Experimental Rheumatology, University Hospital Zurich/Zurich Center of Integrative Human Physiology (ZIHP), Zürich, Switzerland; ⁶Department for Tissue Engineering and Regenerative Medicine, Würzburg University Medical Center, Würzburg, Germany

Background: The complex pathophysiological processes that result in fibrotic tissue remodeling in systemic sclerosis involve interplay between multiple cell types (1). Experimental models of fibrosis are essential to provide a conceptual understanding of the pathogenesis of these diseases and to test antifibrotic drugs. Current models of fibrosis have important limitations: the *in vivo* models rely on species that are phylogenetically distant, whereas the *in vitro* models are oversimplified cultures of a single cell type in an artificial two-dimensional environment of excessive stiffness, which imposes an unphysiological cell polarization (2).

Objectives: Here we evaluated the potential use of vascularized, three-dimensional *in vitro* human skin equivalents as a novel model of skin fibrosis and a platform for the evaluation of antifibrotic drugs.

Methods: Skin equivalents were generated by seeding human endothelial cells, fibroblasts and keratinocytes on a decellularized porcine extracellular matrix with perfusable vascular structure. The skin models were cultured for one month in a system that ensured perfusion of the vascular network at physiological pressure. Fibrotic transformation induced by TGF β and response to nintedanib as an established antifibrotic drug was evaluated by capillary Western immunoassays, qPCR, histology and immunostaining.

Results: The vascularized human skin equivalents formed the major skin structures relevant for the pathogenesis of fibrosis: a polarized, fully matured epidermis, a stratified dermis and a perfused vessel system with small capillaries. Exposure to TGF β led to the fibrotic transformation of the skin equivalents, with activated TGF β downstream pathways, increased fibroblast-to-myofibroblast transition and excessive deposition of extracellular matrix. Treatment of models exposed to TGF β with nintedanib (a drug with proven antifibrotic effects) ameliorated the fibrotic transformation of skin equivalents with reduced TGF β signaling, fibroblast-to-myofibroblast transition and decreased extracellular matrix deposition.