(patients who stop treatment no longer contribute information, which may overestimate true drug effectiveness).

Objectives: To present points to consider (PtC) when analysing and reporting comparative effectiveness with observational data in rheumatology (EULAR-funded taskforce).

Methods: The task force comprises 18 experts: epidemiologists, statisticians, rheumatologists, patients, and health professionals.

Results: A systematic literature review of methods currently used for comparative effectiveness research in rheumatology and a statistical simulation study were used to inform the PtC (table). Overarching principles focused on defining treatment effectiveness and promoting robust and transparent epidemiological and statistical methods increase the trustworthiness of the results.

Points to consider

Reporting of comparative effectiveness observational studies must follow the STROBE guidelines

Authors should prepare a statistical analysis plan in advance

To provide a more complete picture of effectiveness, several outcomes across multiple health domains should be compared

Lost to follow-up from the study sample must be reported by the exposure of interest

The proportion of patients who stop and/or change therapy over time, as well as the reasons for treatment discontinuation must be reported

Covariates should be chosen based on subject matter knowledge and model selection should be justified

The study baseline should be at treatment initiation and a description of how covariate measurements relate to baseline should be included

The analysis should be based on all patients starting a treatment and not limited to patients remaining on treatment at a certain time point

When treatment discontinuation occurs before the time of outcome assessment, this attrition should be taken into account in the analysis.

Sensitivity analyses should be undertaken to explore the influence of assumptions related to missingness, particularly in case of attrition

Conclusion: The increased use of real-world comparative effectiveness studies makes it imperative to reduce divergent or contradictory results due to biases. Having clear recommendations for the analysis and reporting of these studies should promote agreement of observational studies, and improve studies' trustworthiness, which may also facilitate meta-analysis of observational data.

Disclosure of Interests: Delphine Courvoisier: None declared, Kim Lauper: None declared, Sytske Anne Bergstra: None declared, Maarten de Wit Grant/research support from: Dr. de Wit reports personal fees from Ely Lilly, 2019, personal fees from Celgene, 2019, personal fees from Pfizer, 2019, personal fees from Janssen-Cilag, 2017, outside the submitted work., Consultant of: Dr. de Wit reports personal fees from Ely Lilly, 2019, personal fees from Celgene, 2019, personal fees from Pfizer, 2019, personal fees from Janssen-Cilag, 2017, outside the submitted work., Speakers bureau: Dr. de Wit reports personal fees from Ely Lilly, 2019, personal fees from Celgene, 2019, personal fees from Pfizer, 2019, personal fees from Janssen-Cilag, 2017, outside the submitted work., Bruno Fautrel Grant/research support from: AbbVie, Lilly, MSD, Pfizer, Consultant of: AbbVie, Biogen, BMS, Boehringer Ingelheim, Celgene, Lilly, Janssen, Medac MSD France, Nordic Pharma, Novartis, Pfizer, Roche, Sanofi Aventis, SOBI and UCB, Thomas Frisell: None declared, Kimme Hyrich Grant/research support from: Pfizer, UCB, BMS, Speakers bureau: Abbvie, Florenzo Iannone Consultant of: Speaker and consulting fees from AbbVie, Eli Lilly, Novartis, Pfizer, Roche, Sanofi, UCB, MSD, Speakers bureau: Speaker and consulting fees from AbbVie, Eli Lilly, Novartis, Pfizer, Roche, Sanofi, UCB, MSD, Joanna KEDRA: None declared, Pedro M Machado Consultant of: PMM: Abbvie, Celgene, Janssen, Lilly, MSD, Novartis, Pfizer, Roche and UCB, Speakers bureau: PMM: Abbvie, BMS, Lilly, MSD, Novartis, Pfizer, Roche and UCB, Lykke Midtbøll Ørnberg Grant/research support from: Novartis, Ziga Rotar Consultant of: Speaker and consulting fees from Abbvie, Amgen, Biogen, Eli Lilly, Medis, MSD, Novartis, Pfizer, Roche, Sanofi., Speakers bureau: Speaker and consulting fees from Abbvie, Amgen, Biogen, Eli Lilly, Medis, MSD, Novartis, Pfizer, Roche, Sanofi., Maria Jose Santos Speakers bureau: Novartis and Pfizer, Tanja Stamm Grant/research support from: AbbVie, Roche, Consultant of: AbbVie, Sanofi Genzyme, Speakers bureau: AbbVie, Roche, Sanofi, Simon Stones Consultant of: I have been a paid consultant for Envision Pharma Group and Parexel. This does not relate to this abstract., Speakers bureau: I have been a paid speaker for Actelion and Janssen. These do not relate to this abstract., Anja Strangfeld Speakers bureau: AbbVie, BMS, Pfizer, Roche, Sanofi-Aventis, Robert B.M. Landewé Consultant of: AbbVie; AstraZeneca; Bristol-Myers Squibb; Eli Lilly & Co.; Galapagos NV; Novartis; Pfizer; UCB Pharma, Axel Finckh Grant/research support from: Pfizer: Unrestricted research grant, Eli-Lilly: Unrestricted research grant, Consultant of:

Sanofi, AB2BIO, Abbvie, Pfizer, MSD, Speakers bureau: Sanofi, Pfizer, Roche, Thermo Fisher Scientific

DOI: 10.1136/annrheumdis-2020-eular.1162

Multiple issues in inherited connective tissues - more than 'just' hypermobility

OP0200

PATIENTS WITH HYPERMOBILITY RELATED DISORDERS HAVE A SIGNIFICANT NUMBER OF ORTHOPAEDIC INTERVENTIONS ON MULTIPLE SITES AND AT A YOUNG AGE: DATA FROM A TERTIARY REFERRAL CENTRE

S. Wright¹, P. Mehta², J. Parry², H. Kazkaz². ¹University of Dundee, Dundee, United Kingdom; ²University College London, Rheumatology, London, United Kingdom

Background: Mechanisms of pain associated with joint hypermobility are poorly understood and include nociceptive pain from structural joint changes along with soft tissue injuries linked to impaired proprioception; central sensitisation associated with chronic pain and muscle weakness alongside deconditioning. Anxiety and depression are also thought to play a role in patients presenting with pain and hypermobility. We have observed an increase in the rate of orthopaedic surgical procedures undertaken in patients attending the hypermobility clinics compared to those attending the general rheumatology and chronic pain clinics. There is limited published data regarding orthopaedic interventions in patients with hypermobility related disorders especially those with confirmed genetic mutations.

Objectives: We aimed to evaluate the characteristics of patients in our hypermobility cohort focusing on those who had received prior surgical intervention in order to understand the underlying mechanism behind their presentations.

Methods: A retrospective review of medical records was conducted of patients attending a hypermobility clinic at our tertiary referral centre, University College London Hospital, between January 2018 and December 2018.

Results: There were 350 patients (300 females, 50 males) with a mean age of 36 years (range 18-71 years). 63% had a diagnosis of Hypermobility Spectrum Disorder or Hypermobility Syndrome and 37% had a type of Ehlers-Danlos Syndromes (EDS) (hypermobility, classical, vascular or other rare type). 46 patients (13%) had documented genetic mutations. 83 patients (24%) had undergone orthopaedic interventions including 9 who had EDS with confirmed genetic mutations. 54% of patients who had surgical intervention were under the age of 40. The total number of surgical procedures in the cohort was 227 (equating to 0.6485 interventions per patient). Of those requiring operative intervention, the average number of interventions per patient was 2.73. One third of patients had surgery on two or more joint groups, including 8 patients (2%) who had surgery in four or more joint groups. Knees (24%) and hips (23%) were the most common sites for operative intervention with 9% having surgery on their shoulders. 29% of pts had significant hypermobility with a Beighton score of 7 and above but there was no correlation between Beighton score and number of surgical procedures. Only 2% of cases were referred from an orthopaedic team thereby excluding a referral bias.

Conclusion: Patients with hypermobility related disorders have a significant number of orthopaedic surgical procedures on multiple sites and at a young age, with indication of mechanical pathology playing an important role in their symptoms. The Beighton score does not appear to be a reliable predictor of surgical intervention. This is not surprising given that the score only covers 5 joint areas and excludes common surgical sites such as the hips and shoulders. Early diagnosis and a holistic non-operative approach combining physiotherapy and chronic pain management is essential to reduce the need for multiple surgical procedures.

References:

- [1] Chopra P, Tinkle B, Hamonet C, Brock I, Gompel A, Bulbena A, et al. Pain management in the Ehlers-Danlos syndromes. *Am J Med Genet C Semin Med Genet* [Internet]. 2017 [cited 2020 Jan 27];175(1):212–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28186390>
- [2] Shirley ED, Demajo M, Bodurtha J. Ehlers-danlos syndrome in orthopaedics: etiology, diagnosis, and treatment implications. *Sports Health* [Internet]. 2012 Sep [cited 2019 Jan 30];4(5):394–403. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23016112>

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.1486