Objectives: Our aim was to analyze the association of HLA class II with HS in a Caucasian population from Cantabria (northern Spain).

Methods: In this study we analyzed the *HLA-A, -B, -C, DRB1, -DQA1 and -DQB1* allele distribution in 106 HS patients and 262 age- and sex-matched controls from a Caucasian population of Cantabria (northern Spain).

Results: *HLA-A*29* and *B*50* were significantly more frequent in HS patients and *A*30* and *B*37* in controls, but these associations disappeared after correction. On the other hand, *DRB1*07*, *DQA1*02* and *DQB1*02* were significantly more frequent in controls (p 0.026, p 0.0012 and p 0.0005 respectively), and the *HLA* allele *DQB1*03:01* was significantly more frequent in HS patients (p 0.00007) all of them after Bonferroni correction. Furthermore, the *DRB1*07*; *DQA1*02*; *DQB1*02* haplotype was significantly more frequent in controls (p 0.0005).

Conclusion: This is the first study showing an association of HLA-class II with HS. Our results suggest that HLA-II alleles (*DRB1*07*, *DQA1*02*, *DQB1*02* and *DQB1*03:01*) and the *DRB1*07*-*DQA1*02*-*DQB1*02* haplotype could influence on resistance or susceptibility to HS.

References:

- [1] González-López MA. J Am Acad Dermatol. 2016; Aug;75(2):329-35.
- [2] González-López MA. PLoS One. 2018 Jan 4;13(1).
- [3] Vilanova I. J Eur Acad Dermatol Venereol. 2018 May;32(5):820-824.
- [4] Durán-Vian C, et al. J Eur Acad Dermatol Venereol. 2019 Nov;33(11):2131-2136.

Disclosure of Interests: Monica Calderón-Goercke: None declared, J. Gonzalo Ocejó-Vinyals: None declared, Miguel A González-Gay Grant/research support from: Pfizer, Abbvie, MSD, Speakers bureau: Pfizer, Abbvie, MSD, Marcelo A. Fernández-Viña: None declared, Juan Cantos-Mansilla: None declared, Iosune Vilanova: None declared, Ricardo Blanco Grant/research support from: AbbVie, MSD, and Roche, Speakers bureau: AbbVie, Pfizer, Roche, Bristol-Myers, Janssen, and MSD, Marcos González-López: None declared

DOI: 10.1136/annrheumdis-2020-eular.4483

AB0006

HYPOMETHYLATION OF THE PROMOTER REGION OF TLR4 GENE AT A SYSTEMIC LEVEL IN PATIENTS WITH RHEUMATOID ARTHRITIS AND PERIODONTITIS

S. Chatterjee¹, D. Bhattacharjee¹, S. Misra¹, A. Rao¹, A. Chattopadhyay², A. Ghosh¹, ¹Institute of Post Graduate Medical Education and Research, Department of Rheumatology, Kolkata, India; ²Post Graduate Institute of Medical Education & Research, Department of Rheumatology, Chandigarh, India

Background: Periodontitis (PD) has long been linked with Rheumatoid arthritis (RA) [1]. Epigenetic modifications are being recently explored to explain such associations, DNA methylation being one such important mechanism.

Objectives: To study the effect chronic generalized periodontitis on systemic methylation of TLR4 genes in comparison to only RA and RA with PD patients.

Methods: Twenty-three RA patients, among which 11 patients had chronic generalized PD, 20 patients with only PD and 15 healthy individuals recruited. DNA was isolated from PBMCs of the participants blood, then were first bisulphite converted and then methylation specific PCR were performed using primers for methylated and un-methylated promoters of TLR4. The DNA amplifications were checked in horizontal gel electrophoresis. The methylation signatures were verified by DNA sequencing (Sanger) of the amplified products.

Results: The anti-CCP, DAS-CRP and HAQ DI were higher in patients with both RA and PD (220±40, 5.7±0.2, 1.5±0.1 respectively, p <0.05). Control samples had shown amplification bands for methylated primers of TLR4 but not for un-methylated primers of TLR4. However, only RA, only PD and RA with PD samples, had shown amplification for un-methylated primers and not for methylated primers. These results together with DNA sequencing indicated that 4 CpG sites in the promoter of TLR4 genes were hypo-methylated in the PBMCs of patients whereas those remain methylated in healthy individuals.

Conclusion: The observations indicated that though PD is a localised disease of the gingiva there is a systemic involvement of TLR mediated pathways in them which is similar to those in RA. However, further validation in larger cohort and down-stream signalling molecules needs to be studied.

References:

- [1] Ceccarelli F, Saccucci M, Di Carlo G, Lucchetti R, Pilloni A, Pranno N, Luzzi V, Valesini G, Polimeni A. Periodontitis and rheumatoid arthritis: the same inflammatory mediators? Mediators of inflammation. 2019;2019.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.3455

AB0007

VALUE OF ULTRASOUND IN ASSESSMENT OF ACTIVE SACROILIITIS IN PATIENTS WITH AXIAL SPONDYLOARTHRITIS

M. Chen¹, S. M. Dai¹. ¹Shanghai Jiao Tong University Affiliated Sixth People's Hospital, Department of Rheumatology and Immunology, Shanghai, China

Background: The inflammatory of the sacroiliac joints (SIJs) called sacroiliitis, is a characteristic of axial Spondyloarthritis (axSpA). The detection of sacroiliitis is meaningful to prevent irreversible changes. The tool of assessment of sacroiliitis including radiographs, computed tomography (CT) and magnetic resonance imaging (MRI). Ultrasound (US) has also been used in the evaluation of sacroiliitis in recent years.

Objectives: We aimed to evaluate the value of US in the assessment of active sacroiliitis in axSpA patients.

Methods: Fifty-one patients fulfilling Assessment of SpondyloArthritis International Society (ASAS) 2009 criteria for the classification of axSpA were recruited¹. All the patients underwent MRI and US evaluation of bilateral SIJs. MRI was performed using the sequences of T1WI, T2WI and fat suppression T2WI (FS-T2WI). MRI sacroiliitis was defined according to ASAS criteria of active sacroiliitis². The Spondyloarthritis research Consortium of Canada (SPARCC) scoring was used to evaluate the inflammatory lesions in SIJs³. US were performed by an ultrasonographer with 10 years of experience in musculoskeletal ultrasound, and resistive index (RI) value was recorded. The US sacroiliitis was defined as the presence of more flow signals at SIJ with an RI $\leq$ 0.75. The HLA-B27, erythrocyte sedimentation rate (ESR) and hypersensitive C-reactive protein (hsCRP) were also evaluated. Consistency rate, sensitivity, specificity, positive predictive values (PPV) and negative predictive values (NPV) for the diagnosis of sacroiliitis by US were calculated, using MRI as the gold standard.

Results: Of the 51 patients, 24 were female and 27 were male. The HLA-B27 positive rate was 90.2% (46/51). The consistency rate of US and MRI sacroiliitis was 55.88 (57/102). The sensitivity and specificity of US for the diagnosis of sacroiliitis were 55.93 (33/59) and 55.81 (24/43) respectively. The PPV and NPV were 63.46 (33/52) and 48 (24/50) respectively. There was no significant difference in ESR and hsCRP between the US positive sacroiliitis and the others ($P=0.7477$ and 0.2268 , respectively). The SPARCC scores have no significant difference between the US positive sacroiliitis and the others ($P=0.2206$). The RI was not significantly associated with the MRI SPARCC score ($P=0.4236$).

Conclusion: US may be an optional method for preliminary screening sacroiliitis. But its reliability as a diagnostic method needs further verification.

References:

- [1] Rudwaleit M, et al. The development of Assessment of SpondyloArthritis international Society classification criteria for axial spondyloarthritis (part II): validation and final selection. Ann Rheum Dis. 2009; 68(6):777-83
- [2] Rudwaleit M, et al. Defining active sacroiliitis on magnetic resonance imaging (MRI) for classification of axial spondyloarthritis: a consensus approach by the ASAS/OMERACT MRI group. Ann Rheum Dis 2009;68(10):1520-7
- [3] Maksymowych WP, et al. Spondyloarthritis research Consortium of Canada magnetic resonance imaging index for assessment of sacroiliac joint inflammation in ankylosing spondylitis. Arthritis Rheum. 2005;53(5):703-9.

Acknowledgments: This project was supported by grants from National Natural Science Foundation of China (81900795)

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.3990

AB0008

THE IMPACT OF STAT4 RS7574865, IL6 RS1800795, IL6R RS2228145 AND RS4845618 ON RHEUMATOID ARTHRITIS SUSCEPTIBILITY IN BELARUSIAN POPULATION

E. Siniauskaya¹, T. Kuzhir¹, V. Yagur², R. Goncharova¹. ¹Institute of Genetics and Cytology of NAS of Belarus, Minsk, Belarus; ²Belarusian State Medical University, Minsk, Belarus

Background: Rheumatoid arthritis (RA) is a chronic systemic disorder of the connective tissue of still unknown aetiology and complex autoimmune pathogenesis that primarily affects small joints. HLA alleles provide for 11-37% of the RA heritability, suggesting the substantial role of the non-HLA loci in genetic predisposition to RA. Among non-HLA loci, *IL6*, *IL6R* and *STAT4* genes attract attention, however, the data concerning their influence on RA risk are somewhat contradictory.

Objectives: The aim of the study was to analyze the involvement of four SNPs (*STAT4* rs7574865, *IL6* rs1800795, *IL6R* rs2228145 and rs4845618) in RA susceptibility.

Methods: 187 patients diagnosed with RA (mean age 58.2 ± 11.9), and 380 healthy blood donors (mean age 37.18 ± 10.69 years) were included into the