Background: STING (stimulator of interferon genes) is a cytosolic protein that is found in endoplasmic reticulum (ER) membrane, mitochondria and mitochondria-associated membranes. Although it is well established that STING plays an important role in innate immune responses, its potential involvement in rheumatic disease processes remains to be clarified (1).

Objectives: The aims of this study were to evaluate the levels of STING and its relationship with local inflammation in the synovial fluid (SF) of patients with psoriatic arthritis (PsA), rheumatoid arthritis (RA), gout, calcium pyrophosphate (CPP) crystal-induced arthritis (CPP-IA), osteoarthritis (OA) and OA with CPP crystals (OA+CPP).

Methods: SF was collected from the knees of 60 untreated patients: 10 with PsA, 10 with RA, 10 with gout, 10 with CPP-IA, 10 with OA and 10 with OA+CPP. SF was examined under optical light microscopy. White cell count (WBC) and the polymorphonuclear cell (PMN) percentage were determined in SF according to standard procedures. SF IL-8, IL-6, IL-1 β and extra- and intra-cellular STING levels were assayed by ELISA.

Results: The levels of WBC were higher in SFs from gouty patients ($27.7 \pm 20.56 \times 10^3/\text{mm}^3$), while OA and OA+CPP patients showed the lowest WBC count ($0.34 \pm 0.3 \times 10^3/\text{mm}^3$, $0.3 \pm 0.28 \times 10^3/\text{mm}^3$). SFs from inflammatory arthritis contained elevated percentages of PMN (gout: $85.5 \pm 10.86\%$, CPP-IA: $84 \pm 11.31\%$, RA: $80.33 \pm 8.14\%$, PsA: $42.6 \pm 35.97\%$). Extracellular STING was determined in OA ($440 \pm 413.31 \text{ pg/ml}$), OA+CPP ($225 \pm 205.06 \text{ pg/ml}$) and CPP-IA ($475 \pm 7.07 \text{ pg/ml}$) SF, while was not detectable in RA, PsA and gout. Intracellular STING levels were similar and the highest in SFs from gout ($96.4 \pm 35.44 \text{ pg/ml}$) and RA ($90.64 \pm 23.13 \text{ pg/ml}$), while remained under detection limit only in SFs from PsA. SF concentration of IL-6 was lower in OA ($354.87 \pm 377.56 \text{ pg/ml}$) and OA+CPP ($389.56 \pm 104.14 \text{ pg/ml}$) as compared with inflammatory arthritides (PsA: $3807.14 \pm 489.86 \text{ pg/ml}$; RA: $17354 \pm 2334.87 \text{ pg/ml}$; gout: $19359 \pm 84.85 \text{ pg/ml}$; CPP-IA: $20389.56 \pm 104.14 \text{ pg/ml}$). The patients with gout and RA had the highest levels of IL-8 ($2159.54 \pm 347.09 \text{ pg/ml}$; $2039.6 \pm 97.74 \text{ pg/ml}$) and IL-1 β ($35.93 \pm 20.46 \text{ pg/ml}$; $44.36 \pm 23.16 \text{ pg/ml}$), while OA showed the lowest concentrations (IL-8: $23.21 \pm 11.32 \text{ pg/ml}$; IL-1 β : $0.47 \pm 0.13 \text{ pg/ml}$). In the total group of patients, we found a negative correlation between extracellular STING and IL-8 ($r = -0.53$; $p = 0.004$) and IL-1 β ($r = -0.47$; $p = 0.012$). There was a positive correlation between intracellular STING and IL-8 ($r = 0.54$; $p = 0.017$), IL-1 β ($r = 0.77$; $p < 0.001$) and IL-6 ($r = 0.69$; $p = 0.009$).

Conclusion: This study is the first to define the presence of STING in SF of different arthritides. The high levels of extracellular STING in OA, OA+CPP and CPP-IA SFs may be due to the activation of factors that reduce its interaction with the ER. The effect of downregulating factors in PsA might explain the low concentration of intracellular STING in these patients.

References:

- [1] Barber GN. STING: infection, inflammation and cancer. *Nat Rev Immunol* 2015;15:760-70.

Disclosure of Interests: Anna Scanu: None declared, Roberto Luisetto: None declared, Francesca Oliviero: None declared, Paola Galozzi: None declared, Augusta Ortolan: None declared, Mariagrazia Lorenzin: None declared, Mara Felicetti: None declared, Andrea Doria Consultant of: GSK, Pfizer, Abbvie, Novartis, Ely Lilly, Speakers bureau: UCB pharma, GSK, Pfizer, Janssen, Abbvie, Novartis, Ely Lilly, BMS, Roberta Ramonda Speakers bureau: Novartis, Celgene, Janssen, Pfizer, Abbvie, Lilly

DOI: 10.1136/annrheumdis-2020-eular.5734

AB0061 INVOLVEMENT OF CELL-FREE DNA IN THE DIFFERENTIATION OF OSTEOCLASTS – IMPLICATIONS FOR CLINICAL RESEARCH

P. Škubica¹, J. Horinkova², M. Gregová², K. Pavelka², M. Husakova², P. Dankova¹. ¹Faculty of Science, Charles University, Department of Anthropology and Human Genetics, Prague 2, Czech Republic; ²Institute of Rheumatology and Department of Rheumatology, First Faculty of Medicine, Charles University, Prague 2, Czech Republic

Background: Cell-free DNA (cfDNA) is a molecule with undisputable immunogenic potential^[1]. Its role is well established in etiopathogenesis of systemic lupus erythematosus and many reports suggest it may play an important role in other rheumatic diseases^[2]. Patients suffering from rheumatic diseases are at higher risk of osteoporosis^[3], which may be a consequence of increased generation of osteoclasts (OC). Traditionally, increased formation of OC is attributed to changes in cytokine network^[4]. It is not yet known whether cfDNA can also act on mononuclear precursors and influence their differentiation into OC.

Objectives: The aim of this study was to assess role of cfDNA in differentiation of bone resorbing cells in rheumatic diseases. For this purpose, deoxyribonuclease I (DNase) was employed to eliminate cfDNA from sera of patients and healthy controls (HC) used for cell culture.

Methods: In order to evaluate effects of cfDNA on osteoclastogenesis, an *in vitro* experiment of peripheral blood monocytes (PBM) cultures was designed. Axial spondyloarthritis (AxSpA) was chosen as a model of rheumatic disease. Sera of

AxSpA patients (total n=13) and healthy controls (total n=13) were collected and subsequently used in cell cultures. PBM of 13 healthy donors were cultured in presence of 25 ng/ml macrophage colony-stimulating factor (M-CSF), 0.5 ng/ml receptor-activator of nuclear factor kappa B ligand (RANKL) and 10 % of pooled AxSpA or HC sera, respectively. Serum counterpart treated with DNase was used in parallel. After 14 days, cells were stained for tartrate-resistant acid phosphatase (TRAP) and counted. TRAP positive cells with 3 or more nuclei were recognized as OC.

Results: On average, differentiation of OC in presence of AxSpA sera led to higher osteoclastogenesis compared to DNase-treated counterpart (137 vs. 61; $p < 0.01$). In a similar way, treatment of HC sera with DNase greatly diminished osteoclastogenesis compared to untreated sera (17 vs. 111; $p < 0.05$). Overall, the effect of cfDNA depletion in HC sera regarding OC differentiation was more than 2.5x higher than the effect of cfDNA depletion in AxSpA sera ($p < 0.001$).

Conclusion: Results of our pilot study indicate possible involvement of cfDNA in osteoclastogenesis. Decrease in number of OC after cfDNA depletion in both AxSpA sera and HC sera suggests that cfDNA effect in osteoclastogenesis is non-specific with respect to healthy condition and may represent universal link between innate immunity and bone metabolism. Lower effect of cfDNA depletion in AxSpA sera on osteoclastogenesis is very possibly owing to presence of other inflammatory molecules, which influence osteoclastogenesis. Future research might identify cfDNA as useful therapeutic target to decrease osteoporosis risk not only in patients with rheumatic and autoimmune diseases, where pathological bone resorption is imminent, but also in general aging population.

References:

- [1] Pisetsky, D. S. (2012). The origin and properties of extracellular DNA: from PAMP to DAMP. *Clinical Immunology*, 144(1), 32-40.
- [2] Lood, C. J., & Duvvuri, B. (2019). Cell-free DNA as a biomarker in autoimmune rheumatic diseases. *Frontiers in immunology*, 10, 502.
- [3] Maruotti, N., Corrado, A., & Cantatore, F. P. (2014). Osteoporosis and rheumatic diseases. *Reumatismo*, 125-135.
- [4] Amarasekara, D. S., Yu, J., & Rho, J. (2015). Bone loss triggered by the cytokine network in inflammatory autoimmune diseases. *Journal of immunology research*, 2015.

Acknowledgments: Project MH CR 00023728 & MEYS CR Progres Q43

Disclosure of Interests: Patrik Škubica: None declared, Jana Horinkova: None declared, Monika Gregová Consultant of: Novartis, Abbvie, Paid instructor for: Novartis, Speakers bureau: Novartis, Abbvie, MSD, Karel Pavelka Consultant of: Abbvie, MSD, BMS, Egis, Roche, UCB, Medac, Pfizer, Biogen, Speakers bureau: Abbvie, MSD, BMS, Egis, Roche, UCB, Medac, Pfizer, Biogen, Marketa Husakova Speakers bureau: Novartis, Pavlina Dankova: None declared

DOI: 10.1136/annrheumdis-2020-eular.2774

AB0062 CHARACTERIZATION OF IL-12 AND IL-23 REDUCTION BY TOFACITINIB IN MDCs

N. Vincken¹, C. Angioliilli¹, S. Cardoso¹, A. Lopes¹, M. Olde-Nordkamp¹, T. Radstake¹. ¹Center of Translational Immunology, Rheumatology and Clinical Immunology, University Medical Center Utrecht, Utrecht, Netherlands

Background: Psoriatic arthritis (PsA) is a chronic inflammatory auto-immune disease characterized by an excessive production of pathogenic mediators that cause inflammation of the skin, peripheral joints, entheses and the spine. Among these, interleukin (IL)-23, IL-12, the IL-17 family and TNF constitute key players in PsA pathogenesis.^{1,2} IL-23, consisting of IL23A (IL-23p19) and IL23B (IL-23p40) subunits, is predominantly produced by myeloid dendritic cells (mDCs). While the p19 subunit is unique to IL-23, the p40 subunit is shared with IL-12. Together, IL-12 and IL-23 play a crucial role in promoting the differentiation of naïve T lymphocytes into T helper (Th) interferon (IFN)- γ -producing Th1 or IL17-producing Th17 cells, respectively.³

Small-molecule inhibitors, such as the JAK/STAT inhibitor Tofacitinib, have recently shown promising therapeutic potential in PsA clinical trials.⁴ The inhibition of JAKs by Tofacitinib results in the direct suppression of multiple intracellular signaling pathways which constitute key hubs in the cytokine network.⁵ However, whether Tofacitinib is able directly target IL-12/IL-23 production by mDCs has not yet been documented. Suppression of these canonical inflammatory pathways would provide further evidence that Tofacitinib is an effective drug in halting both innate and adaptive immune responses.

Objectives: To evaluate the transcriptional and molecular events underlining IL-12 and IL-23 regulation by Tofacitinib in mDCs.

Methods: Peripheral blood mononuclear cells from healthy donors were isolated by Ficoll gradient. Monocytes and myeloid dendritic cells (mDCs) were isolated by using magnetic beads on autoMACS. Monocytes were cultured for 6 days in the presence of IL-4 and granulocyte-macrophage colony-stimulating factor (GM-CSF) to generate monocyte-derived dendritic cells (moDCs). MoDCs were harvested, washed and put to rest for 1 day prior to stimulation, while mDCs

were stimulated on the same day of isolation. Both moDCs and mDCs were pre-treated with Tofacitinib and then stimulated with either lipopolysaccharide (LPS) or combination of LPS with IFN- γ for 4 hours. Cytokines were measured using enzyme-linked immunosorbent assay (ELISA) and gene expression was assessed using quantitative polymerase chain reaction (qPCR).

Results: Treatment of both mDCs and moDCs with Tofacitinib led to a decreased mRNA expression of IL-12 p40 (*IL12B*) in the presence of TLR4 and IFN γ co-stimulation. The decreased *IL12B* mRNA expression also resulted in lower production of IL-12 p40 and IL-23 proteins in mDCs.

Conclusion: In this work, we demonstrated for the first time that Tofacitinib can suppress the production of IL-23/IL-12 p40 subunit in mDCs, upon the condition that an active type II IFN signalling is also present in these cells. This observation indicates that specific factors, such as endogenous IFN- γ levels in the serum of PsA patients, can possibly predict differential responses to Tofacitinib treatment.

References:

- [1] Gaffen SL. et al. The IL-23-IL-17 immune axis: from mechanisms to therapeutic testing. *Nat Rev Immunol.* 2014 Sep;14(9):585-600
- [2] Bravo A, Kavanaugh A. Bedside to bench: defining the immunopathogenesis of psoriatic arthritis. *Nat Rev Rheumatol.* 2019 Nov;15(11):645-656
- [3] Floss DM. et al. Insights into IL-23 biology: From structure to function. *Cytokine Growth Factor Rev.* 2015 Oct;26(5):569-78
- [4] Berekmeri A. et al. Tofacitinib for the treatment of psoriasis and psoriatic arthritis. *Expert Rev Clin Immunol.* 2018 Sep;14(9):719-730
- [5] T Virtanen E. et al. Selective JAKinibs: Prospects in Inflammatory and Auto-immune Diseases. *BioDrugs.* 2019 Feb;33(1):15-32

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.3307

OA, aetiology, pathology and animal models

AB0063

AGING CARTILAGE IN WILD-TYPE MICE: AN OBSERVATIONAL STUDY

J. Akoum¹, K. Tahiri¹, F. Etienne¹, M. T. Corvol¹, F. Rannou^{1,2}, C. Nguyen^{1,2}.

¹Université de Paris - UFR Mathématiques et Informatique, Paris, France;

²Hospital Cochin, Rheumatology Institute, Rehabilitation Department, Paris, France

Background: Many animal models of osteoarthritis (OA) have been used to study the pathogenesis of cartilage degeneration¹. In mice, spontaneous OA can occur in wild-type or genetically modified animals. The first report of spontaneous OA developing in wild-type mice was published in 1956² and changes affecting the knee joint were further related to OA by using ultrastructural-histochemical analyses. However, a quantitative assessment of age-related evolution of OA-type cartilage lesions is lacking. The OA Research Society International (OARSI) grading score was adapted to semi-quantify histopathological changes occurring in OA animal models, including mice³. The OARSI score has been used to describe changes occurring in induced or genetic OA mouse models but not to describe spontaneous age-related evolution of OA-type cartilage lesions in wild-type mice.

Objectives: We aimed to describe the spontaneous evolution of age-related changes affecting knee joint articular cartilage, walking speed and a serum biomarker of cartilage remodeling in C57BL/6 wild-type male mice.

Methods: Histological changes were assessed by the OARSI score in newborn, 1-week- and 1-, 3-, 6-, 9- and 12-month-old C57BL/6 wild-type male mice, walking speed by the Locotronic system, and serum C-terminal telopeptide of type II collagen (CTX-II) content by ELISA in 1-, 3-, 6-, and 9-month-old C57BL/6 wild-type male mice.

Results: Mean (SD) OARSI score increased from 0.2(0.3) to 1.3(0.6) ($p=0.03$) between 1 and 3 months of age and from 1.3(0.6) to 3.3(0.6) ($p=0.04$) between 3 and 6 months of age. Mean walking speed was stable between 1 and 6 months of age but decreased from 11.4(1.8) to 3.2(0.8) cm.s⁻¹ ($p=0.03$) between 6 and 9 months of age. Serum CTX-II content was maximal at 1 month of age, then decreased from 12.2(8.5) to 2.4(8.4) pg/ml ($p=0.02$) between 1 and 3 months of age, remaining low and stable thereafter.

Conclusion: C57BL/6 wild-type male mice showed continuously increasing osteoarthritic changes but delayed decreasing walking speed with age. These variations were maximal between 3 and 9 months of age. Maximal serum CTX-II content preceded these changes.

References:

- [1] McCoy AM. Animal Models of Osteoarthritis: Comparisons and Key Considerations. *Vet Pathol.* 2015;52(5):803-18.
- [2] Sokoloff L. Natural history of degenerative joint disease in small laboratory animals. I. Pathological anatomy of degenerative joint disease in mice. *AMA Arch Pathol.* 1956;62(2):118-28.

- [3] Glasson SS. The OARSI histopathology initiative - recommendations for histological assessments of osteoarthritis in the mouse. *Osteoarthritis Cartilage.* 2010;18 Suppl 3:S17-23.

Table 1. Evolution of cartilage changes, walking speed and serum C-terminal telopeptide of type II collagen (CTX-II) concentrations in wild-type C57BL/6 male mice.

Age	New-born	1 week	1 month	3 months	6 months	9 months	12 months
OARSI score (0 to 6)	0.0 (0.0)	0.0 (0.0)	0.2 (0.3)	1.3 (0.6)*	3.3 (0.6)*	3.7 (0.6)	4.3 (0.6)
Walking speed (cm.s⁻¹)	-	-	10.5 (1.5)	11.3 (4.3)	11.4 (1.8)	3.2 (0.8)*	-
CTX-II concentrations (pg/ml)	-	-	12.2 (8.5)	2.4 (8.4)*	1.1 (4.0)	4.0 (3.8)	-

N ≥ 3 per timepoint. All results are means (standard deviation). * $p<0.05$ as compared to the previous timepoint using the non-parametric Mann-Whitney U-test.

Figure 1. (A) New born, 1-week- and 1-, 3-, 6-, 9- and 12-month-old wild-type C57BL/6 male mice (N=3 per timepoint) knees were evaluated for cartilage changes following OARSI recommended guidelines by 3 independent readers. Each point represents the mean of OARSI score per mice as scored by 3 independent readers. **(B)** 1-, 3-, 6- and 9-month-old wild-type C57BL/6 male mice were evaluated for walking speed using the Locotronic® system. Each point represents the mean of 3 measures of walking speed per mice. All results are means (SD). * $p<0.05$ as compared to the previous timepoint using the non-parametric Mann-Whitney U-test.

Disclosure of Interests: Joulmar Akoum: None declared, Khadija Tahiri: None declared, François Etienne: None declared, Marie-Thérèse Corvol: None declared, François Rannou Grant/research support from: Pierre Fabre, Fidia, MSD, Pfizer, Bone Therapeutics, Expanscience, Grunenthal, Thuasne, Genévrier, Fondation Arthritis, Consultant of: Pierre Fabre, Fidia, MSD, Pfizer, Bone Therapeutics, Expanscience, Grunenthal, Thuasne, Genévrier, Speakers bureau: Pierre Fabre, Fidia, MSD, Pfizer, Bone Therapeutics, Expanscience, Grunenthal, Thuasne, Christelle Nguyen: None declared

DOI: 10.1136/annrheumdis-2020-eular.5989

AB0064

CHANGES IN THE VASCULAR ENDOTHELIAL GROWTH FACTOR A (VEGFA) SPLICING AXIS IN HUMAN SYNOVIUM ARE RELATED TO INFLAMMATION IN ARTHRITIS

D. Amanitis^{1,2}, S. Shahtaheri^{2,3}, D. Daniel McWilliams^{2,3}, N. Beazley-Long¹, D. Walsh^{2,3}, L. Donaldson^{1,2}. ¹University of Nottingham, School of Life Sciences, Nottingham, United Kingdom; ²University of Nottingham, Pain Centre Versus Arthritis, Nottingham, United Kingdom; ³University of Nottingham, School of Medicine, Academic Rheumatology, Nottingham, United Kingdom

Background: VEGF-A is a key regulator of rheumatoid (RA) and osteoarthritis (OA). During articular inflammation in OA and RA there is increased synovial angiogenesis and upregulation of angiogenic growth factors such as VEGF-A. VEGF-A comprises two splice variant families, VEGF-A_{xxx}a and VEGF-A_{xxx}b (xxx represents the number of amino acids, from 121 to 206), resulting from alternative splice site selection in exon 8. This site selection is controlled by Serine/Arginine Rich Splicing Factor Kinase 1 (SRPK1), which phosphorylates Serine/Arginine Rich Splicing Factor 1 (SRSF1), inducing it to translocate to the nucleus. In most normal tissues, VEGF-A_{xxx}b isoforms predominate, with anti-nociceptive and anti-angiogenic functions. In contrast, in pathological conditions such as inflammation and solid tumours, VEGF-A_{xxx}a isoforms predominate, with pro-nociceptive and pro-angiogenic functions. VEGF-A has been proposed as