Figure 1. Effects of LOR on the quantity of the cartilage catabolism end products glycosaminoglycan (GAG) and nitric oxide (NO) in supernatants. Knee cartilage explant cultures stimulated with pro-inflammatory cytokines were subsequently treated with DMSO (control) or LOR as shown. N=22; * $p<0.05$, ** $p<0.01$, *** $p<0.001$ vs. DMSO by one-way ANOVA.

Disclosure of Interests: Vishal Deshmukh Shareholder of: Samumed, LLC, Employee of: Samumed, LLC, Shawn Grogan: None declared, Tim Seo Shareholder of: Samumed, LLC, Employee of: Samumed, LLC, Deepti Bhat Shareholder of: Samumed, LLC, Employee of: Samumed, LLC, William Bugbee: None declared, Darryl D'Lima: None declared, Yusuf Yazici Shareholder of: Samumed, LLC, Grant/research support from: Bristol-Myers Squibb, Celgene, and Genentech, Consultant of: Celgene and Sanofi, Employee of: Samumed, LLC
DOI: 10.1136/annrheumdis-2020-eular.6346

AB0071

THERAPEUTIC EFFECTS OF BONE MARROW MESENCHYMAL STEM CELLS-DERIVED EXOSOMES ON OSTEOARTHRITIS

C. Dong¹, Y. Liu¹, A. Deng¹, J. Ji¹, W. Zheng¹, Z. Gu¹. ¹Affiliated Hospital of Nantong University, Nantong, China

Background: Mesenchymal stem cells (MSCs) have shown chondroprotective effects in clinical models of osteoarthritis (OA)^[1].

Objectives: The study aimed to investigate the therapeutic potential of exosomes from human bone marrow MSCs (BM-MSCs) in alleviating OA.

Methods: The anterior cruciate ligament transection (ACLT) and destabilization of the medial meniscus (DMM) surgery were performed on the knee joints of a rat OA model, followed by intra-articular injection of BM-MSCs or their exosomes. The beneficial effects were evaluated by histological staining, OARSI scores and micro-CT. Furthermore, BM-MSCs-derived exosomes were administered to primary human chondrocytes to observe the functional and molecular alterations. In addition, lncRNA MEG3 was investigated in chondrocytes to explore the biological contents accounting for anti-OA effects of BM-MSCs-derived exosomes.

Results: Based on the observation in the rat OA model, both of BM-MSCs and BM-MSCs-derived exosomes alleviated cartilage destruction, reduced joint damage and restored the trabecular bone of OA rats. In addition, *in vitro* assays showed that BM-MSCs- exosomes could maintain the chondrocyte phenotype by increasing collagen type II synthesis and inhibiting IL-1 β - induced senescence and apoptosis. Furthermore, exosomal lncRNA MEG3 also reduced the senescence and apoptosis of chondrocytes induced by IL-1 β , indicating that lncRNA MEG3 might partially account for the anti-OA effects of BM-MSC exosomes.

Conclusion: The exosomes from BM-MSCs exerted beneficial therapeutic effects on OA by reducing the senescence and apoptosis of chondrocytes, suggesting that MSCs-derived exosomes might provide a candidate therapy for OA treatment.

References:

- [1] McKinney J M, Doan T N, Wang L, et al. Therapeutic efficacy of intra-articular delivery of encapsulated human mesenchymal stem cells on early stage osteoarthritis[J]. Eur Cell Mater, 2019, 37: 42-59.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.6040

AB0072

A MULTICOMPONENT MEDICATION PROMOTES CHONDROGENESIS AND REDUCES MMP-13 IN PRIMARY ARTICULAR CHONDROCYTES FROM KNEE OSTEOARTHRITIS PATIENTS IN VITRO

C. Sanchez¹, K. Hemmer², N. Kroemmelbein², B. Seilheimer², J. E. Dubuc³, C. Antoine⁴, Y. Henrotin^{1,4}. ¹University of Liège, Bone and Cartilage Research Unit, Liège, Belgium; ²Biologische Heilmittel Heel GmbH, Baden-Baden, Germany; ³Cliniques Universitaires Saint-Luc, Orthopedic Department, Brussels, Belgium; ⁴Artialis, Liège, Belgium

Background: HE-1100 is a multicomponent medicinal product. Initial preclinical data potentially suggest a preventive effect on cartilage degradation.

Objectives: This study aims to understand the mode of action of HE-1100 on OA chondrocytes *in vitro*.

Methods: Primary chondrocytes were obtained from 10 knee osteoarthritis (OA) patients undergoing knee replacement surgery. The cultures were treated with 20% (v/v) HE-1100 or placebo. Samples were collected for subsequent RNA extraction using standard methods. The reads were generated with Illumina NextSeq5000 sequencer and aligned to the human reference genome (UCSC hg19) to generate the transcriptome. Differential expression analysis between HE-1100 and placebo was made in R using the DESeq2 package to identify the differentially expressed genes in the OA-associated regulatory pathways. The protein production of the selected genes was quantified by ELISA in 10 independent human OA chondrocytes cultures.

Results: According to the DESeq2 analysis, HE-1100 significantly modified the expression of 13 genes in OA chondrocytes by at least 10% with an adjusted p-value < 0.05: EGR1 (+93%), FOS (+87%), NR4A1 (+43%), DUSP1 (+18%), ZFP36 (+18%), ZFP36L1 (+14%), NFKB1 (+16%) and CYR61 (+14%) were upregulated and ATF7IP (-10%), TXNIP (-11%), C10orf10 (-12%), CLEC3A (-12%) and MMP13 (-18%) were downregulated after 24h HE-1100 treatment. HE-1100 significantly increased (2.3 fold $\pm$ 1.2 after 24h, $p=0.0444$ and 2.3-fold $\pm$ 1.0 after 72h, $p=0.0239$) the CYR61 protein production by human OA chondrocytes. After 72h, HE-1100 slightly but not significantly increased aggrecan production by $14 \pm 19\%$ ($p=0.1117$) and significantly increased type II collagen pro-peptide production by $27 \pm 20\%$ ($p=0.0147$). For both time points CYR61 production by OA chondrocytes was positively and significantly correlated with aggrecan ($r=0.66$, $p=0.0004$) and type II collagen pro-peptide ($r=0.64$, $p=0.0008$) production. In alginate beads culture, pro-MMP-13 was significantly decreased by HE-1100 treated cultures from day 7 to day 14 (from -16 to -25 %, $p<0.05$) and from day 17 to 21 (-22 %, $p=0.0331$) in comparison to controls.

Conclusion: HE-1100 significantly modified the expression of DUSP1, C10orf10, ZFP36/L1 and CLEC3A, which are pathway mediators involved in MMP-13 expression and activation. Further, long-term (28 days) treatment with HE-1100 significantly reduced the production of pro-MMP-13, the inactive precursor of the metalloproteinase MMP-13 involved in type II collagen degradation. HE-1100 also promoted extracellular matrix formation probably through CYR61 production, a growth factor well correlated with type II collagen and aggrecan production.

References:

Acknowledgments: We would like to thank the staff of the GIGA ULiège Genomic Next Generation Sequencing platform for performing the RNA sequencing and Benoit Charleatoux for his help in RNAseq data analysis.

Disclosure of Interests: christelle sanchez: None declared, Kathrin Hemmer Employee of: Heel, Natascha Kroemmelbein Employee of: Heel, Bernd Seilheimer Employee of: Heel, Jean-Emile Dubuc: None declared, Christophe Antoine Employee of: Artialis, Yves Henrotin Grant/research support from: HEEL, TILMAN

DOI: 10.1136/annrheumdis-2020-eular.2106

AB0073

BOSWELLIA SERRATA EXTRACT AND CURCUMIN INCREASE GDF15 PRODUCTION BY HUMAN PRIMARY OSTEOARTHRITIS CHONDROCYTES: A NEW MECHANISM OF ACTION

C. Sanchez¹, J. Zappia¹, Y. Dierckxsens², J. P. Delcour³, Y. Henrotin^{1,4}.

¹University of Liège, Bone and Cartilage Research Unit, Liège, Belgium;

²Tilman, Baillonville, Belgium; ³Centre Hospitalier du Bois de l'Abbaye, Seraing, Belgium; ⁴Princess Paola Hospital, Marche-en-Famenne, Belgium

Background: Boswellia serrata extract (BSE) and curcumin are used to relief symptoms in osteoarthritis (OA).

Objectives: This study aims to better understand the mode of action of these compounds on OA chondrocytes *in vitro*.

Methods: Therapeutic plasmatic concentrations of the different components of BSE correspond to an *in vitro* range from 25 to 100 μ g/ml of total BSE (100 μ g/ml of BSE corresponds to 9.2 μ M of 11-keto- β -boswellic acid (KBA), 5.2 μ M of acetylKBA, 22 μ M de α BA, 34 μ M de β BA, 4.4 μ M de acetyl α BA and 13.2 acetyl β BA), and between 2 to 10 μ M for bioavailability-increased curcumin. BSE (5-100 μ g/ml) and curcumin (0.04 to 4 μ g/ml corresponding to 0.1 to 10 μ M) were tested separately on primary chondrocytes from 3 OA patients. Lactate Dehydrogenase LDH, nitrite (NO₂), interleukin (IL)-6 and Growth Differentiation Factor (GDF)15 were quantified in 72h-treated supernatant using enzyme activity, Griess reaction and ELISAs, respectively.

Results: No mortality was observed at the tested concentrations. BSE and curcumin both decreased concentration-dependently NO₂ and IL-6 production, and increased GDF15 production. For NO₂ production, the decrease was observed from 0.2 μ g/ml of curcumin and 10 μ g/ml of BSE. For IL-6 production,