- The belief that HCQ is "Very important" (12.9% "low") rather than "important" (22.1% "low") or "not important / useless" (33.1% "low").
 - Taking many different medications (9.8% "low" for Patients indicating more than 7 medications vs 19.8% for those listing 3 or less)
 Childhood onset of the SLE was associated with a lower adherence (30.0% "low" vs. 17.4% for later onset SLE ($p < .001$))

658 patients (29.6%) reported having experienced side effects. 42.6% of them stopped taking HCQ (patient led 161, doctor led 110, unclear 9). Amongst those continuing HCQ despite experiencing side effects, the proportion of non adherent patients increased to 24.7%, compared to 15.2% in the group of patients that have not experienced side effects ($p = .0001$). The 232 patients who talked with their Doctor and felt listened to appear to adhere better (22.0% low adherence) than the 84 who did not feel heard (31.0% low adherence), but the significance is only directional ($p < .015$)

523 patients have used HCQ in the past. 206 (39.4%) consider the decision to stop HCQ as doctor initiated, 272 (52.0%) as patient initiated, and 36 (6.9%) as a joint decision.

When stopping was patient initiated, 59.9% was due to experiencing a significant side effect attributed by the patient to HCQ, 6.7% due to concern of a potential side effect, 11.2% "tested" stopping and noticed no difference, 10.0% were not convinced that it worked, 8.2% felt their lupus was less active, 2.6% wanted to reduce pill consumption. (Side effects attributed to HCQ may relate to age, disease activity or other factors).

Conclusion: Doctors can help HCQ adherence by boosting patient's confidence in the importance of HCQ. Better patient education may contribute to avoid up to 40% of patient initiated decision to stop HCQ treatment.

References:

- [1] Fanouriakis A et al. 2019 update of the EULAR recommendations for the management of systemic lupus erythematosus. *Ann Rheum Dis*. 2019 Jun; 78(6): 736-745.
- [2] Costedoat N et al, Treatment adherence in systemic lupus erythematosus and rheumatoid arthritis: time to focus on this important issue, *Rheumatology*, Vol 57 (9), Sep 2018, 1507–1509.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.3233

AB0372

PREGNANCY OUTCOME IN WOMEN WITH SJÖGREN SYNDROME IS OFTEN COMPLICATED BY PLACENTA INSUFFICIENCY

I. Troester¹, F. Kollert², A. Zbinden², L. Raio³, F. Foerger². ¹University Hospital and University of Bern, Inselspital, Pediatrics, Bern, Switzerland; ²University Hospital and University of Bern, Inselspital, Rheumatology, Immunology and Allergology, Bern, Switzerland; ³University Hospital and University of Bern, Inselspital, Obstetrics, Bern, Switzerland

Background: Chronic inflammatory rheumatic diseases are often associated with a negative effect on pregnancy outcome. Most obstetrical complications are placenta-mediated such as preterm delivery and growths restrictions. In women with Sjögren syndrome, data on placenta-mediated complications are scarce and conflicting (1,2).

Objectives: To analyse neonatal outcome in women with Sjögren syndrome with focus on preterm delivery and growth restriction.

Methods: We retrospectively analysed 23 pregnancies of 16 patients with Sjögren syndrome that were followed at our centre with regard to pregnancy outcome, medication and disease characteristics. Small for gestational age was defined as birthweight percentile $< 10^{\text{th}}$. Preterm delivery was defined as delivery before 37, early term as delivery between 37-39 and term as delivery between 39-42 weeks of gestation.

Results: Of 23 pregnancies, one ended in a miscarriage and 22 resulted in live births including one set of twins. Treatment used during pregnancy was hydroxychloroquine (20 pregnancies), prednisone (8), azathioprine (5) and cyclosporine (2). Concomitant treatment with low-dose aspirin was used in 9 pregnancies.

Of the 22 live births, 17 were born at early term and 5 at term. There were no preterm deliveries. Median birth weight was 2820g (range 2095-3845g). Nine newborns (40.9%) were small for gestational age (SGA). Maternal treatment during these pregnancies was hydroxychloroquine in all cases and additional low-dose aspirin in three cases. Elevated CRP levels during pregnancy were found in 57% of the cases with SGA outcome. Only one woman with an SGA infant had positive anti-phospholipid antibodies.

Regarding delivery mode, most patients had caesarean sections.

Conclusion: In our cohort of women with Sjögren syndrome the prevalence of small for gestational age infants was high despite maternal treatment with hydroxychloroquine. Inflammatory markers could help to identify the patients

at risk for placental insufficiency, yet prospective studies of larger cohorts are needed.

References:

- [1] Gupta S et al; Sjögren Syndrome and Pregnancy: A literature review. *Perm J* 2017; 21:16-047
- [2] De Carolis S et al; The impact of primary Sjögren's syndrome on pregnancy outcome: Our series and review of the literature. *Autoimmun Rev* 2014; 13(2):103-7

Disclosure of Interests: Isabella Troester: None declared, Florian Kollert Employee of: Novartis, Astrid Zbinden: None declared, Luigi Raio: None declared, Frauke Foerger Grant/research support from: unrestricted grant from UCB, Consultant of: UCB, GSK, Roche, Speakers bureau: UCB, GSK

DOI: 10.1136/annrheumdis-2020-eular.5725

AB0373

TREATMENT OF SLE WITH THE IMMUNOPROTEASOME INHIBITOR KZR-616: RESULTS FROM THE FIRST 4 COHORTS OF THE MISSION STUDY, AN OPEN-LABEL PHASE 1B DOSE ESCALATION TRIAL

R. Furie¹, S. Parikh², A. Maiquez³, A. Khan⁴, O. Moreno³, M. Soneira⁴, C. Kirk⁵, D. Bombardieri⁶, K. Harvey⁵, M. K. Farmer⁵. ¹Northwell Health, Great Neck, NY, United States of America; ²The Ohio State University Wexner Medical Center, Columbus, United States of America; ³SouthCoast Research Center, Inc, Miami, United States of America; ⁴Accurate Clinical Research, Houston, United States of America; ⁵Kezar Life Sciences, South San Francisco, United States of America

Background: Non-specific proteasome inhibitors, such as bortezomib (BTZ), target the constitutive proteasome and immunoproteasome and are approved treatments for multiple myeloma¹. BTZ has also been used to treat systemic lupus erythematosus (SLE) and lupus nephritis (LN); however, treatment emergent adverse events (TEAEs), such as gastrointestinal (GI) effects, hematologic abnormalities, asthenia and peripheral neuropathy, limit its use as a long-term treatment option for chronic autoimmune disease². KZR-616 is a first-in-class selective immunoproteasome inhibitor and is highly active in murine SLE³. Subcutaneous (SC) administration of KZR-616 (30 and 45 mg weekly [QW]) was demonstrated as safe and well-tolerated, and successfully achieved target levels of immunoproteasome inhibition in healthy volunteers^{4,5}.

Objectives: We report the preliminary safety and efficacy of KZR-616 in the first 4 cohorts of the Phase 1b portion of Study KZR-616-002 in patients with active SLE (NCT03393013).

Methods: SLE patients (per SLICC Classification Criteria) with SLEDAI ≥ 4 despite stable background immunosuppressant, anti-malarial, and/or corticosteroid (≤ 20 mg prednisone equivalent) therapy in this open-label multicentric dose-escalation trial received KZR-616 at doses of 45mg (Cohort 1), 60mg (Cohort 2), or 30mg with escalation to 60mg (Cohorts 2a and 2b) SC weekly through Week 13 (W13) with 12 weeks of follow-up. Efficacy measures included SLEDAI, Cutaneous Lupus Erythematosus Disease Area and Severity Index, 28 tender and swollen joint counts, Physicians Global Assessment, Patient Global Assessment, and Patient Assessment of Pain, in evaluable patients (those who received ≥ 1 month of KZR-616).

Results: As of 16 January 2020, 33 patients had enrolled and received at least 1 dose of KZR-616. The majority of TEAEs have been mild or moderate with no reported peripheral neuropathy, prolonged GI-related AEs, and no clinically significant laboratory AEs. There were 3 treatment emergent SAEs, one each of thrombotic microangiopathy (Cohort 2), localized herpes zoster (Cohort 2a), and systemic inflammatory response syndrome (Cohort 2a) with the latter 2 patients completing the full 13 weeks of treatment after resolution. When compared to baseline, improvement in all measures of disease activity were seen at W13 and maintained or improved during the follow-up period, and 94% of evaluable patients had improvements on at least 2 measures/assessments of disease activity. A single patient with active class IV/V nephritis was enrolled on prednisone 10 mg, leflunomide 10 mg, and hydroxychloroquine 200mg/day; nephrotic-range proteinuria at baseline (3.85 g/day) decreased to 0.6 g/day 4 weeks after the last dose of KZR-616.

Conclusion: Weekly subcutaneous administration of KZR-616 at 45 and 60 mg was safe and well-tolerated. Evidence of disease suppression at W13 in active SLE patients on stable background therapy was observed. In addition, one study participant with active proliferative nephritis was enrolled with significant reduction in proteinuria. The Phase 2 portion of this study in active proliferative LN is open for enrollment.

References:

- [1] Scott K. et al. *Cochrane Database Syst Rev* 2016;4:CD010816