Figure. Cumulative mortality of patients with idiopathic inflammatory myopathy and matched controls without IIM in early versus late cohorts (1999-2006 versus 2007-2014).

Disclosure of Interests: Kristin D'Silva: None declared, Leo Lu: None declared, Alexis Ogdie Grant/research support from: Pfizer to Penn, Novartis to Penn, Amgen to Forward/NDB, Consultant of: Abbvie, Amgen, Bristol-Myers Squibb, Celgene, Corrona, Janssen, Eli Lilly, Novartis, Pfizer, Antonio Aviña: None declared, Hyon Choi Grant/research support from: Ironwood, Horizon, Consultant of: Takeda, Selecta, Horizon, Kowa, Vaxart, Ironwood
DOI: 10.1136/annrheumdis-2020-eular.2230

OP0248 PREMATURE MORTALITY BURDEN FOR SYSTEMIC SCLEROSIS: NATIONWIDE POPULATION-BASED STUDY

E. Yen¹, D. Singh¹, M. Wu¹, B. Singh¹. ¹UCLA, Los Angeles, United States of America

Background: Premature mortality is an important way to quantify disease burden. Patients with systemic sclerosis (SSc) can die prematurely of disease, however, the premature mortality burden of SSc is unknown. The years of potential life lost (YPLL), in addition to age-standardized mortality rate (ASMR) in younger ages, can be used as measures of premature death.

Objectives: To evaluate the premature mortality burden of SSc by calculating: 1) the proportions of SSc deaths as compared to deaths from all other causes (non-SSc) by age groups over time, 2) ASMR for SSc relative to non-SSc-ASMR by age groups over time, and 3) the YPLL for SSc relative to other autoimmune diseases.

Methods: This is a population-based study using a national mortality database of all United States residents from 1968 through 2015, with SSc recorded as the underlying cause of death in 46,798 deaths. First, we calculated the proportions of deaths for SSc and non-SSc by age groups for each of 48 years and performed joinpoint regression trend analysis¹ to estimate annual percent change (APC) and average APC (AAPC) in the proportion of deaths by age. Second, we calculated ASMR for SSc and non-SSc causes and ratio of SSc-ASMR to non-SSc-ASMR by age groups for each of 48 years, and performed joinpoint analysis to estimate APC and AAPC for these measures (SSc-ASMR, non-SSc-ASMR, and SSc-ASMR/non-SSc-ASMR ratio) by age. Third, to calculate YPLL, each decedent's age at death from a specific disease was subtracted from an arbitrary age limit of 75 years for years 2000 to 2015. The years of life lost were then added together to yield the total YPLL for each of 13 preselected autoimmune diseases.

Results: 23.4% of all SSc deaths as compared to 13.5% of non-SSc deaths occurred at <45 years age in 1968 ($p < 0.001$, Chi-square test). In this age group, the proportion of annual deaths decreased more for SSc than for non-SSc causes: from 23.4% in 1968 to 5.7% in 2015 at an AAPC of -2.2% (95% CI, -2.4% to -2.0%) for SSc, and from 13.5% to 6.9% at an AAPC of -1.5% (95% CI, -1.9% to -1.1%) for non-SSc. Thus, in 2015, the proportion of SSc and non-SSc deaths at <45 year age was no longer significantly different. Consistently, SSc-ASMR decreased from 1.0 (95% CI, 0.8 to 1.2) in 1968 to 0.4 (95% CI, 0.3 to 0.5) per million persons in 2015, a cumulative decrease of 60% at an AAPC of -1.9% (95% CI, -2.5% to -1.2%) in <45 years old. The ratio of SSc-ASMR to non-SSc-ASMR also decreased in this age group (cumulative -20%, AAPC -0.3%). In <45 years old, the YPLL for SSc was 65.2 thousand years as compared to 43.2 thousand years for rheumatoid arthritis, 18.1 thousand years for dermatomyositis, 146.8 thousand years for myocarditis, and 241 thousand years for type 1 diabetes.

Conclusion: Mortality at younger ages (<45 years) has decreased at a higher pace for SSc than from all other causes in the United States over a 48-year period. However, SSc accounted for more years of potential life lost than

rheumatoid arthritis and dermatomyositis combined. These data warrant further studies on SSc disease burden, which can be used to develop and prioritize public health programs, assess performance of changes in treatment, identify high-risk populations, and set research priorities and funding.

References:

[1] Yen EY....Singh RR. *Ann Int Med* 2017;167:777-785.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.2270

OP0249 LONG-TERM EXTENSION RESULTS OF RISE-SSC, A RANDOMIZED TRIAL OF RIOCIQUAT IN PATIENTS WITH EARLY DIFFUSE CUTANEOUS SYSTEMIC SCLEROSIS (DCSSC)

D. Khanna¹, J. Pope², M. Matucci-Cerinic³, M. Kuwana⁴, C. Denton⁵, Y. Allanore⁶, M. Wosnitzer⁷, M. E. Truchetet⁸, G. Szűcs⁹, W. Stevens¹⁰, V. Steen¹¹, C. Stagnaro¹², V. Smith¹³, R. Silver¹⁴, E. Schiopu¹⁵, V. Riccieri¹⁶, F. Kramer⁷, S. Johnson¹⁷, O. Ishikawa¹⁸, T. Ishii¹⁹, E. Hachilla²⁰, E. De Langhe²¹, L. Czirják²², R. Bečvář²³, T. Atsumi²⁴, O. Distler²⁵. ¹Univ Michigan, Ann Arbor, United States of America; ²Univ Western Ontario, Ontario, Canada; ³Univ Florence, Florence, Italy; ⁴Nippon Med School, Tokyo, Japan; ⁵UCL, London, United Kingdom; ⁶Paris Descartes Univ, Paris, France; ⁷Bayer AG, Wuppertal, Germany; ⁸Bordeaux Univ Hosp, Bordeaux, France; ⁹Univ Debrecen, Debrecen, Hungary; ¹⁰St Vincent's Hosp, Melbourne, Australia; ¹¹Georgetown Univ, Washington, United States of America; ¹²Univ Pisa, Pisa, Italy; ¹³Ghent Univ Hosp, Ghent, Belgium; ¹⁴Medical Univ South Carolina, Charleston, United States of America; ¹⁵Michigan Medicine Univ Hosp, Ann Arbor, United States of America; ¹⁶Sapienza Univ Rome, Rome, Italy; ¹⁷Univ Toronto, Ontario, Canada; ¹⁸Gunma Univ, Maebashi, Japan; ¹⁹Tohoku Univ, Sendai, Japan; ²⁰Hôpital Claude Huriez, CHU, Lille Univ, Lille, France; ²¹Univ Hosp Leuven, Leuven, Belgium; ²²Univ Pécs Medical School, Pécs, Hungary; ²³Charles Univ, Prague, Czech Republic; ²⁴Hokkaido Univ, Sapporo, Japan; ²⁵Univ Hosp Zurich, Zurich, Switzerland

Background: RISE-SSc (NCT02283762) was a multicenter Phase IIb trial of riociguat in pts with early (duration ≤ 18 months) dcSSc and modified Rodnan skin score (mRSS) 10–22 units. Pts were randomized double-blind to placebo or riociguat 0.5–2.5 mg t.i.d. for 52 weeks. The primary endpoint, mRSS change from baseline to Week (Wk) 52, did not reach statistical significance ($p = 0.08$, riociguat vs placebo), but there were favorable trends in some other outcomes.

Objectives: To present open-label long-term extension (LTE) results of RISE-SSc.

Methods: Pts who completed Wk 52 of double-blind therapy could enter LTE on riociguat. Endpoints included mRSS, adverse events (AEs), and serious AEs (SAEs).

Results: Of 60 pts randomized to riociguat and 61 to placebo, 42 (riociguat-riociguat group) and 45 (former placebo group), respectively, entered LTE. At LTE start, mean $\pm$ SD mRSS was 16.4 $\pm$ 3.2 and 16.3 $\pm$ 4.2 units, and mean disease duration was 8.9 $\pm$ 7.8 and 8.9 $\pm$ 5.8 months, in the riociguat-riociguat and former placebo groups, respectively. Other demographics/disease characteristics were also comparable. Median duration of riociguat treatment was 1092 d in riociguat-riociguat pts and 649 d in former placebo pts. Throughout the study, mRSS decreased in both groups (Figure 1). From Wk 52 to last visit, mRSS fell by -3.02 ± 5.51 in riociguat-riociguat patients and -3.96 ± 5.43 in former placebo pts. Rates of mRSS regression (decrease by >5 units and $\geq 25\%$ from Wk 52 to last visit) and of % declines in mRSS were similar in the two groups (Figure 2). mRSS progression (increase by >5 units and $\geq 25\%$ from Wk 52 to last visit) occurred in 1 pt (2%) in each group. During the entire study, rescue therapy agents were used in 15 (36%) riociguat-riociguat pts and 17 (38%) former placebo pts. AEs were reported from Wk 52 to last visit in 82 pts (94%): 40 (95%) riociguat-riociguat and 42 (93%) former placebo. Most common AEs overall: nasopharyngitis (24%), gastroesophageal reflux disease (17%), diarrhea (15%), and hypotension (14%). AEs of special interest (dizziness, postural dizziness, or hypotension) occurred in 5 riociguat-riociguat pts (12%) and 4 former placebo pts (9%). SAEs were reported in 21 (24%) pts: 10 (24%) riociguat-riociguat pts and 11 (24%) former placebo pts, with no SAE reported in >1 patient, no SAEs of special interest, and no deaths.

Conclusion: During LTE riociguat treatment, mRSS decreased in both groups from Wk 52 onwards and mRSS progression was uncommon. Riociguat had acceptable safety, similar to the main study, with no new safety signal.

Figure 1. Mean change from baseline in mRSS by time point from baseline to the last visit (long-term safety analysis set)

BL, baseline; mRSS, modified Rodnan skin score

Figure 2. Changes in mRSS from Week 52 to last visit

^aDecrease by ≥5 units and ≥25% from Week 52 to last visit
mRSS, modified Rodnan skin score

Acknowledgments: RISE-SSc was jointly funded by Bayer AG and Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Kenilworth, NJ, USA.

Disclosure of Interests: Dinesh Khanna Shareholder of: Eicos, Grant/research support from: NIH NIAID, NIH NIAMS, Consultant of: Acceleron, Actelion, Bayer, BMS, Boehringer Ingelheim, Corbus, Galapagos, Genentech/Roche, GSK, Mitsubishi Tanabe, Sanofi-Aventis/Genzyme, UCB Pharma, Janet Pope Grant/research support from: AbbVie, Bristol-Myers Squibb, Eli Lilly & Company, Merck, Roche, Seattle Genetics, UCB, Consultant of: AbbVie, Actelion, Amgen, Bayer, Boehringer Ingelheim, Bristol-Myers Squibb, Eicos Sciences, Eli Lilly & Company, Emerald, Gilead Sciences, Inc., Janssen, Merck, Novartis, Pfizer, Roche, Sandoz, Sanofi, UCB, Speakers bureau: UCB, Marco Matucci-Cerinic Grant/research support from: Actelion, MSD, Bristol-Myers Squibb, Speakers bureau: Actelion, Lilly, Boehringer Ingelheim, Masataka Kuwana Grant/research support from: Actelion, Consultant of: Actelion, Bayer, Chugai, Corbus Pharmaceuticals, CSL Behring and Reata Pharmaceuticals. He was a member of the SENSICIS trial Steering Committee (Boehringer Ingelheim), Christopher Denton Grant/research support from: GlaxoSmithKline, CSL Behring, and Inventiva, Consultant of: Medscape, Roche-Genentech, Actelion, GlaxoSmithKline, Sanofi Aventis, Inventiva, CSL Behring, Boehringer Ingelheim, Corbus Pharmaceuticals, Acceleron, Curzion and Bayer, Yannick Allanore Grant/research support from: BMS, Inventiva, Roche, Sanofi, Consultant of: Actelion, Bayer AG, BMS, BI, Melanie Wosnitza Employee of: Bayer AG, Marie-Elise Truchetet: None declared, Gabriella Szűcs: None declared, Wendy Stevens: None declared, Virginia Steen Grant/research support from: The associated affiliation has received grants/research from Boehringer Ingelheim, Corbus Pharmaceuticals, CSL Behring, Eicos, Galapagos, Immune Tolerance Network, Reata, Consultant of: Virginia Steen has acted as a consultant for Boehringer Ingelheim, Corbus, CSL Behring, Eicos, Forbuis, Chiara Stagnaro: None declared, Vanessa Smith Grant/research support from: The affiliated company received grants from Research Foundation - Flanders (FWO), Belgian Fund for Scientific Research in Rheumatic diseases (FWRO), Boehringer Ingelheim Pharma GmbH & Co and Janssen-Cilag NV, Consultant of: Boehringer-Ingelheim Pharma GmbH & Co, Speakers bureau: Actelion Pharmaceuticals Ltd, Boehringer-Ingelheim Pharma GmbH & Co and UCB Biopharma Sprl, Richard Silver: None declared, Elena Schiopu: None declared, Valeria Ricciari: None declared, Frank Kramer Employee of: Bayer AG, Sindhu Johnson Grant/research support from: Boehringer Ingelheim, Corbus Pharmaceuticals, GlaxoSmithKline, Roche, Merck, Bayer, Consultant of: Boehringer Ingelheim, Ikaria, Osamu Ishikawa: None declared, Tomonori Ishii: None declared, Eric Hachilla: None declared, Ellen De Langhe Consultant of: member of advisory board for Boehringer, László Cziráj Consultant of: Actelion, BI, Roche-Genentech, Lilly, Medac, Novartis, Pfizer, Bayer AG, Radim Bečvář Consultant of: Actelion, Roche, Tatsuya Atsumi Grant/research support from: Eli Lilly Japan K.K., Alexion Pharmaceuticals, Inc., Bristol-Myers Squibb Co., AbbVie Inc., Daiichi Sankyo Co., Ltd., Pfizer Inc., Chugai Pharmaceutical Co., Ltd., Mitsubishi Tanabe Pharma Co., Astellas Pharma Inc., Consultant of:

Gilead Sciences, Inc., Eli Lilly Japan K.K., UCB Japan Co. Ltd., AbbVie Inc., Daiichi Sankyo Co., Ltd., Pfizer Inc., Chugai Pharmaceutical Co., Ltd., Speakers bureau: Eli Lilly Japan K.K., UCB Japan Co. Ltd., Bristol-Myers Squibb Co., AbbVie Inc., Eisai Co. Ltd., Otsuka Pharmaceutical Co., Ltd., Daiichi Sankyo Co., Ltd., Pfizer Inc., Chugai Pharmaceutical Co., Ltd., Mitsubishi Tanabe Pharma Co., Takeda Pharmaceutical Co., Ltd., Astellas Pharma Inc., Oliver Distler Grant/research support from: Grants/Research support from Actelion, Bayer, Boehringer Ingelheim, Competitive Drug Development International Ltd. and Mitsubishi Tanabe; he also holds the issued Patent on mir-29 for the treatment of systemic sclerosis (US8247389, EP2331143), Consultant of: Consultancy fees from Actelion, Acceleron Pharma, AnaMar, Bayer, Baecon Discovery, Blade Therapeutics, Boehringer, CSL Behring, Catenion, ChemomAb, Curzion Pharmaceuticals, Ergonex, Galapagos NV, GSK, Glenmark Pharmaceuticals, Inventiva, Italfarmaco, IQvia, medac, Medscape, Mitsubishi Tanabe Pharma, MSD, Roche, Sanofi and UCB, Speakers bureau: Speaker fees from Actelion, Bayer, Boehringer Ingelheim, Medscape, Pfizer and Roche

DOI: 10.1136/annrheumdis-2020-eular.3671

OP0250

EFFICACY AND SAFETY OF ROMILKIMAB IN DIFFUSE CUTANEOUS SYSTEMIC SCLEROSIS (DCSSC): RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED, 24-WEEK, PROOF OF CONCEPT STUDY

Y. Allanore¹, P. Wung², C. Soubrane³, C. Esperet³, M. Frederic³, R. Bejuit³, A. Lahmar², D. Khanna⁴, C. Denton⁵, ¹Cochin Hospital, Rheumatology, Paris, France; ²Sanofi R&D, Bridgewater, United States of America; ³Sanofi R&D, Chilly Mazarin, France; ⁴University of Michigan, Ann Arbor, United States of America; ⁵University College London, London, United Kingdom

Background: Systemic sclerosis (SSc) is a progressive, multi-organ disease with limited treatment options. Interleukin-4 (IL-4) and IL-13 have been implicated in the fibrotic pathway and pathogenesis of SSc and are promising targets. Romilkimab (RKB) is an engineered humanized bispecific Ig-G4 antibody that binds and neutralizes both IL-4/IL-13. We report a Phase IIa randomized, double-blind, placebo-controlled trial (NCT02921971, Sanofi funded) employing RKB in SSc.

Objectives: To evaluate the efficacy and safety of RKB in dcSSc.

Methods: Patients with dcSSc duration ≤36 months, mRSS 10-35, with or without immunosuppressive background therapy were randomized (1:1) to subcutaneous RKB 200mg or placebo (PBO) for 24 weeks and stratified on history of SSc-ILD. Primary endpoint was mean change in mRSS at Week 24 and FVC/DLCO and HAQ-DI were secondary endpoints. All analyses used a 1-sided p-value <0.05 as reaching statistical significance.

Results: Ninety-seven patients with similar baseline characteristics between arms, including use of background therapy (RKB 59.2% vs. PBO 52.1%) were randomized. Six (12.2%) and 4 (8.3%) patients discontinued study treatment early in the PBO and RKB arms, respectively. Primary endpoint showed an absolute change in mRSS of -2.45 (0.85) and -4.76 (0.86) for PBO and RKB groups, respectively with a difference of -2.31 (1.21) favoring RKB (p=0.029). Subgroup analysis based on background therapy showed a similar treatment effect with a PBO subtracted difference in mRSS of -2.69 (1.83) and -2.38 (1.59), suggesting an additive effect between background therapy and RKB. Secondary endpoints did not show a statistically significant difference between RKB vs. PBO arms, although there was numerically less decline in FVC with RKB with a PBO subtracted difference of 70ml (p=0.06). Exploratory endpoints suggested possible effect of RKB on overall pain, Raynaud's, digital ulcers, and EQ-5D-5L. Post-hoc analysis was undertaken to determine time to progression (first event defined as death, ≥10% relative decline in % predicted FVC, ≥15% relative decline in % predicted DLCO, ≥20% increase or +5 in mRSS, or other events: cardiac, SRC, PAH development) and showed a benefit for RKB (HR: 0.47 p=0.04). Adverse events were balanced between the two groups (RKB 83.3% vs. PBO 83.7%). There were 5 and 4 SAEs in the PBO and RKB arms, respectively. One death occurred in each arm (SRC – RKB, cardiomyopathy – PBO).

Conclusion: Patients with dcSSc who were treated with RKB showed a statistically significant reduction in mRSS compared to those receiving PBO. Secondary outcomes were not met, although RKB was associated with a smaller decline in FVC than PBO. Post-hoc analysis showed a possible reduction on time to progression with RKB. RKB was well tolerated with no major safety concerns.

References: None.

Disclosure of Interests: Yannick Allanore Grant/research support from: Yannick Allanore has received grants from Inventiva, Roche and Sanofi, Consultant of: Yannick Allanore has received fees from Actelion, Bayer, Boehringer Ingelheim, Bristol-Myers Squibb, Curzion, Inventiva, Roche, Sanofi, Peter Wung Shareholder of: I own Sanofi stock, Employee of: I work for Sanofi, Christina Soubrane Employee of: I work for Sanofi., Corinne Esperet Employee of: I work for Sanofi., MARRACHE Frederic Employee of: I work for Sanofi,