AB0670

AXIAL MANIFESTATIONS IN PATIENTS WITH AXIAL SPONDYLOARTHRITIS AND PSORIATIC ARTHRITIS: ARE THEY SIMILAR?

D. Benavent¹, V. Navarro-Compán¹, C. Plasencia¹, D. Peiteado¹, A. Villalva¹, A. Balsa¹. ¹Hospital La Paz, Idipaz, Rheumatology Unit, Madrid, Spain

Background: Spondyloarthritis (SpA) is a group of heterogeneous diseases that includes axial SpA (axSpA), such as ankylosing spondylitis and axial non-radiographic SpA, and Psoriatic Arthritis (PsA) with peripheral and/or axial involvement (axPsA). Currently, it is not well known if the characteristics and burden of the disease in patients with axPsA are similar to that of patients with axSpA.

Objectives: To compare the demographic, clinical and structural features between patients with axSpA and axPsA.

Methods: Data from an observational prospective cohort including all patients with SpA initiating biological therapy because of predominant axial manifestations from 2002-2019 in a university hospital were analyzed. AxSpA and axPsA were defined in clinical practice according to the prescribing rheumatologist, based on clinical features and complementary examinations. Demographic information, laboratory tests, disease presentation, sacroiliitis according to modified New York criteria in the pelvis X-ray, disease activity indexes (ASDAS and BASDAI) and concomitant treatment before starting biological drug were collected from the electronic medical record and biologic database. In the statistical analysis, chi square or the exact Fisher's test was used for categorical and t-student or U-Mann Whitney for continuous variables, according to the distribution of the data. Then, the association between demographic and clinical features and each disease was analysed using univariable and multivariable logistic regression models.

Results: Out of 352 included patients, 287 (81.5%) had axSpA, and 65 had axPsA (18.5%). Baseline characteristics are shown in Table 1. Mean baseline ASDAS was 3.3±0.9 and 3.1±1.0 for axSpA and axPsA, respectively. Biological therapies initiated can be seen in Figure 1. No significant differences at baseline were observed between axSpA and axPsA for most of the characteristics including: gender, age at diagnosis, age at starting biologic, disease duration before biologic, smoking habit, CRP, disease activity, enthesitis, dactylitis, inflammatory bowel disease (IBD), patient global assessment and sulfasalazine use. However, there were differences between diseases in some relevant characteristics. AxSpA patients had less peripheral involvement (41.5 vs. 78.5 %, p=0.004), more uveitis (15.3 vs. 3.1 %, p=0.03) and were more frequently HLA-B*27 positive (72.3 vs 34.1 %, p<0.001), in comparison to axPsA patients. They also had better physician global assessments (PhGA) (37.4 vs 44.4, p=0.02), and a higher grade of radiographic sacroiliitis. AxSpA patients used less global baseline concomitant therapy (p=0.001), methotrexate (p<0.001) and prednisone (p<0.01), whereas they used more sulfasalazine (p=0.003) than axPsA patients in our cohort. After running multivariate analyses, the absence of peripheral manifestations (OR=4.7; p<0.001) and the positivity of HLA-B27 (OR=5.4; p<0.001) were independently associated with axSpA.

Table 1. Baseline stratified characteristics. Results are shown as absolute numbers (percentages) or mean ± standard deviation.

	AxSpA & AxPsA n= 352	AxSpA n= 287	axPsA n= 65	p-value
Sex (male)	223 (60.6)	180 (62.7)	43 (66.2)	0.7
Age (years):				
At diagnosis	35.9 ± 13.4	35.7 ± 13.7	36.9 ± 12.1	0.9
At biologic starting	44.4 ± 13.2	44.1 ± 13.4	45.8 ± 11.6	0.3
Disease duration before biologic (years)	17.9 ± 10.3	7.9 ± 11.3	8.9 ± 9.0	0.7
Current smoking habit	158 (44.9)	129 (44.9)	29 (44.6)	0.9
HLA B27 positive	219/322(67.8)	204/281 (72.3)	16/47 (34.1)	<0.001
CRP(mg/dL)	12.4 ± 17.9	12.6 ± 18.9	11.1 ± 12.7	0.5
Clinical involvement:				
Only axial	170 (48.2)	168 (58.5)	14 (21.5)	<0.001
Axial and peripheral	182 (51.7)	119 (41.5)	51 (78.5)	
Psoriasis	74 (21.3)	11 (4.2)	63 (97)	<0.001
ASDAS	3.3 ± 0.9	3.3 ± 1.0	3.1 ± 1.0	0.1
BASDAI (0-10)	5.9 ± 4.2	6.1 ± 4.5	5.23 ± 2.1	0.1
Enthesitis	85 (41.5)	73 (25.4)	12 (28.6)	0.07
Dactylitis	10 (2.7)	7 (2.4)	3 (4.6)	0.4
IBD	9 (2.6)	8 (2.8)	1 (1.5)	0.7
Uveitis	46 (13.6)	44 (15.3)	2 (3.1)	0.03
PGA (0-100)	63.2 ± 21.8	64.1 ± 21.5	58.8 ± 23.2	0.1
PhGA (0-100)	39.1 ± 21.5	37.4 ± 13.7	44.4 ± 22.6	0.02
Radiographic sacroiliitis, mNY criteria	227 (64.5)	203 (70.7)	24 (36.9)	< 0.001
Concomitant therapy:				
Monotherapy	193 (52.4)	145 (50.5)	48 (73.8)	0.001
Only MTX	66(20.7)	36 (13.9)	30 (46.2)	< 0.001
Only SZS	82 (25.6)	73 (28.2)	9(13.8)	0.03
Prednisone use	32 (9.5)	20 (7.7)	12 (20)	0.004

Conclusion: Despite being spondyloarthritis with many common traits, axSpA and axPsA present some differences in clinical practice. Whereas axSpA patients are more frequently HLA-B27 positive, axPsA have more peripheral involvement. These differences in clinical presentation between both diseases may contribute to variances in therapeutic management, such as increased use of baseline concomitant therapy in axPsA patients who initiate biological therapy.

Figure 1. Biological therapies initiated in axSpA and axPsA

Disclosure of Interests: Diego Benavent: None declared, Victoria Navarro-Compán Consultant of: Abbvie, Lilly, Novartis, Pfizer, UCB, Speakers bureau: AbbVie, MSD, Lilly, Novartis, Pfizer, UCB, Chamaida Plasencia: None declared, Diana Peiteado: None declared, Alejandro Villalva: None declared, Alejandro Balsa Grant/research support from: BMS, Roche, Consultant of: AbbVie, Gilead, Lilly, Pfizer, UCB, Sanofi, Sandoz, Speakers bureau: AbbVie, Lilly, Sanofi, Novartis, Pfizer, UCB, Roche, Nordic, Sandoz

DOI: 10.1136/annrheumdis-2020-eular.1703

AB0671

PREVALENCE OF EXTRA-ARTICULAR MANIFESTATIONS AND IMPACT ON TARGETED DRUG PRESCRIPTION IN PATIENTS WITH SPONDYLOARTHRITIS: A RETROSPECTIVE ANALYSIS OF A REAL-LIFE COHORT

M. Biggiongero¹, E. G. Favalli¹, A. Marchesoni¹, R. Caporali^{1,2}. ¹Gaetano Pini-CTO Institute, Department of Rheumatology, Milan, Italy; ²Università degli Studi di Milano, Department of Clinical Sciences & Community Health, Milan, Italy

Background: Extra-articular manifestations (EAMs), such as uveitis, inflammatory bowel diseases (IBD), and psoriasis (PSO) can frequently complicate the disease course of patients with spondyloarthritis (SpA), although prevalence data on this regard are still controversial. The occurrence of EAMs may influence the decision to introduce a targeted therapy and also drive the choice of the most appropriate drug.

Objectives: The aim of this study is to retrospectively evaluate the prevalence of EAMs in a real-life cohort of SpA patients who were eligible to receive a targeted therapy and to investigate their impact in the choice of targeted treatment.

Methods: Clinical data of SpA (axial SpA [axSpA], peripheral SpA, and psoriatic arthritis [PsA]) patients treated with a biologic or targeted synthetic Disease-Modifying Anti-Rheumatic Drug (DMARD) between December 1999 and December 2019 were extracted from a local registry. Prevalence of main SpA-related EAMs (uveitis, IBD and PSO) was calculated at the time of drug prescription, evaluating their distribution according to treatment subgroups. Comparisons between disease and treatment subgroups were made using the Fisher's test.

Results: The study included 629 patients with SpA (axSpA 26%, peripheral SpA 24%, PsA 50%), 266 [42%] women, mean age [±SD] 52 [±13.2] years, mean disease duration 7.8 [±7.9] years), receiving a total of 1106 lines of targeted treatment (I-line n=629, II-line n=258, ≥ III-line n=219) with etanercept (n=177), anti-TNF monoclonal antibodies (397 infliximab, 273 adalimumab, 38 certolizumab pegol, and 130 golimumab), secukinumab (n=46), ustekinumab (n=28), or apremilast (n=18). At the time of drug introduction, 13% of SpA patients showed at least one EAM. The prevalence of uveitis was higher in axSpA (11.8%) compared with both peripheral SpA (5.5%, p=0.01) and PsA (2.8%, p<0.0001), while IBD was more frequent in peripheral SpA (15.6%) than in axSpA (8.1%, p=0.008) and PsA (4.7%; p<0.0001). The prevalence of PSO was similar in axial and peripheral SpA (8.4 versus 6.3%, respectively;