Conclusion: In pCJIA pts, tofacitinib vs PBO resulted in significantly fewer disease flares, and improved time to flare, disease activity and physical functioning. Tofacitinib safety was consistent with that in RA pts.

Disclosure of Interests: Nicolino Ruperto Grant/research support from: Bristol-Myers Squibb, Eli Lilly, F Hoffmann-La Roche, GlaxoSmithKline, Janssen, Novartis, Pfizer, Sobi (paid to institution), Consultant of: Ablynx, AbbVie, AstraZeneca-Medimmune, Biogen, Boehringer Ingelheim Bristol-Myers Squibb, Eli Lilly, EMD Serono, GlaxoSmithKline, Hoffmann-La Roche, Janssen, Merck, Novartis, Pfizer, R-Pharma, Sanofi, Servier, Sinergie, Sobi, Takeda, Speakers bureau: Ablynx, AbbVie, AstraZeneca-Medimmune, Biogen, Boehringer Ingelheim, Bristol-Myers Squibb, Eli Lilly, EMD Serono, GlaxoSmithKline, Hoffmann-La Roche, Janssen, Merck, Novartis, Pfizer, R-Pharma, Sanofi, Servier, Sinergie, Sobi, Takeda, Olga Synoverska Speakers bureau: Sanofi, Tracy Ting: None declared, Carlos Abud-Mendoza Speakers bureau: Eli Lilly, Pfizer Inc, Alberto Spindler Speakers bureau: Eli Lilly, Yulia Vyzhga Grant/research support from: Pfizer Inc, Katherine Marzan Grant/research support from: Novartis, Vladimir Keltsev: None declared, Irit Tirosh: None declared, Lisa Imundo: None declared, Rita Jerath: None declared, Daniel Kingsbury: None declared, Betül Sözeri: None declared, Sheetal Vora: None declared, Sampath Prahalad Grant/research support from: Novartis, Elena Zholobova Grant/research support from: Novartis and Pfizer Inc, Speakers bureau: AbbVie, Novartis, Pfizer Inc and Roche, Yonatan Butbul Aviel: None declared, Vyacheslav Chasnvyk: None

P. Quartier¹, E. Feist², D. J. Lovell³, H. Umabayashi⁴, N. Ruperto⁵, H. Brunner³, C. Dunger-Baldauf⁶, S. Novioello⁷, S. Whelan⁸. ¹Necker Hospital, Université de Paris, Paris, France; ²Helios Department of Rheumatology, Vogelsang-Gommern, Germany; ³Cincinnati Children's Hospital Medical Center, Cincinnati, United States of America; ⁴Miyagi Children's Hospital, Miyagi, Japan; ⁵Clinica Pediatrica e Reumatologia, IRCCS Istituto Giannina Gaslini, Genoa, Italy; ⁶Novartis Pharma AG, Basel, Switzerland; ⁷Novartis Pharmaceuticals Corporation, East Hanover, United States of America; ⁸Novartis Ireland Limited, Dublin, Ireland

Background: Still's disease is a systemic auto-inflammatory disease with a pediatric form, sJIA, and an analogous condition in adults, adult-onset Still's disease (AOSD). The role of interleukin-1 (IL-1) in the pathophysiology of Still's disease is well established. Canakinumab, a monoclonal antibody against IL-1 β , is approved to treat patients with Still's disease in Europe (sJIA and AOSD) and the United States (sJIA).

Objectives: To study the efficacy of canakinumab in sJIA patients categorized by age, we performed an intention-to-treat analysis of pooled data from 5 clinical trials, as an addition to a previously communicated analysis including 3 of the studies¹

Methods: The age categories were children (2-<12 years), young adolescents (12-<16 years) and older adolescents and young adults (16-<20 years). We pooled efficacy results from patients with active disease at baseline treated during a 12-week period with canakinumab (4mg/kg every 4 weeks), including the presence of intermittent fever, serum concentrations of C reactive protein (CRP), improvement of sJIA (adapted pediatric ACR 30, 70 and 100 responses) and JIA ACR inactive disease status. Safety was assessed by analysis of reported adverse events (AEs).

Results: 302 children, 82 young adolescents and 34 older adolescents and young adults were included in the analysis, with a mean disease duration of 922, 1708 and 2615 days, respectively. Prior therapy with other biologics was common, with anakinra used in 33%, 35% and 47% of patients in each group. Disease severity was comparable among groups, with the mean number of active joints ranging from 11.8 to 13.7. Adapted pediatric ACR responses revealed a rapid response to canakinumab, with all groups showing similar rates of responders at most time points (Table 1). In each age group, the proportion of patients with inactive disease progressively increased to Day 57. At all time points after Day 15, the 16-<20 years group presented the highest proportion of patients with inactive disease. Median CRP levels decreased from baseline to reach values in the normal range (<10mg/L) from Day 29 onwards in the three groups, with improvements more marked in the 16-<20 years group. The safety profile was similar in the three age groups analyzed, with a lower proportion of 16-<20 years old patients experiencing serious AEs (28%) as compared to children (35%) and young adolescents (42%).

Table 1. Percentages of patients with Adapted pediatric ACR responses and inactive disease status over time*

	Time of treatment (Days)	2 - <12 years ^a	12 - <16 years ^b	16-<20 years ^c
ACR30 (%)	15	72.7	81.7	83.9
	29	77.5	84.1	82.4
	57	76.2	82.9	88.2
	85	65.5	74.5	83.3
	15	51.5	58.3	64.5
ACR70 (%)	29	61.9	62.2	70.6
	57	65.2	58.5	79.4
	85	58.6	61.8	75.0
	15	21.6	25.0	12.9
	29	29.5	30.5	35.3
ACR100 (%)	57	36.1	34.1	38.2
	85	34.1	30.9	33.3
	15	19.0	30.0	19.4
	29	34.1	34.1	47.1
	57	39.4	36.6	55.9
Inactive disease (%)	85	36.7	43.4	52.2

*Some studies did not include visits at Day 15 and/or 85. For Day 15, 29, 57 and 85 the respective denominators for each age group were: ^aN = 231, 302, 302, 232; ^bN = 60, 82, 82, 55; ^cN = 31, 34, 34, 24.

Conclusion: The efficacy and safety profile of canakinumab was consistent in children, adolescents and young adults with sJIA. Since sJIA and AOSD represent pediatric- and adult- onset variants of the Still's disease continuum, these results further support the therapeutic effect of canakinumab 4 mg/kg every 4 weeks in both children and adults with Still's disease.

References:

[1] Feist et al. *Clin Exp Rheumatol*. 2018;36(4):668-75.

Disclosure of Interests: Pierre Quartier Consultant of: AbbVie, Chugai-Roche, Lilly, Novartis, Sanofi, Sobi, Speakers bureau: AbbVie, BMS, Chugai-Roche, Novartis, Pfizer, Sobi, Eugen Feist Consultant of: Novartis, Roche, Sobi, Lilly, Pfizer, Abbvie, BMS, MSD, Sanofi, Speakers bureau: Novartis, Roche, Sobi, Lilly, Pfizer, Abbvie, BMS, MSD, Sanofi, Daniel J Lovell Consultant of: Abbott (consulting and PI), AbbVie (PI), Amgen (consultant and DSMC Chairperson), AstraZeneca, Boehringer Ingelheim, Bristol-Myers Squibb (PI), Celgene, Forest Research (DSMB Chairman), GlaxoSmithKline, Hoffman-La Roche, Janssen (co-PI), Novartis (consultant and PI), Pfizer (consultant and PI), Roche (PI), Takeda, UBC (consultant and PI), Wyeth, Employee of: Cincinnati Children's Hospital Medical Center, Speakers bureau: Wyeth, Hiroaki Umebayashi: None declared, Nicolino Ruperto Grant/research support from: Bristol-Myers Squibb, Eli Lilly, F Hoffmann-La Roche, GlaxoSmithKline, Janssen, Novartis, Pfizer,

Sobi (paid to institution), Consultant of: Ablynx, AbbVie, AstraZeneca-Medimmune, Biogen, Boehringer Ingelheim, Bristol-Myers Squibb, Eli Lilly, EMD Serono, GlaxoSmithKline, Hoffmann-La Roche, Janssen, Merck, Novartis, Pfizer, R-Pharma, Sanofi, Servier, Sinergie, Sobi, Takeda, Speakers bureau: Ablynx, AbbVie, AstraZeneca-Medimmune, Biogen, Boehringer Ingelheim, Bristol-Myers Squibb, Eli Lilly, EMD Serono, GlaxoSmithKline, Hoffmann-La Roche, Janssen, Merck, Novartis, Pfizer, R-Pharma, Sanofi, Servier, Sinergie, Sobi, Takeda, Hermine Brunner Consultant of: Hoffman-La Roche, Novartis, Pfizer, Sanofi Aventis, Merck Serono, AbbVie, Amgen, Alter, AstraZeneca, Baxalta Biosimilars, Biogen Idec, Boehringer, Bristol-Myers Squibb, Celgene, EMD Serono, Janssen, MedImmune, Novartis, Pfizer, and UCB Biosciences, Speakers bureau: GSK, Roche, and Novartis, Cornelia Dunger-Baldauf Employee of: Novartis, Stephanie Novello Employee of: Novartis, Sarah whelan Employee of: Novartis
DOI: 10.1136/annrheumdis-2020-eular.3759

OP0293

USE OF WHOLE-BODY MAGNETIC RESONANCE TO IDENTIFY POTENTIAL DIAGNOSTIC CLUES IN CHILDREN WITH FEVER OF UNKNOWN ORIGIN (FUO)

S. Signa^{1,2}, R. Caorsi¹, G. Stagnaro³, F. Minoia⁴, P. Picco⁴, A. Ravelli^{2,4}, M. B. Damasio³, M. Gattorno¹. ¹IRCCS Istituto Giannina Gaslini, UOSD Centro Malattie Autoinfiammatorie ed Immunodeficienze, Genoa, Italy; ²University of Genoa, DINOGMI, Genoa, Italy; ³IRCCS Istituto Giannina Gaslini, UOC Radiologia, Genoa, Italy; ⁴IRCCS Istituto Giannina Gaslini, UOC Clinica Pediatrica e Reumatologia, Genoa, Italy

Background: Whole-body magnetic resonance imaging (WBMRI) is a fast and accurate method to detect diseases throughout the entire body without exposure to ionizing radiation. Possible emerging applications for this technique include rheumatologic field and evaluation of fever of unknown origin (FUO).

Objectives: To evaluate the ability of WBMRI to identify significant potential diagnostic clue (PDC) in patients presenting a non specific inflammatory clinical picture.

Methods: We retrospectively collected cases of pediatric patients followed in a single pediatric rheumatology center who underwent WBMRI between January 2010 and December 2015 for the following indications: i) FUO (temperature greater than 38.3°C for more than three weeks or failure to reach diagnosis after one week of investigations), ii) recurrent fever (febrile episodes separated by periods of normal temperature), iii) Inflammation of unknown origin, IUO (an illness of at least 3 weeks' duration, with raised inflammatory markers and fever below 38.3°C). WBMRI studies were acquired with coronal and sagittal planes (slice thickness 5mm) with acquisition of several image sets with automatic direct image realignment after acquisition creating a whole-body scan. Sequences include short τ inversion recovery (STIR) and T1-weighted. All studies have been evaluated twice, the second time according to a predefined checklist, defined by an experienced radiologist, considering systematically single/multifocal bone lesion, bone marrow, joint effusion, soft tissues, adenopathies, parenchymal and vessels looking for PDC. We considered as a Potential Diagnostic Clue each alteration of the examined district that can potentially guide the diagnosis. Each alteration found is a PDC. We retrospectively evaluated patients' clinical history and final diagnosis and we classified the PDCs identified during both first evaluation and re-evaluation as: Not useful (the identified PDC did not guide the diagnosis and is not coherent with the final diagnosis), consistent (the identified PDC is congruent with the patient's final diagnosis) or diagnostic (the identification of the considered PDC strongly orient the final diagnosis).

Results: We collected 104 patients who underwent WBMRI; 24 (23%) of them presenting FUO, 28 (27%) presenting recurrent fever and 52 (50%) presenting IUO. The mean age of onset symptoms was 6 years and nine months (range: 2 weeks old- 17 years and 6 months). The mean age of execution of WBMRI was 9 years (range: 5 months old- 19 years). After the whole diagnostic work-out a final diagnosis was achieved in 44 patients (42%). PDCs were identified at the first evaluation in 78/104 cases (75%). In 22 cases (21%) the identified PDCs were consistent with the diagnosis, whereas in 9 cases (8.5%) the identified PDCs were considered diagnostic. Globally we can consider that at first evaluation PDCs were somehow contributory to the diagnosis in 31 cases (30%); 6 JIA, 7 systemic infections, 5 monogenic inflammatory diseases, 4 ALPS, 2 Goldbloom's Syndrome, 2 Vasculitis, 1 eosinophilic fasciitis, 1 hystiocytosis, 3 oncologic diagnosis). Blind re-evaluation of WBMRI allowed the identification of additional PDCs in 52 patients (12 of them previously negative). In 10 cases the PDC found after re-evaluation were consistent with the final diagnosis (2 JIA, one infectious disease, one neuroblastoma, 3 ALPS, 1 monogenic inflammatory disease, 1 Takayasu arteritis, 1 Goldbloom's syndrome).

Conclusion: WBMRI can be a powerful diagnostic tool in patients with FUO. A predefined checklist increases sensitivity of WBMRI in the identification of PDC.

Disclosure of Interests: Sara Signa: None declared, Roberta Caorsi: None declared, Giorgio Stagnaro: None declared, Francesca Minoia: None declared, Paolo Picco: None declared, Angelo Ravelli: None declared, Maria Beatrice