Conclusion: Demographic and clinical factors differentiate axSpA from nr-ax-SpA patients. Diagnostic delay was higher in rad-axSpA compared with nr-ax-SpA despite the same treatment. Some lesions of spine/SI at CR and MRI, and psoriasis, were mostly associated with diagnostic delay and sex.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.3356

AB1086

POWER DOPPLER AND SPECTRAL DOPPLER ULTRASOUND IN SUSPECTED ACTIVE SACROILIITIS: A COMPARISON WITH MAGNETIC RESONANCE IMAGING AS GOLD STANDARD

P. Falsetti¹, E. Conticini¹, C. Baldi¹, M. Bardelli¹, S. Gentileschi¹, R. D'aleandro¹, L. Cantarini¹, B. Frediani¹. ¹University of Siena, Rheumatology Unit, Siena, Italy

Background: SIJ involvement is a characteristic feature of Spondylarthritis (SpA). Magnetic Resonance imaging (MRI) has been included in the new Assessment of SpA International Society (ASAS) criteria for the classification of Axial SpA. Gray scale US, Color Doppler ultrasound (CDUS), contrast-enhanced CDUS, and spectral Doppler (SD) US has been used in few works to evaluate the inflammatory activity of the SIJ with not conclusive results. Power Doppler ultrasound (PDUS) was not yet applied to the study of SIJ with active SI.

Objectives: The aim of this work was to study with PDUS and SD US the SIJ of patients with suspected active SI, to describe inflammatory flows with spectral wave analysis (SWA) in duplex Doppler US, and to correlate US data with clinical characteristics and the presence of bone marrow edema (BME) in MRI.

Methods: 22 patients (18 females and 4 males, mean age 35 years) with new onset of inflammatory back pain (IBP), were included. Every patient underwent an US examination in prone position. The sonographers were blinded to the clinical data of the patient. A Esaote Twice US machine, equipped with a convex multifrequency 1-8 MHz probe, was used, with standardized parameters: 1-5 MHz for gray scale, 1.9-2.3 MHz frequency for Doppler with Pulse Repetition Frequency (PRF) of 1.0 KHz and a color gain just under the artifact limit. SIJ was located as the hypoechoic triangle delimited between the sacrum and iliac bone, and the posterior SI ligament as the upper margin. The first sacral foramen was always localized to avoid measurement of the normal pre-sacral arteries. The PDUS was applied, and if any signals were detected in the SIJ, they were scored with a 3-points scale: 0= absence of signals, 1= isolate vessels, 2= more than one vessel. The signals were also classified as intra-articular or peri-articular. The same vessels were also evaluated using quantitative SD calculating the Resistive Index (RI=peak of systolic flow- end diastolic flow/peak systolic flow), ranging between 0 and 1. Every patient underwent MRI of SIJ within the same week, before treatment. A statistical analysis was performed, estimating the sensitivity and specificity against the gold standard (presence of BME in the same SIJ according to ASAS criteria). The Spearman rank non-parametric test was applied to correlate the presence and grading of BME with PDUS grading and RI. A regression analysis was applied between PDUS results and clinical characteristics.

Results: In 14/22 SIJ MRI revealed BME. In 13 of them, PDUS confirmed abnormal hypervascularisation in the intrarticular portion of SI, and in 3 in the peri-articular site too. Two SIJ showed hypervascularisation at PD with no BME in MRI. A significant correlation was demonstrated between positivity and grading of PD and presence of BME in MRI ($p=0.0005$). SD analysis demonstrated low Resistance Index (RI) values in 14 SIJ (mean 0.57). An inverse correlation was demonstrated between RI and grading of BME in MRI ($r=-0.6229$, $p=0.044$). The diagnostic accuracy of SD for detection of active SI varied on the basis of RI cut-off value. The best values of sensitivity (62.5%) and specificity (61.5%) were obtained with a RI cut-off values of 0.60. A multiple regression model demonstrated a significant relationship between PDUS signals and ASDAS ($p=0.0382$), but not with inflammatory reactants.

Conclusion: PDUS and SD US of SIJ can be useful as first imaging assessment in suspected active SI, demonstrating a good diagnostic accuracy compared with MRI. Intra-articular low RI values (<0.60) on SD indicate active SI with good accuracy. Moreover, PDUS signals into the SIJ correlate with clinical symptoms but not with inflammation reactants.

Figure 1. Doppler US in SI. Right SIJ with a Doppler signal along the posterior SIJ ligament, and another Doppler signal into the joint, where SD analysis gave a RI of 0.62.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.2281

AB1087

DETECTING AXIAL AND PERIPHERAL NEW BONE FORMATION IN SPONDYLOARTHRITIS PATIENTS USING [18F]FLUORIDE PET-CT IMAGING

J. De Jongh¹, R. Hemke², G. C. J. Zwezerijnen³, M. Yaqub³, I. Van der Horst-Bruinsma¹, M. G. H. Van de Sande⁴, A. Van Kuijk⁵, I. Bultink¹, L. Burgemeister⁵, N. M. A. Van Dillen⁵, A. Voskuy¹, C. J. Van der Laken¹. ¹Amsterdam UMC Location VUmc, Rheumatology and Clinical Immunology, Amsterdam, Netherlands; ²Amsterdam UMC Location AMC, Radiology and Nuclear Medicine, Amsterdam, Netherlands; ³Amsterdam UMC Location VUmc, Radiology and Nuclear Medicine, Amsterdam, Netherlands; ⁴Amsterdam UMC Location AMC, Rheumatology and Clinical Immunology, Amsterdam, Netherlands; ⁵Reade, Rheumatology, Amsterdam, Netherlands

Background: Bone formation in spondyloarthritis (SpA) is presumably related to local enthesitis/peri-articular inflammation and ultimately may lead to functional limitation (1,2). X-rays only allow long-term monitoring of bone formation (≥ 2 years) (3). Imaging techniques that can visualize bone formation at an early stage would therefore be valuable. Positron Emission Tomography (PET) using [¹⁸F]Fluoride can visualize and quantify (early changes in) bone formation at molecular level (4).

Objectives: To investigate the feasibility of [¹⁸F]Fluoride to assess new bone formation at axial and peripheral enthesial sites in SpA patients.

Methods: Thus far, 5 of the total of 15 patients with clinically active ankylosing spondylitis (AS) (according to modified New York criteria and BASDAI ≥ 4) and 8 of the 25 patients with active psoriatic arthritis (PsA) (according to CASPAR criteria and ≥ 1 clinically active enthesitis) were included. Of each patient, a whole body [¹⁸F]Fluoride PET-CT scan was performed. All scans were visually judged and scored dichotomously by one reader (blinded for clinical data) for PET-positive lesions in the spine, peripheral entheses sites and joints. Low dose CT was used for anatomical reference.

Results: The study is ongoing, with whole body [¹⁸F]Fluoride PET-CT scans available in five AS patients and eight PsA patients. In 4/5 AS scans, at least (≥ 1) PET positive lesions were found in the cervical, thoracic and/or lumbar vertebrae. These were mainly found in anterior corners of vertebrae and bridging syndesmophytes (Fig. 1A). In all eight PsA patients, at least 1 PET positive lesion was visualized, projected either at the site of a tendon attachment (fascia plantaris, achilles- and patella tendon (Fig 1B)) or peri-articularly (in the ankle or wrist).

Fig 1. [^{18}F]Fluoride uptake in the cervical, thoracic and lumbar spine in a clinically active AS patient (A) and in the patella tendon of the right knee in a clinically active PsA patient (B)

Conclusion: [^{18}F]Fluoride PET uptake, reflecting new bone formation, can be visualized at heterogeneously distributed entheses and (peri-)articular sites in AS- and PsA patients. The technique therefore is sensitive to visualize new bone formation and may reflect local disease activity. Additional scans will be collected and analyzed quantitatively, also after anti-TNF or Secukinumab treatment, to further investigate the applicability of [^{18}F]Fluoride PET for monitoring of therapeutic effects on bone formation in SpA.

References:

- [1] Maksymowych WP, Mallon C, Morrow S, Shojania K, Olszynski WP, Wong RL, et al. Development and validation of the Spondyloarthritis Research Consortium of Canada (SPARCC) Enthesitis Index. *Ann Rheum Dis*. 2009;68(6):948-53.
- [2] Rezvani A, Bodur H, Ataman S, Kaya T, Bugdayci DS, Demir SE, et al. Correlations among enthesitis, clinical, radiographic and quality of life parameters in patients with ankylosing spondylitis. *Mod Rheumatol*. 2014;24(4):651-6.
- [3] Rudwaleit M, Khan MA, Sieper J. The challenge of diagnosis and classification in early ankylosing spondylitis: do we need new criteria? *Arthritis Rheum*. 2005;52:1000-8.
- [4] Bruijnen ST, Verweij NJF, van Duivenvoorde L, Bravenboer N, Baeten D, van Denderen JC, et al. [^{18}F]Fluoride PET-CT imaging of bone formation in ankylosing spondylitis before and after 12 weeks of anti-TNF treatment. 2017.

Acknowledgments: We thank EULAR Forum, Pfizer and Novartis for financial support of this investigator initiated study.

Disclosure of Interests: Jerney de Jongh: None declared, Robert Hemke: None declared, Gerben C.J. Zwezerijnen: None declared, Maqsood Yaqub: None declared, Irene van der Horst-Bruinsma Grant/research support from: AbbVie, Novartis, Eli Lilly, Bristol-Myers Squibb, MSD, Pfizer, UCB Pharma, Consultant of: AbbVie, Novartis, Eli Lilly, Bristol-Myers Squibb, MSD, Pfizer, UCB Pharma, Marleen G.H. van de Sande Grant/research support from: Novartis, Eli Lilly, UCB, Jansen, Consultant of: AbbVie, Novartis, Eli Lilly, MSD, Arno Van Kuijk: None declared, Irene Bultink: None declared, Lot Burgemeister: None declared, Nancy M.A. van Dillen: None declared, Alexandre Voskuyl: None declared, Conny J. van der Laken: None declared

DOI: 10.1136/annrheumdis-2020-eular.4216

AB1088

NAIL PSORIASIS: THE UNDERESTIMATED DISORDER IN PSORIASIS AND PSORIASIS ARTHRITIS. CAN ULTRASOUND AND CAPILLARY MICROSCOPY IN PATIENTS WITH NAIL PSORIASIS SPEED UP OUR DIAGNOSIS AND THERAPY?

B. C. Dettler¹, S. El-Nawab-Becker², M. Toellner¹, A. Kleinheinz², T. Witte⁴.

¹Clinic Dr. Hancken GmbH, Stade, Germany; ²Rheumatologic Practice,

Hamburg, Germany; ³Klinik für Dermatologie, Buxtehude, Germany;

⁴Medizinische Hochschule Hannover, Hannover, Germany

Background: Nail psoriasis is an extreme diagnostic and therapeutic challenge and represents an enormous physical and psychological burden for affected patients. 50% of patients with psoriasis vulgaris develop nail involvement (NailPso) during the course of their disease. NailPso is the strongest predictor of psoriatic arthritis (PsA). Through the synovio-enthesis concept we have learned that there is an anatomical-pathophysiological relationship between DIP joint, extensor tendon and nail matrix. We have observed in daily practice that hypervascularization (HV) in ultrasound Power Doppler (US-PD) the nail matrix may be a pathognomonic element in its own right. There are no data on this in the literature.

Objectives: Is there a difference in the ultrasound PD examination of the DIP joint and nail area and in the capillary microscopy of the corresponding nail fold in patients with psoriasis vulgaris and nail psoriasis versus patients with psoriasis vulgaris without nail psoriasis.

Methods: Monocentric prospective study of all consecutive patients with psoriasis vulgaris who have come to a rheumatic practice to clarify a PsA. In addition to demographic data, assessments (PASI, DLQI, CASPAR, GEPARD, DAS28, SJ, TJ, FFBH), clinical examination, a standardized ultrasound PD examination and capillary microscopy of the affected fingertips in PsO patients suffering from nail psoriasis was performed as well as corresponding examinations of the 2nd and 3rd finger right in PsO patients without nail involvement.

Results: 79 patients could be included during the study period. Thereof 25 PsO patients without nail involvement and 44 PsO patients with nail involvement. Since the patients were examined consecutively, the difference results. There was no difference in age, BMI and sex in both groups (PsO and NailPso). The Caspar criteria as classification criteria for a PsA were positive in 65% of the NailPso patients and positive in 50% of all PsO patients without nail involvement. Hypervascularization in the US-PD examination in the area of the nail matrix could be seen significantly more frequently in NailPso compared to non-NailPso patients. Such a difference did not exist in the HV of the extensor tendons. Capillary microscopy showed a significant difference in the number of torsions/twist capillaries in NailPso compared to PsO patients without NailPso. Hypervascularization of the nail matrix is seen significantly more frequently in patients with psoriasis of the nail than in patients without psoriasis of the nail. Such a difference does not exist in DIP joint -extensor tendon- enthesitis. At the same time, torsions are significantly more frequently seen in capillary microscopy in NailPso than in patients without NailPso.

Conclusion: The US-PD examination is a simple and non-invasive procedure which can be performed routinely in daily practice. The hypervascularization of the nail matrix should also make one think of nail psoriasis in the early stage of PsO, in order to be able to start early an appropriate therapy for this very stigmatizing and therapeutically extremely difficult manifestation of PsO. It seems to occur independently of extensor tendon synovitis as an independent manifestation phenomenon.

The occurrence of torsions in capillary microscopy >50 % also seems to be groundbreaking for a NailPso, whereby capillary microscopy is a temporal challenge in daily routine.

References: § The present study (7734-BO-S2018 Ethics Commission of the MHH, Medical School Hannover, Germany) contains parts of the PhD thesis of M. Töllner

Disclosure of Interests: Becker-Capeller Detlef Grant/research support from: Novartis, Speakers bureau: Pfizer, Soham El-Nawab-Becker: None declared, Malo Toellner: None declared, Andreas Kleinheinz: None declared, Torsten Witte: None declared

DOI: 10.1136/annrheumdis-2020-eular.186

AB1089

EVALUATION OF JOINT RHEUMATOLOGY/ RADIOLOGY MDT OUTCOMES & THEIR IMPACT ON RHEUMATOLOGY SERVICE

A. Ediriweera¹, W. P. Lee¹, M. Steward¹, O. Savanovic-Abel¹. ¹University Hospital Plymouth NHS Trust, Plymouth, United Kingdom

Background: Multidisciplinary team (MDT) discussion between rheumatology and radiology is vital in diagnostic and prognostic management of patients' outcome. Nevertheless, discrepancies of the radiology report and clinical history cause unnecessary confusion and distresses to clinicians and patients. This could potentially affect subsequent clinical management.

Objectives: This survey was aimed to evaluate outcomes of rheumatology / radiology MDT discussion and to identify any discrepancies between original reports of the radiology images and amended reports following MDT. We also looked for potential reasons for the discrepancies and their impact on patient and health care resources due to erroneous original reporting.

Methods: We looked at all types of images which were discussed in rheumatology/ radiology MDT of University Hospital Plymouth from October 2016 to