Table 1. Adjusted R^2 of the models and indices. R^2_m : adjusted R^2 of the models; R^2_i : adjusted R^2 of the index included in the model; LR: Likelihood-Ratio test comparing HUPI models vs other indices¹.

OUTCOME	INDEX	ACT-RAY		LR	PEARL		LR
		R^2_m	R^2_i		R^2_m	R^2_i	
RADIOGRAPHIC PROGRESSION	HUPI	0.025	0.024	ref	0.110	0.020	ref
	DAS28	0.031	0.030	<0.000	0.102	0.011	<0.000
	SDAI	0.051	0.050	<0.000	0.109	0.019	<0.000
HAQ	HUPI	0.353	0.323	ref	0.477	0.448	ref
	DAS28	0.329	0.298	<0.000	0.472	0.442	<0.000
	SDAI	0.334	0.303	<0.000	0.486	0.457	<0.000
IL6	HUPI	-	-	-	0.212	0.184	ref
	DAS28	-	-	-	0.204	0.176	<0.000
	SDAI	-	-	-	0.201	0.172	<0.000

Figure 1. Box plots comparing the adjusted R^2 of each index estimated in models for HAQ in ACT-RAY.

Figure 2. Curves for comparison of predicted serum IL6 using fractional polynomials for HUPI and DAS28.

Conclusion: Although all indices explained the outcomes' variability similarly, HUPI did it better than DAS28 and SDAI for almost all outcomes except for Δ Genant and HAQ in PEARL.

References:

- [1] Gonzalez-Alvaro I, et al. PLoS ONE 14(4): e0214717

Disclosure of Interests: Sebastian C Rodriguez-García Speakers bureau: Novartis Farmaceutica, S.A., Merck Sharp & Dohme España, S.A., Sanofi Aventis, UCB Pharma, Nuria Montes: None declared, José Ivorra: None declared, Ana Triguero-Martinez: None declared, Luis Rodríguez Rodríguez: None declared, Isabel Castrejón: None declared, Loreto Carmona Grant/research support from: Novartis Farmaceutica, SA, Pfizer, S.L.U., Merck Sharp & Dohme España, S.A., Roche Farma, S.A, Sanofi Aventis, AbbVie Spain, S.L.U., and Laboratorios Gebro Pharma, SA (All through institution), Isidoro González-Alvaro Grant/research support from: Roche Laboratories, Consultant of: Lilly, Sanofi, Paid instructor for: Lilly, Speakers bureau: Abbvie, MSD, Roche, Lilly

DOI: 10.1136/annrheumdis-2020-eular.713

AB1253

EVALUATION OF CYSTEINE-RICH 61 IN RHEUMATOID ARTHRITIS AS A DIAGNOSTIC MARKER AND PREDICTOR OF ATHEROSCLEROSIS

H. A. Esaily¹, D. M. Serag², M. S. Rizk³, A. Sonbol⁴, D. Salem¹. ¹Menoufia University, Egypt, Physical Medicine, Rheumatology and Rehabilitation Department, Faculty of Medicine, Shebin El Kom, Egypt; ²Menoufia University, Egypt, Radiology Department, Faculty of Medicine, Shebin El Kom, Egypt; ³Menoufia University, Egypt, Medical Biochemistry and Molecular Biology Department, Faculty of Medicine, Shebin El Kom, Egypt; ⁴Menoufia University, Egypt, Clinical Pathology Department, Faculty of Medicine, Shebin El Kom, Egypt

Background: Matricellular protein Cysteine-rich protein 61 (Cyr61) is involved in chronic inflammatory disorders like rheumatoid arthritis (RA) and atherosclerosis.

Objectives: This study aimed to assess the value of serum Cyr61 in diagnosis of rheumatoid arthritis, evaluating its correlation with disease activity and its relation to atherosclerosis.

Methods: Cross-sectional study included 105 RA patients classified into active and inactive groups according to disease activity score (DAS28) with 50 healthy age and gender-matched controls. Full clinical and laboratory assessment was done including enzyme-linked immunosorbent assay (ELISA) measurement of Cyr61. Bilateral assessment of carotid intima-media thickness (CIMT) was done using high resolution-ultrasonography. Comparison of Cyr61 between RA patients and controls, correlation between Cyr61 and disease activity and CIMT were analyzed with appropriate statistical analyses.

Results: Significant elevation of Cyr61 in RA patients compared to controls (235.62±62.5 vs. 73.11±18.2) respectively. The cut off value of Cyr61 was 99.25 pg/ml, with area under the curve (AUC) =0.995, P <0.001, 98 % sensitivity and 95% specificity. Cyr61 was inversely correlated with DAS28 and its components in RA patients (r=-0.92, r=-0.94) (p<0.001). There was a significant positive correlation between Cyr61 levels and CIMT in inactive and active RA patients (r=0.88, r=0.47) respectively.

Conclusion: Serum Cyr61 as a potential diagnostic biomarker in RA is inversely correlated with disease activity. High Cyr61 in RA is a risk factor for atherosclerosis. Disruption of serum Cyr61 is engaged in the pathogenesis of both rheumatoid arthritis and atherosclerosis which is a clue for a future treatment strategy of RA.

References:

- [1] Smolen JS, Aletaha D, Barton A, Burmester GR, Emery P, Firestein GS, Kavanaugh A, McInnes IB, Solomon DH, Strand V, Yamamoto K (2018) Rheumatoid arthritis. Nature reviews Disease primers 4:18001. doi:10.1038/nrdp.2018.1
- [2] Pelechas E, Kaltsonoudis E, Voulgari PV, Drosos AA (2019) Rheumatoid Arthritis. In: Pelechas E, Kaltsonoudis E, Voulgari PV, Drosos AA (eds) Illustrated Handbook of Rheumatic and Musculo-Skeletal Diseases. Springer International Publishing, Cham, pp 45-76. doi:10.1007/978-3-030-03664-5_3
- [3] Sparks JA (2019) Rheumatoid Arthritis. Annals of Internal Medicine 170 (1):ITC1-ITC16. doi:10.7326/aitc201901010
- [4] Abd El-Monem S, Ali A, Hashaad N, Bendary A, Abd El-Aziz H (2019) Association of rheumatoid arthritis disease activity, severity with electrocardiographic findings, and carotid artery atherosclerosis. Egyptian Rheumatology and Rehabilitation 46 (1):11-20. doi:10.4103/err.err_36_18
- [5] Rawla P (2019) Cardiac and vascular complications in rheumatoid arthritis. Reumatologia 57 (1):27-36. doi:10.5114/reum.2019.83236
- [6] de Brito Rocha S, Baldo DC, Andrade LEC (2019) Clinical and pathophysiological relevance of autoantibodies in rheumatoid arthritis. Advances in Rheumatology 59 (1):2. doi:10.1186/s42358-018-0042-8

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.752

AB1254

PHENOTYPING OF MULTIPLE BIOFLUIDS FOR LIQUID BIOMARKERS FOR DIAGNOSTICS AND PERSONALIZED MEDICINE OF RHEUMATOID ARTHRITIS, SPONDYLOARTHRITIS AND OSTEOARTHRITIS

T. B. G. Poulsen¹, J. S. Andersen¹, M. K. Kristiansen¹, S. Rasmussen², L. Arent-Nielsen¹, C. H. Nielsen³, A. Stensballe¹. ¹University of Aalborg, Aalborg, Denmark; ²Aalborg University Hospital South, Aalborg, Denmark; ³Rigshospitalet, København, Denmark

Background: Inflammatory and autoimmune diseases include multifactorial pathomechanisms and systemic responses. The etiology of the joint diseases rheumatoid arthritis (RA), spondyloarthritis (SpA) in relation to osteoarthritis (OA) remain incomplete and establishing the correct diagnose remains nontrivial. Advances in high-throughput molecular technologies have increased investigations into the