In the group of patients treated with physical rehabilitation (n=33), a significant improvement of the VAS of pain and of the physical mean score of SF36 was noted only on day 18 (Table 2). A significant improvement of the mental mean score of SF 36 was noted on day 18 and 6 months (Table 2).

Table 2. Tendencies of the clinical indexes after treatment by physical rehabilitation

	D0 (n=37)	D18 (n=33)	3 months (n=28)	6 months (n=29)	12 months (n=25)
VAS ± SD	64 ± 19,5	47,4 ± 24,6	55,7 ± 18,1	50,5 ± 27,5	62,2 ± 22,2
P		0,002	0,091	0,057	0,567
ODI ± SD	34,3 ± 12,3	31,1 ± 17,3	31,9 ± 15,6	30,3 ± 18,8	31,1 ± 12
P		0,268	0,258	0,296	0,150
SF 36 physical mean score ± SD	38,1 ± 16,1	53,8 ± 22	42,6 ± 16,4	45,1 ± 18,7	42,6 ± 16,2
P		0,002	0,284	0,202	0,185
SF36 mental mean score ± SD	39,3 ± 17,5	61,6 ± 23,5	45,1 ± 18,9	49,6 ± 24,6	46,9 ± 22,1
P		0,000	0,130	0,033	0,09
Schober test ± SD	3,2 ± 1,4	3,5 ± 1,3	3,4 ± 1,4	2,7 ± 1,2	2,9 ± 1,3
P		0,845	0,983	0,091	0,346

VAS: visual analog scale; ODI: Oswestry Disability Index; SF36: short form 36

Comparison of the two types of tested treatments showed a certain number of statistically significant differences. The VAS and physical mean score of SF 36 at day 18 and mental mean score of SF-36 at 3 months were significantly lower on the thermal cure group, while schober test at 3 months was significantly lower in the physical rehabilitation group.

Conclusion: Our study suggests a superiority of thermal cure in relieving pain and improving quality of life at short term among patient with LBP and a comparable efficacy in improving lumbar spine function. However, the effectiveness of physical rehabilitation was greater on spine mobility.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.2212

AB1261 DOUBLE-BLIND PLACEBO-CONTROLLED TRIAL OF PULSED ELECTROMAGNETIC FIELD THERAPY IN KNEE OSTEOARTHRITIS (PRELIMINARY RESULTS)

D. Karateev¹, A. Makevnina¹, A. Tangieva¹, E. Luchikhina¹, H. Hamhoeva¹.

¹Moscow Regional Research and Clinical Institute ("MONIKI"), Rheumatology Department, Moscow, Russian Federation

Background: Pulsed electromagnetic field (PEMF) therapy is widely used in different areas of medicine. There are a lot of portable PEMF therapeutic devices on the market around the world. Nevertheless, the role of PEMF in treatment of rheumatic conditions is not clear. Current evidence is of low and very low quality.

Objectives: To study efficacy and safety of PEMF therapy in primary and secondary osteoarthritis (OA) of the knee in controlled clinical trial.

Methods: This abstract presents the preliminary results of an ongoing double-blind placebo-controlled trial of PEMF portable therapeutic device "ALMAG+" (Certificate EN ISO 13485:2012+AC:2012 reg.-No 44221 117836, Yelatma Instrument-Making Enterprise, Russian Federation, Reg. Num.: 3007075140). The device is intended for physiotherapeutic treatment and rehabilitation with a low-frequency low-intensity PEMF in medical institutions, as well as at home after the recommendation of a doctor. Patients with primary and secondary (as a part of inflammatory rheumatic disease) OA of knee with Kellgren-Lawrence Grade I-III included in the study (in patients with inflammatory conditions disease activity should be minimal on stable drug therapy). Three courses of PEMF of 20 procedures for 1 year planned in active treatment group and placebo (inactive device) group of 35 patients each. Efficacy is evaluated by pain VAS, WOMAC, Lequesne index, and quality of life studied using SF-36, EuroQoL 5D tools. Instrumental control with knee ultrasound and MRI investigations will be performed. The study protocol was approved by local Ethical Committee.

Results: To date 23 patients (7 males, 16 females, age 54,6±11,2 years, primary knee OA – 16 pts, RA- 6 pts, AS – 1 pt) completed 1st course of PEMF. Table presents differences (Δ) in the main clinical parameters in active treatment and placebo device groups. No significant difference in ESR or CRP levels was found. No treatment-related adverse events has been reported.

Table. Differences (Δ) in the main clinical parameters after a 1st course of PEMF.

Parameter	Active device (n=11)	Placebo device (n=12)	p
Δ VAS pain in movement	15 [0; 38,5]	5 [1,25; 10,5]	0,053
Δ VAS pain in rest	10 [0; 34]	1 [0; 2,75]	0,043
Δ WOMAC	3 [2; 10]	2 [0; 4,5]	0,174
Δ Lequesne index	3 [0; 4]	1 [0,25; 2,5]	0,258

Conclusion: In this preliminary analysis pulsed electromagnetic field (PEMF) therapy showed significant impact on pain in rest in knee OA after one course of procedures.

References: Negm A, Lorbergs A, Macintyre NJ. Efficacy of low frequency pulsed subsensory threshold electrical stimulation vs placebo on pain and physical function in people with knee osteoarthritis: systematic review with meta-analysis. Osteoarthritis Cartilage. 2013 Sep;21(9):1281-9. doi: 10.1016/j.joca.2013.06.015

Disclosure of Interests: Dmitry Karateev Consultant of: Abbvie, Pfizer, Biocad, Sanofi, Novartis, Lilly, Speakers bureau: Abbvie, Roche, Pfizer, Biocad, MSD, Sanofi, Johnson & Johnson, Glaxo, UCB, Celgene, Novartis, Lilly, Bayer, Alexandra Makevnina Speakers bureau: Sanofi, Aminat Tangieva: None declared, Elena Luchikhina Consultant of: Abbvie, Biocad, Sanofi, Celgene, Speakers bureau: Abbvie, Roche, Pfizer, Biocad, MSD, Sanofi, Johnson & Johnson, Glaxo, UCB, Celgene, Novartis, Hava Hamhoeva: None declared

DOI: 10.1136/annrheumdis-2020-eular.6396

AB1262 EFFECTS OF TAI CHI EXERCISE ON PAIN, FUNCTIONAL STATUS AND QUALITY OF LIFE IN PEOPLE WITH OSTEOARTHRITIS AND INFLAMMATORY ARTHRITIS

N. Kalebota¹, M. Milošević², N. Laktašić Žerjavić¹, A. Andreić³, B. Končar⁴, M. Vedriš⁴, P. Turković⁴, N. Žura¹, K. Kovač Durmiš¹, I. Žagar¹, P. Perić¹.

¹Clinical Hospital Centre Zagreb, University Department for Rheumatology and Rehabilitation, Zagreb, Croatia; ²University of Zagreb, School of Medicine, Andrija Štampar School of Public Health, Zagreb, Croatia; ³Patient Advocacy Association, Zagreb, Croatia; ⁴Croatian Wushu Federation, Zagreb, Croatia

Background: Tai Chi as a type of physical activity (PA) is included in recommendations for PA in people with inflammatory arthritis (IA) and osteoarthritis (OA).¹

Objectives: The aim of this study was to evaluate the effect of Tai Chi exercise program on pain, functional status and quality of life in people with OA and IA.

Methods: A study co-funded by the Erasmus + European Union Program for Sporting Activities called #WushuElixir was conducted in Croatia, Slovenia, Belgium and Spain. We present the results of the research conducted in Croatia from October 2018 to April 2019. A total of 44 participants (Pts) entered the Tai Chi exercise program. Majority of Pts were female with OA (Table 1). Tai Chi exercise program was performed for 60 minutes, twice a week, with a total of 50 training sessions. In data analysis where included only those Pts who completed at least 36% of exercise program, a total of 28 (63.6%) Pts, out of which 24 (85.7%) completed at least 70% of exercise program. An average attendance was 39 (78%) sessions. Outcome measures were: pain (VAS scale), stability and balance (Functional reach test; FRT), chest mobility (breathing index), spinal mobility (cervical and thoracic sagittal indices and Schober's test), muscle strength (Wall sit test), physical function (Lequesne Index of severity for OA of the hip, and Stanford Health Assessment Questionnaire (HAQ) 20-Item for IA) and quality of life (15D® The health-related quality of life instrument; 15D QL). For statistical analysis t-test for paired samples was used with statistical significance set on P<0.05.

Table 1. Descriptive statistics of participants

		N	%
Gender	Male	4	14.3%
	Female	24	85.7%
Age groups	≤45 years	3	10.7%
	46-60 years	9	32.1%
	61-70 years	11	39.3%
	≥71 years	5	17.9%
		5	17.9%
Diagnosis	Osteoarthritis of the spine, hip and knee	19	67.9%
	Inflammatory arthritis	5	17.9%
	Other	4	14.3%

Results: There was significant improvement in breathing index (P<0.001), sagittal mobility of cervical (P<0.001) and thoracic spine (P=0.007), Schober's

test ($P<0.001$), Wall sit test ($P=0.042$), 15D QL ($P=0.048$) and Lequesne index ($P=0.030$). There were no significant changes in VAS pain level ($P=0.651$), FRT value ($P=0.442$) and Stanford HAQ-20 scale result ($P=0.186$).

Conclusion: Tai Chi can improve functional status and quality of life in people with OA and IA.

References:

- [1] Rausch Osthoff AK, Niedermann K, Braun J, et al. 2018 EULAR recommendations for physical activity in people with inflammatory arthritis and osteoarthritis. *Ann Rheum Dis*. 2018;77(9):1251-1260.

Acknowledgments: Goran Pažin, Tatjana Božac, Damir Grbac, Kristina Spajić, Ana Karić.

Disclosure of Interests: Nataša Kalebota Speakers bureau: Krka, Milan Milošević: None declared, Nadica Laktašić Žerjavić Speakers bureau: Abbvie., Roche, MSD, Eli Lilly, Pfizer, Mylan, Amgen, Fresenius Kabi, Antun Andreić: None declared, Biljana Končar: None declared, Mladen Vedriš: None declared, Petar Turković: None declared, Nikolino Žura: None declared, Kristina Kovač Durmiš Speakers bureau: Abbvie, Fresenius Kabi, Iva Žagar Speakers bureau: Abbvie, Roche, MSD, Eli Lilly, Pfizer, Porin Perić: None declared

DOI: 10.1136/annrheumdis-2020-eular.5843

AB1263

THE EFFICACY OF PULMONARY REHABILITATION IN PATIENTS WITH ANKYLOSING SPONDYLITIS

N. Sahin¹, N. Sanoğlu², B. Görcan¹. ¹Balikesir University, Physical Medicine and Rehabilitation, Balikesir, Turkey; ²Balikesir University, Chest Diseases, Balikesir, Turkey

Background: Ankylosing spondylitis (AS) is a rheumatic disease that can cause a restrictive lung condition. Pulmonary rehabilitation increases lung capacity in patients with lung disease. In this study, we aimed to show increase in exercise capacity, decrease in dyspnea and change in quality of life with pulmonary rehabilitation in patients with AS.

Objectives: In this study, we included 20 AS patients of whom admitted to our outpatient clinic. At the beginning and end of the study, patients some findings such as SFT, FEV1 values, saturations, arterial blood pressures and heart rates were recorded separately. Also, BODE index results, 6-minute walk test (MWT) values, BORG dyspnea scores, Modified Medical Research Council (MMRC) dyspnea scales, Anxiety Stress Scales (ASS) were examined. Short Form-36 (SF-36) questionnaires were filled out.

Methods: Patients underwent rehabilitation for 8 weeks, two days in a week at hospital and one day at home. According to their exercise tolerances in rehabilitation, patients were given upper and lower limb endurance training, strength training, flexibility and stretching exercises, balance exercises, inspiratory muscles training.

Results: In our study, patients with AS showed significant improvement in MMRC, borg dyspnea scales and 6 MWT at the end of rehabilitation compared to beginning of rehabilitation ($p<0.05$). It is shown that there was significant improvement in the values of PEF % and FVC%, when the difference averages before and after rehabilitation were compared ($p < 0.05$) but there was no difference in FEV1 %, FEV1/FVC values ($p > 0.05$). There was also significant improvement in BASDAI-BASFI values ($P=0.006$, $P=0.016$, respectively). The bode index showed no significant difference ($p > 0.05$). There was significant improvement in anxiety on ASS. Only the mental parameter showed significant improvement when parameters evaluated by the SF-36 survey ($p < 0.05$). There was no significant difference in the rest of all parameters ($p>0.05$).

	Before treatment n=20 (Mean ±SD)	After treatment n=20(Mean ±SD)	p*
FVC	84,35±13,92	86,40±13,42	,007
FEV1	84,50±13,03	85,30±11,13	,431
FEV1 FVC	105,30±9,57	85,30±11,13	,837
PEF	64,80±17,14	67,60±16,65	,003
6-Minute Walk Test	415,5500±124,88625	439,900±122,364	,000
MMRC Dyspnea Scale	1,350±1,136	1,150±0,87509	,046
BORG Dyspnea Score	2,400±1,273	1,650±0,988	,000
BODE Index	,7500±1,86025	,700±1,688	,317
BASDAI	3,622±1,837	3,140±1,392	,006
BASFI	4,350±1,692	6,065±8,638	,016
ASS	9,00±5,036	7,050±3,034	,006
SF36			
General Health	47,000±15,165	49,125±14,171	,485
Energy	47,000±20,026	52,000±18,020	,120
SocialFunctioning	55,000±22,725	58,875±18,217	,129
Role limitations due to emotional problems	47,765±46,185	64,515±36,463	,117
Mental	52,450±19,527	62,800±17,572	,041

P* Wilcoxon signed rank test. n, number of participants; SD, standard deviation; FVC,forced vital capacity;FEV1,forced expiratory volume;PEF,pik expiratory volume; MMRC, Modified Medical Research Council;O2,oxygen; BASDAI,Bath Ankylosing Spondylitis Disease Activity Index; BASFI,Bath Ankylosing Spondylitis Disease Activity Index;SF-36, Short Form-36.

Conclusion: After pulmonary rehabilitation in patients with AS dyspnea symptoms decreased. As a result of this study, it is thought that pulmonary rehabilitation applied to patients with AS can minimize the adverse effects of the disease over the respiratory system.

Key words: pulmonary rehabilitation, ankylosing spondylitis, respiratory function test

References:

- [1] Zão A, Cantista P.The role of land and aquatic exercise in ankylosing spondylitis: a systematic review. *Rheumatol Int*. 2017; 37: 1979-1990.
- [2] Fuschillo S, De Felice A, Martucci M et al. Pulmonary Rehabilitation Improves Exercise Capacity in Subjects with Kyphoscoliosis and Severe Respiratory Impairment. *Respir Care*. 2015; 60(1): 96-101.
- [3] Spruit MA, Pitta F, McAuley E et al. Pulmonary Rehabilitation and Physical Activity in Patients with Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med*. 2015; 15;192(8): 924-33.
- [4] Drăgoi RG, Amarica E, Drăgoi M et al. Inspiratory muscle training improves aerobic capacity and pulmonary function in patients with ankylosing spondylitis: A randomized controlled study. *Clin Rehabil*. 2016; 30(4):340-346.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.2299

AB1264

THE EFFECTIVENESS OF KINESIOTHERAPY FOR PAIN SHOULDER SYNDROME

O. Tikhoplav¹, E. Guryanova^{1,2}, O. Moskovskaya¹, A. Moskovskij¹. ¹Chuvash State University, Cheboksary, Russian Federation; ²Institute for Advanced Medical Studies, Cheboksary, Russian Federation

Background: Pain shoulder syndrome is one of the most common conditions that is inherent in patients with various diseases and conditions. This problem is faced by both patients of a rheumatological profile (rheumatoid arthritis), and patients of other profiles: neurological (stroke, hernia of the cervical spine), traumatological (periartthritis). The regular use of painkillers and NSAID is associated with a risk of serious adverse reactions in patients.

PNF therapy can have a fairly quick positive effect. Kinesiotaping is a common and easily accessible treatment method that, due to the inclusion of local neuromuscular adaptation mechanisms, can have an analgesic effect.

Objectives: The purpose of the study is to compare the effectiveness of traditional treatment methods in combination.

Methods: Patients with pain shoulder syndrome who were admitted to the doctor of physical and rehabilitation medicine were randomly assigned to one of three groups: group G1 - 20 patients who received combined treatment, including massage, acupuncture, PNF, kinesiotherapy.

Patients of group G2 (20 people) who received massage and acupuncture sessions, kinesiotherapy, but did not deal with PNF.

Patients in group G3 (20 people) received massage, acupuncture, and PNF, excluding kinesiotaping.

All patients received treatment 5 times a week (Monday to Friday, excluding Saturday and Sunday), the total duration of the rehabilitation course is 20 days. The program included: 30 minutes an acupuncture session, 20 minutes massage, 45 minutes physical exercises with a physical therapist. Classes using the PNF technique were carried out separately every other day for 45 minutes using the standard method for shoulder pain.

VAS was used to monitor efficacy. Pain was assessed at the beginning of the study, with the third, sixth, ninth and last visit to the clinic.

Results: The age composition of the patient group is from 31 to 70 years. The gender composition is 35 (58.33%) women, and 25 (41.67%) men.

The average time between the onset of clinical symptoms and the first treatment session was 47.3 days. In 90% of patients, a history of pain lasted from 6 weeks or more. 52 patients (86.67%) completed the treatment completely. Of these, G1 is 19 people (90% of this group), 16 (80%) in G2, and 17 of 20 in the G3 group (85%).

The average VAS score in G1 was 5.15 at the start of the study, and after the tenth session, it dropped to 2.78. A significant decrease in pain is also observed in the G2 group (from 5.17 at the beginning of the study to 3.19 after the tenth session). The G3 group in terms of pain reduction almost equaled the G1 group, where the level of pain from 5.16 decreased to 2.71.

Patients within 6 months after treatment evaluated the level of pain on their own and reported the data to the doctor. In patients of group G1, after 6 months, the average VAS score is 3.71. G3 patients after 6 months, the VAS score is 4.25 points. Patients of the G2 group at 3 months noted a return of the pain syndrome, after 3 months, the VAS score was 5.11.