Table 1. Distribution of KL grades in the training and testing sets.

KL grades	0	1	2	3	4	Total
Training set	10893	4582	6114	3320	799	25,978
	41.9%	18.7%	23.5%	12.8%	3.1%	
Testing set	2472	1353	1696	775	194	6,490
	38.1%	20.8%	26.1%	11.9%	3.0%	

Table 2. Performance matrices of the Deep-Ten and ResNet18 models to detect osteoarthritis

	Deep-TEN	ResNet18
Sensitivity	62.29% (95% CI, 60.42%–64.13%)	59.14% (95% CI, 57.24%–61.01%)
Specificity	90.07% (95% CI, 89.07%–91.00%)	94.09% (95% CI, 93.30%–94.82%)
Positive predictive value	81.37% (95% CI, 79.81%–82.84%)	87.46% (95% CI, 85.96%–88.82%)
Negative predictive value	77.42% (95% CI, 77.64%–79.65%)	76.77% (95% CI, 75.93%–77.59%)

Conclusion: This study demonstrates that the bone texture model performs reasonably well to detect radiographic osteoarthritis with a similar performance to the bone contour model.

References:

- [1] Bertalan Z, Ljuhar R, Norman B, et al. Combining fractal- and entropy-based bone texture analysis for the prediction of osteoarthritis: data from the multicenter osteoarthritis study (MOST). *Osteoarthritis Cartilage* 2018;26:S49.
- [2] Lindner C, Wang CW, Huang CT, et al. Fully Automatic System for Accurate Localisation and Analysis of Cephalometric Landmarks in Lateral Cephalograms. *Sci Rep* 2016;6:33581.
- [3] Zhang H, Xue J, Dana K. Deep TEN: Texture Encoding Network. The IEEE Conference on Computer Vision and Pattern Recognition (CVPR) 2017:708-17.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.2858

OP0063

QUANTITATIVE COMPUTED TOMOGRAPHY PREDICTS 10-YEAR MORTALITY IN INTERSTITIAL LUNG DISEASE RELATED TO SYSTEMIC SCLEROSIS

A. Ariani¹, E. Bravi², M. De Santis³, V. Hax⁴, S. Parisi⁵, F. Lumetti⁶, F. Girelli⁷, M. Saracco⁸, F. De Gennaro⁹, A. Giollo¹⁰, M. Abdel Jaber¹¹, F. Bozzao¹², M. Silva¹³, M. C. Ditto⁵, C. Lomater⁸, F. Mozzani¹, D. Santilli¹, E. Di Donato¹, A. Beccioli¹, F. Pucciarini², L. Canziani³, F. C. Bodini¹⁴, E. Arrigoni², M. Bredemeier¹⁵, R. Mendonça Da Silva Chakr⁴, A. Spinella⁶, L. Idolazzi¹⁰, R. Bortolotti¹¹, P. Tomietto¹², E. Baratella¹⁶, S. Tollot¹⁶, D. Giuggioli⁶, F. Fischetti¹², E. Fusaro⁵, N. Sverzellati¹³, C. A. Scirè¹⁷. ¹AOU, Internal Medicine and Rheumatology Unit, Parma, Italy; ²Ospedale G. Da Saliceto, Internal Medicine and Rheumatology Unit, Piacenza, Italy; ³Humanitas Clinical and Research Center, Rheumatology and Clinical Immunology, Rozzano, Italy; ⁴University of Rio Grande do Sul, Hospital de Clínicas, Division of Rheumatology, Porto Alegre, Brazil; ⁵AOU Città della Salute e della Scienza, Rheumatology Department, Turin, Italy; ⁶University of Modena and Reggio Emilia, AOU, Policlinico, Rheumatology Unit, Modena, Italy; ⁷Ospedale GB Morgagni - AUSL, Internal Medicine Unit-Rheumatology Service, Forlì, Italy; ⁸Ospedale Mauriziano - Umberto I, Rheumatology Unit, Turin, Italy; ⁹AO "Istituti Ospitalieri", Rheumatology Unit, Cremona, Italy; ¹⁰University of Verona, AO, Rheumatology Unit, Verona, Italy; ¹¹Ospedale Santa Chiara, Rheumatology Unit, Trento, Italy; ¹²University of Trieste, Cattinara Teaching Hospital, Department of Medical Sciences, Trieste, Italy; ¹³University of Parma, Unit of Surgical Sciences, Section of Radiology, Parma, Italy; ¹⁴Ospedale G. Da Saliceto, Radiology Unit, Piacenza, Italy; ¹⁵Hospital Nossa Senhora da Conceição, Rheumatology Service, Porto Alegre, Brazil; ¹⁶ASUI, Department of Radiology, Trieste, Italy; ¹⁷University of Ferrara - AOU Sant'Anna, Section of Rheumatology, Ferrara, Italy

Background: Interstitial lung disease (ILD) is the main cause of death in Systemic Sclerosis (SSc). Chest CT is the gold standard in detecting ILD although it is not easy to understand its prognostic value. ILD qualitative assessment is almost worthless. Goh et al. semi quantitative score of ILD extent is related to mortality risk but it is burdened by relevant inter/intra-readers variability. An operator independent algorithm based on voxel-wise analysis proved to identify SSc patients with an increased risk of mortality according to prediction models.

Objectives: To verify if quantitative analysis of chest CT (QCT) predict 10 years-mortality in SSc patients.

Methods: SSc patients with availability of a chest CT were enrolled in 13 different centers. The CT voxel-wise analysis with a free software (www.horosproject.com) provided QCT indexes: kurtosis, skewness, mean lung attenuation and standard deviation. Patients characteristics, autoimmune profile and pulmonary function test were collected. The follow-up interval lasted from the date of chest CT to the one of the last visit or death. Each QCT index cutoff, established in a previous study (1), clustered patients in two groups. Kaplan-Meier analysis estimated and compared survival in the above mentioned groups. $p < 0.05$ was considered statistically significant.

Results: Five hundred sixty three SSc patients were enrolled (35938 patient-months); 52.4% had ILD detectable at CT scan. For each QCT index cutoff the cohort was split in two subgroups without differences in terms of sex, age, disease duration, autoimmune profile. All QCT indexes' cutoff selected subgroups with statistically different survival rate (e.g in Figure 1).

Figure 1

Conclusion: QCT can arise as the new gold standard in identifying SSc patients with poor prognosis. The real possibility to stratify SSc subjects according mortality risk will have a pivotal role in ILD treatment decisional process with the incoming anti-fibrotic drugs.

References:

- [1] Ariani A et al. *Rheumatology* 2017

Disclosure of Interests: Alarico Ariani: None declared, Elena Bravi: None declared, Maria De Santis: None declared, Vanessa Hax: None declared, Simone Parisi: None declared, Federica Lumetti: None declared, Francesco Girelli: None declared, Marta Saracco: None declared, Fabio De Gennaro: None declared, Alessandro Giollo: None declared, Masen Abdel Jaber: None declared, Francesco Bozzao: None declared, Mario Silva: None declared, Maria Chiara Ditto: None declared, Claudia Lomater: None declared, Flavio Mozzani: None declared, Daniele Santilli: None declared, eleonora Di Donato: None declared, Andrea Beccioli: None declared, Sanofi-Genzyme, UCB and AbbVie, Francesco Pucciarini: None declared, Lorenzo Canziani: None declared, Flavio Cesare Bodini: None declared, eugenio arrigoni: None declared, M Bredemeier: None declared, Rafael Mendonça da Silva Chakr: None declared, Amelia Spinella: None declared, Luca Idolazzi: None declared, Roberto Bortolotti: None declared, Paola Tomietto: None declared, Elisa Baratella: None declared, Saverio Tollot: None declared, Dilia Giuggioli: None declared, Fabio Fischetti: None declared, Enrico Fusaro: None declared, Nicola Sverzellati: None declared, Carlo Alberto Scirè: None declared

DOI: 10.1136/annrheumdis-2020-eular.2239

OP0064

AUTOMATIC SCORING OF ARTHRITIS DISEASE ACTIVITY ON ULTRASOUND IMAGES FROM RHEUMATOID ARTHRITIS PATIENTS WITH CASCADED CONVOLUTIONAL NEURAL NETWORKS

A. Christensen¹, S. A. Just², J. K. H. Andersen¹, T. R. Savarimuthu¹. ¹University of Southern Denmark, Maersk Mc-Kinney Møller Institute, Odense, Denmark; ²Odense University Hospital, Svendborg Hospital - Department of Medicine, Section of Rheumatology, Svendborg, Denmark

Background: Systematic Power or Color Doppler (CD) ultrasound (US) of joints can be used for early detection of Rheumatoid Arthritis (RA), predicting radiographic progression and early detection of disease flare in established