Figure 2. Adherence to Clinical Guidelines (%) reported in the Clinical Vignette

Conclusion: Our findings reveal an overall sub-optimal adherence to OA-guidelines in a sample of Italian physiotherapists. Italian physiotherapists appear to be aware of the importance of exercise and education for patients suffering from OA. Instead, a sub-optimal level of knowledge was found regarding the criteria for OA clinical diagnosis and on the role of other non-surgical interventions in the management of OA. These results identified some gaps between evidence and clinical practice, which may lead to an incorrect management of patients suffering from OA.

References:

- [1] GBD 2017 Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392:1789–858.
- [2] Fernandes, L. et al. EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis. *Ann Rheum Dis*. 2013;72:1125–35.
- [3] Basedow M et al. Assessing appropriateness of osteoarthritis care using quality indicators: a systematic review. *J Eval Clin Pract*. 2015;21:782–9.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.987

THU0644-HPR TRENDS IN DIAGNOSIS AND TREATMENT OF FIBROMYALGIA AMONG MEXICAN PHYSICIANS.

R. I. Arvizu-Rivera¹, N. Escobedo-Zuñiga¹, I. J. Colunga-Pedraza², A. Cárdenas², G. Serna-Peña². ¹Hospital Universitario "Dr. Jose Eleuterio Gonzalez", Internal Medicine Department, Monterrey, Mexico; ²Hospital Universitario "Dr. Jose Eleuterio Gonzalez", Rheumatology Division, Monterrey, Mexico

Background: Fibromyalgia (FM) is a disease characterized by widespread pain which affects 2-8% of the population. Previous studies have shown lack of awareness of classification criteria and accompanying symptoms of FM, as well as heterogeneity in the management of these patients among both rheumatologist and non-rheumatologist physicians.

Objectives: The objective of this study was to explore the trends in diagnosis and management of fibromyalgia among general practitioners, family physicians, psychiatrists, neurologists and rheumatologists in northeastern Mexico

Methods: We designed an online survey to yield data on perception of FM, knowledge of existing classification criteria including the ACR 1990, ACR 2010, ACR 2010 modified and AATP classification criteria, as well as pharmacologic and nonpharmacologic therapy for the treatment of FM. Participants should have finished their residence at least in 2019. General practitioners, family physicians, psychiatrists, neurologists and rheumatologists were included.

Results: A total of 236 participants were included, most of the participants were general practitioners, 149 (59.3%). Other specialties included were rheumatologists 21 (8.9%), neurologists 18 (7.6%), psychiatrists 8 (3.4%), and family physicians 49 (20.8%). FM was considered a clinical entity by 208 (88.1%) participants. Participants' characteristics are shown in Table 1. Twenty-eight (11.9%) participants didn't know any classification criteria for FM, and 38 (16.1%) participants answered that they didn't use any classification criteria to make a formal diagnosis of FM. The 1990 ACR classification criteria was used the most, 62 (26.3%); closely followed by the 2010 modified ACR classification criteria, 61 (25.8%). A total of 101 (42.8%) participants made a formal diagnosis of FM in the previous year and 179 (75.8%) referred the patient to another specialist. Most patients were referred to a rheumatologist, 126 (53.4%). One-hundred and

fifty-eight (66.9%) participants believe rheumatologist should be the main care providers for patients with FM.

Table 1. Participants' characteristics.

Variable	
Age, median (25p-75p)	33 (27-38)
Women, n (%)	203 (86)
Rheumatologists, n (%)	21 (8.9)
General practitioners, n (%)	140 (59.3)
Public practice, n (%)	113 (47.9)
Public and private practice, n (%)	51 (21.6)
< 5 years of practice, n (%)	100 (42.4)
> 15 years of practice, n (%)	30 (12.7)

Conclusion: A total of 88.1% of physicians know at least one classification criteria for the diagnosis of FM. In Mexico, rheumatologists are considered the main care providers for patients with FM. Regarding therapies with level 1A evidence for efficacy in the treatment of FM, only cognitive behavioral therapy and patient education were used by more than 50% of physicians. Serotonin-norepinephrine reuptake inhibitor and tricyclic antidepressants (both of which have level 1A evidence) were only used by 20.8 and 29.2%, respectively.

References:

- [1] Fibromyalgia: a clinical review. *JAMA*. 2014 Apr 16;311(15):1547-55.

Table 2. Pharmacologic and non-pharmacologic treatment of fibromyalgia among physicians.

Variable	n (%)
Exercise (any kind), n (%)	91 (38.6)
Aerobic exercise (aerobic), n (%)	86 (36.4)
Cognitive behavioral therapy, n (%)	127 (53.8)
Patient education, n (%)	142 (60.2)
Gabapentinoids, n (%)	107 (45.3)
Tricyclic antidepressants, n (%)	69 (29.2)
Selective serotonin reuptake inhibitors, n (%)	121 (51.3)
Serotonin-norepinephrine reuptake inhibitor, n (%)	49 (20.8)
Muscle relaxers, n (%)	85 (36)
NSAID, n (%)	137 (58.1)
Opioids, n (%)	27 (11.4)
Mindfulness, n (%)	42 (17.8)

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.4052

THU0645-HPR ULTRASOUND EDUCATION FOR RHEUMATOLOGIST IN RUSSIA: REALITIES AND WHAT IS NEEDED TO IMPROVE.

N. Dudnikova¹, A. Mikhailova¹, O. Alekseeva¹, A. Demina¹, A. Volkov¹. ¹VA Nasonova Research Institute of Rheumatology, Moscow, Russia, Moscow, Russian Federation

Background: Musculoskeletal ultrasonography (MSUS) is a valuable imaging modality for the practicing rheumatologist. In everyday practice, physical examination is limited by its sensitivity and subjectivity of assessment. MSUS is inexpensive, available, and allows repeat examination as soon as necessary. Thus, MSUS has become the "third eye" of the rheumatologist, in that it allows more detailed examination of muscles, bones, and joints, just as the stethoscope provides further details about the respiratory and cardiovascular systems. Most ultrasound examinations in Russia, including abdominal and vascular sonography, echocardiography, are conducted by trained physician - radiologist, called "ultrasonography specialists". Nevertheless, it is very difficult to perform MSUS, mainly because standardized ultrasound protocols for same region are different for various diseases and often do not provide physicians with the information they need, and also may be mistake. The conduct of MSUS by rheumatologists themselves is limited to the need for 4-month radiology (ultrasound) certification. However, the number of rheumatologists conducting MSUS themselves is increasing. Training week small group courses are provided by VA Nasonova Research Institute of Rheumatology for the practicing rheumatologist and other physicians.

Objectives: We analyzed the training and awareness of the rheumatologists and radiologists in MSUS, as well as their commitment for participation in our courses.

Methods: We studied a one and a half-year period, from June 2018 to December 2019. During this time, 57 applications were submitted to participate in the courses. All participants who arrived were tested before starting the course. Test included 40 general questions about MSUS. Test results as well as commitment to training were analyzed in two groups: rheumatologists and radiologists. Non-parametric statistics were used for data processing.

Results: Among 57 applications submitted for the courses, 26 were from rheumatologists and 31 from radiologists, but the courses were attended by 13 rheumatologists and 26 radiologists (68%). Thus, commitment was reliably higher in radiologists (84%) than in rheumatologists (50%). Fisher's methods show high statistically significance ($\chi^2 = 7.51$, $p = 0.006$). Although the test focused on ultrasound, it was surprise for us that the median percentage of correct answers was higher in rheumatologists - 64 (54; 80)%, than radiologists ("ultrasonography specialists") 48 (40; 62)%. Difference also were significantly ($p = 0.04$).

Conclusion: Thus, despite better preparedness of rheumatologist, motivation to study MSUS prevails among radiologists. It is necessary to actively introduce MSUS into general rheumatologic educational programs in order to motivate rheumatologists to study MSUS. Changes in professional standard will also encourage wide use of MSUS by rheumatologists.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.6158

THU0646-HPR EXPLORING THE ROLE OF NURSE IN RHEUMATIC CARE AND FEASIBILITY OF ENHANCING TREAT TO TARGET STRATEGIES IN JAPAN.

M. Fusama^{1,2}, M. Uda³, H. Matsumura⁴, Y. Van Eijk-Hustings⁵, S. Oliver⁶, H. Nakahara². ¹Yukioka Hospital, Osaka, Japan; ²Osaka Yukioka College of Health Science, Osaka, Japan; ³Kyoto College of Nursing, Kyoto, Japan; ⁴Higami Internal Medicine Rheumatism and Diabetes, Nara, Japan; ⁵Department of Patient & Care/Clinical Epidemiology and Medical Technology Assessment (KEMTA), Maastricht, Netherlands; ⁶Susan Oliver Associates, London, United Kingdom

Background: The role of rheumatology nurses is considered important for the implementation of T2T [1]. For nurses' contribution to implementation of the T2T strategy, it is necessary to explore the nurses' opinion on their roles in real clinical practice.

Objectives: The aim of this study is to evaluate what is required for nurses to implement T2T in real clinical practice in Japan.

Methods: Registered nurses engaged in rheumatic care in clinical practice in Japan were enrolled. Focus group interviews were conducted exploring 'What is necessary for RA nurses to implement T2T' using semi-structured interviews. Data analysis was used with Krippendorff's content analysis method.

Results: 24 nurses (all females) from 10 hospitals were enrolled in this study. The results of the qualitative analysis were categorized in 10 main categories, derived from 37 subcategories based on 64 different codes: (1) provide basic knowledge of RA, (2) provide knowledge of RA drugs, (3) provide knowledge and skills of self-monitoring, (4) enhance self-efficacy and support self-management, (5) support decision-making, (6) psychological and social support, (7) understand the diversity and feelings of patients and their families, (8) support based on individual needs, (9) ensure continuing educational opportunities for nurses to enable the provision of high quality care, (10) collaborate with multi-disciplinary teams.

These categories are mostly covered in the contents of the 2018 updated EULAR recommendations for the role of nurses except "evidence-based rheumatic care", "telehealth" and "comprehensive participation in disease management".

Conclusion: These findings indicate the areas of exploration including further educational and training needs, attitudes and the professional scope for nurses to extend their roles to provide greater value to patient care.

In Japan, evidence-based RA nursing and telehealth systems have not yet been established. In addition, therapeutic intervention by nurses and nurse-led clinic are not permitted. Our results might reflect this situation and possibly elucidates the gap between EULAR's evidence-based recommendations and opinions of Japanese nurses working in daily clinical practice. As evidence-based nursing is considered to be crucial from both cost-effectiveness and improvement of patients QOL, this result also might shed light on what we need for future better rheumatic nursing in Japan.

References:

[1] van Eijk-Hustings Y, et al. Ann Rheum Dis 2012;71: 13–19.

Disclosure of Interests: MIE FUSAMA: None declared, Miyabi Uda: None declared, Harumi Matsumura: None declared, Yvonne van Eijk-Hustings Grant/

research support from: grand from sanofi and UCB, Consultant of: fee from amgen, Susan Oliver: None declared, Hideko Nakahara: None declared

DOI: 10.1136/annrheumdis-2020-eular.5315

THU0647-HPR SHARED DECISION MAKING AMONGST RHEUMATOLOGISTS IN TRAINING IN THE NETHERLANDS: WHERE DO WE STAND?

L. Kranenburg¹, M. Dankbaar², N. Basoski³, W. Van den Broek², J. Hazes¹.

¹Erasmus MC, Rheumatology, Rotterdam, Netherlands; ²Erasmus MC, Institute for Medical Education Research Rotterdam, Rotterdam, Netherlands;

³Maasstad hospital, Rheumatology, Rotterdam, Netherlands

Background: The training curriculum for rheumatologists in training in the Netherlands describes competences and entrusted professional activities (EPA) to monitor the progress in learning. However, this training program does not discuss training of Shared Decision Making. As the basis for shared care and patient participation is made during these years, the question arises how rheumatologist in training think about Shared Decision Making and how they use this in daily practice.

Objectives: Inventory of vision, experience and self-evaluation of skills related to Shared Decision Making amongst rheumatologists in training in the Netherlands in order to identify barriers in the implementation of Shared Decision Making in daily practice.

Methods: Qualitative data was collected from an online survey amongst rheumatologists in training who were registered in January 2018 by the Dutch Society of Rheumatology.

Results: Forty-two rheumatologists in training from various years of training responded (60%). Respondents think that Shared Decision Making is important. A third applies Shared Decision Making on a regular basis in daily practice. Self rating of skills for Shared Decision Making varies from sufficient to good. However, respondents are uncertain about their performance due to a lack of feedback and unclearness of the concept. They indicate that Shared Decision Making is not possible for all patients and find it difficult to assess whether the patient has a clear understanding of the options. Patient's preferences are discussed only by 33% of the doctors on a regular basis when starting new treatment.

Conclusion: Rheumatologists in training agree on the importance of Shared Decision Making, but are uncertain about their performance. Unclearness of the concept is described as a known barrier in literature^{1,2} and is frequently mentioned by respondents. Rheumatologist in training indicate that not all patients are fit for Shared Decision Making. Regarding the limited training on the subject this could also be a misjudgment of patients preferences and lack of experience how to deal with different patient types. There is a clear plea for more training and feedback on the subject. Training should be integrated in the curriculum focusing on how to assess patients preferences and how to apply Shared Decision Making also for patients who indicate to leave decisions up to their doctor.

References:

[1] van Veenendaal, H. et al. Accelerating implementation of shared decision-making in the Netherlands: An exploratory investigation. *Patient Educ Couns* 101, 2097-2104 (2018).

[2] Legare, F., Ratte, S., Gravel, K. & Graham, I. D. Barriers and facilitators to implementing shared decision-making in clinical practice: update of a systematic review of health professionals' perceptions. *Patient Educ Couns* 73, 526-535 (2008).

Disclosure of Interests: Laura Kranenburg Grant/research support from: Pfizer and UCB for the development of the Reuma App, a tool to support selfmanagement for patients. This is not used for the research related to the submitted abstract., Mary Dankbaar: None declared, Natalja Basoski: None declared, Walter Van den Broek: None declared, Johanna Hazes: None declared

DOI: 10.1136/annrheumdis-2020-eular.4142

THU0648-HPR STUDY OF PAIN ATTITUDES & BELIEFS BETWEEN RHEUMATOLOGISTS, PHYSICAL THERAPISTS AND PATIENTS FOR FIBROMYALGIA- A CROSS-SECTIONAL SURVEY

S. K. Paramasivam^{1,2}, C. S. Kumar² on behalf of Chronic Pain Research Group of Academy of Orthopaedic Manual Physical Therapists (AOMPT). ¹Academy of Orthopaedic Manual Physical Therapists (AOMPT), Bethel Medical Mission, Physiotherapy, Bengaluru, India; ²Bethel Medical Mission, Physiotherapy, Bengaluru, India