Methods: cohort retrospective study, the data being gathered from RRB, which includes all the patients treated with biologics and targeted synthetic DMARDs in the country. Inclusion criteria: at least one treatment course of JAK inhibitors, until end of 2019. The collected data: demographic and disease characteristics, efficacy parameters (DAS28) and safety data (exposure to JAK inhibitors in person-years (PY) and reported adverse events (AE)).

Results: the RA cohort treated with JAK inhibitors included 416 patients (7.6% of all RA patients registered in RRB) exposed to baricitinib (306 pts) or tofacitinib (110 pts) for 66255 PY. The mean age was 57.91 years, 365 (88%) women, 41 (10%) smokers, mean RA duration 11.32 years, 316 (76%) RF positive, 254 (61%) ACPA positive, 356 (85.5%) patients associate a synthetic DMARD, 76 (18%) use steroids (66% <7.5mg prednisone/day); 268 (65%) patients have cardiovascular comorbidities, 84 pts (20%) have modified lipid profile, 89 patients (21.4%) have latent TB, 86 (20.7%) have resolved HBV infection state. JAK inhibitors were the first treatment option for 178 patients (43%), all other patients have been exposed previously to at least one TNF-inhibitor or a biologic with another mode of action, being insufficient responders, as follows: 96 patients – to 1 biologic, 80 patients – to 2 biologics, 30 patients – to 3 biologics, 23 patients – to 4 biologics, 8 patients – to 5 biologics and 1 patient - to 7 molecules before the JAK inhibitors start. Of all patients exposed to JAK inhibitors, 252 (60.5%) patients have continued the therapy since the start summarising 61481PY of exposure. The efficacy parameters reported to this category revealed a mean DAS28 score of 3.08, significantly lower compared to 6 months before (4.77). Regardless the number of biologic agents used before starting a JAK inhibitor, there is no significant difference in DAS28 score when comparing it with DAS28 efficacy of the first treatment option scenario (3.10 vs 2.94, $p > 0.05$). Steroid use, the presence of RF or disease duration have no influence on the efficacy DAS28. On contrary, the ACPA positivity subgroup has a significantly lower DAS28 score, compared to ACPA negativity (3.18 vs 3.82, $p < 0.05$); smoker status has a significantly higher DAS28 compared to non-smokers (3.58 vs 2.94, $p < 0.05$). A total number of 46 AEs were reported for a total exposure of 66255PY. Infections were the most frequently reported AE (0.04/100PY, with 27 events); 6 cases of herpes zoster infection (0.009/100PY, all mild); 3 MACE (0.004/100PY). There were 3 severe adverse events (0.004/100PY) (1 hepatitis non-viral, 2 MACE) with one fatal (acute myocardial infarction).

Conclusion: JAK inhibitors represent a safe and efficacious treatment option for patients with active RA.

Disclosure of Interests: Catalin Codreanu Consultant of: Speaker and consulting fees from AbbVie, Accord Healthcare, Alfasigma, Egis, Eli Lilly, Ewopharma, Genesis, Mylan, Novartis, Pfizer, Roche, Sandoz, UCB, Speakers bureau: Speaker and consulting fees from AbbVie, Accord Healthcare, Alfasigma, Egis, Eli Lilly, Ewopharma, Genesis, Mylan, Novartis, Pfizer, Roche, Sandoz, UCB, Corina Mogosan Consultant of: Speaker and consulting fees from AbbVie, Novartis, Pfizer, Roche, Sandoz, Speakers bureau: Speaker and consulting fees from AbbVie, Novartis, Pfizer, Roche, Sandoz, Denisa Predeteanu Consultant of: Speaker fee for: Pfizer, Abbvie, Roche, MSD, Speakers bureau: Speaker fee for: Pfizer, Abbvie, Roche, MSD, Simona Rednic: None declared, Magda Parvu Consultant of: Speaker fee and consultant: Pfizer, Novartis, Roche, Abbvie, UCB, Eli-Lilly, Speakers bureau: Speaker fee and consultant: Pfizer, Novartis, Roche, Abbvie, UCB, Eli-Lilly, Elena Rezus: None declared, Ruxandra Ionescu Consultant of: Consulting fees from Abbvie, Eli-Lilly, Novartis, Pfizer, Roche, Sandoz, Speakers bureau: Consulting and speaker fees from Abbvie, Eli-Lilly, Novartis, Pfizer, Roche, Sandoz

DOI: 10.1136/annrheumdis-2020-eular.4563

FRI0120 ORAL GLUCOCORTICOID USE IS ASSOCIATED WITH HYPERTENSION IN PATIENTS WITH RHEUMATOID ARTHRITIS

R. E. Costello¹, B. Birlie Yimer¹, M. Jani¹, W. Dixon¹. ¹The University of Manchester, Centre for Epidemiology Versus Arthritis, Manchester, United Kingdom

Background: Oral glucocorticoids (GC) are frequently prescribed to patients with rheumatoid arthritis (RA), however GC use is associated with a number of potential side effects. Hypertension is cited as a possible side effect, but few studies have specifically investigated GC-associated hypertension in patients with RA with conflicting results.

Objectives: The aim of this study was to determine whether GCs were associated with an increased risk of incident hypertension in a cohort of patients with RA.

Methods: A retrospective cohort of patients with incident RA and no hypertension at RA diagnosis were identified from UK primary care electronic health records (Clinical Practice Research Datalink). GC prescriptions were used to determine time-varying GC use and dose, categorised as: no use, >0–4.9mg/day, 5–7.4mg/day, 7.5–14.9mg/day, ≥15mg/day. A 3-month risk attribution model was used where patients continued to remain at risk for 3 months after the end of prescriptions. Hypertension was identified if a patient had either: 1) 2 consecutive systolic blood pressure (BP) measurements >140mmHg within a year, 2) 2 consecutive diastolic BP measurements

>90mmHg within a year or 3) antihypertensive prescriptions on at least two occasions and a Read code for hypertension. Unadjusted and adjusted Cox proportional hazards (PH) regression models were fitted to determine if there was an association between GC use and hypertension. Models were adjusted for baseline age, gender, baseline body mass index, baseline ever smoking, time-varying synthetic disease-modifying anti-rheumatic drug use, time-varying non-steroidal anti-inflammatory drug use and baseline Charlson comorbidity index.

Results: There were 17,760 patients with incident RA and no hypertension. The cohort had a mean age of 56.3 ± 12.7 years and were predominantly female (68%). 7,421 (42%) were prescribed GCs during follow-up. There were 6,243 cases of incident hypertension over 97547 person years (pyrs) of follow-up, giving an incident rate of 64.1 per 1000 pyrs. Of those 1321 cases were in those exposed to GCs and 4922 were in those unexposed, giving incident rates of 87.6 per 1000 pyrs and 59.7 per 1000 pyrs, respectively. The adjusted Cox PH model indicated that recent GC use was associated with a 17% increased hazard of hypertension (hazard ratio: 1.17 (95% CI 1.10 to 1.24)). When categorised by dose, the adjusted model indicated only doses above 7.5mg were significantly associated with hypertension (Table 1).

Table 1. Unadjusted and adjusted Cox proportional hazards regression model results

	Unadjusted HR (95% CI)	Age and gender adjusted HR (95% CI)	Fully adjusted* HR (95% CI)
Recent GC use	1.44 (1.35 to 1.53)	1.23 (1.16 to 1.31)	1.17 (1.10 to 1.24)
Recent GC dose	Reference	Reference	Reference
No GC use	1.35	1.13	1.10
>0 – 4.9mg	(1.21 to 1.53)	(1.01 to 1.28)	(0.98 to 1.24)
5mg – 7.4mg	1.40 (1.22 to 1.60)	1.11 (0.97 to 1.27)	1.07 (0.93 to 1.23)
7.5mg – 14.9mg	1.44 (1.33 to 1.57)	1.26 (1.16 to 1.38)	1.18 (1.08 to 1.29)
15mg and over	1.60 (1.40 to 1.84)	1.45 (1.27 to 1.66)	1.36 (1.18 to 1.56)

* Adjusted for: Baseline age, gender, baseline body mass index, baseline ever smoking, synthetic disease-modifying anti-rheumatic drug use (time-varying), non-steroidal anti-inflammatory drug use (time-varying) and baseline Charlson comorbidity index.

Conclusion: In this large cohort of patients with RA and without hypertension, recent GC use was associated with incident hypertension. In particular doses ≥7.5mg were associated with hypertension while the association with lower doses was inconclusive. Clinicians need to consider cardiovascular risk when prescribing GCs and ensure BP is regularly monitored.

Disclosure of Interests: Ruth E Costello: None declared, Belay Birlie Yimer: None declared, Meghna Jani Speakers bureau: Grifols, William Dixon Consultant of: Bayer and Google

DOI: 10.1136/annrheumdis-2020-eular.1000

FRI0121 STEROID-SPARING EFFECT OF JAK INHIBITORS IN RHEUMATOID ARTHRITIS PATIENTS FOLLOWED UP IN A REAL LIFE SETTING

I. Duca¹, F. R. Spinelli¹, F. Ceccarelli¹, C. Garufi¹, S. Mancuso¹, C. Alessandri¹, R. Scivo¹, R. Priori¹, V. Ricciari¹, M. Di Franco¹, F. Conti¹. ¹Sapienza Università di Roma, Rheumatology Unit, Rome, Italy

Background: Glucocorticoids (GCs) are a milestone of Rheumatoid Arthritis (RA) treatment; EULAR recommendations on the management of medium to high dose glucocorticoids remind to evaluate comorbidities and risk factors for adverse events when planning GCs treatment. Tofacitinib and Baricitinib are Janus kinases inhibitors (JAKi) registered for RA treatment. About 60% of patients enrolled in randomized clinical trials with JAKi were co-treated with GCs; however, little is known about tapering and percentage of withdrawal both in clinical trials and real life.

Objectives: To evaluate the steroid-sparing effect of JAKi in patients with RA.

Methods: We prospectively enrolled consecutive adult patients with RA starting JAKi. At baseline and after 4, 12 and 24 weeks we calculated C-Reactive Protein based Disease Activity score 28 (DAS28_{CRP}). Daily dose of GCs was recorded at each visit as prednisone (PDN)-equivalent dose. Data are expressed as median (IQR). Continuous variables were compared by Mann Whitney test while dichotomous ones by Chi-square test. P values < 0.05 were considered statistically significant.

Results: Between January 2018 and January 2020, 108 patients started JAKi: 67 patients Baricitinib, 41 patients Tofacitinib. The analysis was restricted to 64 RA patients (50 female, 14 male) who had at least 6 months of follow-up. Table 1 shows the demographic, clinical and clinimetric characteristics of the cohort.