Objectives: To compare the epidemiological, clinical, biological, radiological, bacteriological profile as well as the management and prognosis of spondylodiscitis over the last 20 years.

Methods: Retrospective study including 113 patients hospitalized in our department between 1999 and 2018. The diagnosis was based on clinical, biological, radiological and bacteriological data. Spondylodiscitis were divided into two groups: spondylodiscitis occurring between 1998 and 2008 (1st decade) and those between 2009 and 2018 (2nd decade).

Results: Of the 113 identified patients, 54 (47.8%) occurred between 1998 and 2008, with male predominance in both groups (50.2 and 59.3% respectively; $p=0.3$). The average age was 55 years ($p=0.7$). Diabetes was more frequent in the 1st decade but without a statistically significant difference ($p=0.1$). On the other hand, consumption of unpasteurized dairy products was more frequent during the 2nd decade ($p<0.001$) as well as the presence of intercurrent infection ($p=0.01$).

The estimated delay between symptom onset and diagnosis was longer between 1998 and 2008 (5.6 months and 3.2 months respectively, $p=0.005$). The presence of neurological signs was more frequent between 2009 and 2018 ($p=0.001$), especially radiculalgia ($p=0.02$). The sedimentation rate was higher in the 1st decade (93mm and 72mm respectively, $p=0.002$).

We found no statistically significant difference in the location of spondylodiscitis ($p=0.4$) and the multifocal involvement ($p=0.5$).

Radiographic signs were significantly more prevalent in the 1st decade ($p=0.002$), in particular disc space narrowing ($p=0.03$) and irregularity of the intervertebral plates ($p=0.01$). Computed tomography was more often performed during the 1st decade ($p=0.008$), unlike magnetic resonance imaging, performed in 88.1% of cases during the 2nd decade ($p<0.001$). In contrast, the frequency of para-vertebral abscesses, epiduritis and spinal cord compressions were similar in the two groups ($p=0.6$; $p=0.9$ and $p=0.3$, respectively).

Tuberculosis was more frequent in the 1st decade ($p=0.009$). Disco-vertebral biopsy was more often performed in the 1st decade ($p=0.04$), but its efficiency was similar between the two groups ($p=0.1$). For pyogenic spondylodiscitis, blood cultures were more often positive in the 2nd decade but without a statistically significant difference ($p=0.6$). On the other hand, the anatomopathological aspect was more frequently suggestive of a pyogenic germ in the 2nd decade ($p=0.04$).

A surgical procedure was more frequently performed between 2009 and 2018 but with no statistically significant difference ($p=0.2$). All patients received antibiotic therapy. A change in treatment was made in 23.7% of cases between 2009 and 2018 ($p=0.01$). The prognosis was better during the 1st decade ($p=0.01$). On the other hand, the occurrence of immediate complication during tuberculous spondylodiscitis was more frequent during the 2nd decade ($p=0.03$), in particular the toxicity of anti-tuberculosis treatment ($p=0.04$).

Conclusion: In the last decade, the diagnosis of spondylodiscitis has been made earlier. However, these are more severe forms with an increased frequency of neurological complications, probably due to an increase in the proportion of virulent germs. Moreover, the effectiveness of the bacteriological investigation does not differ between the two periods studied.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.4814

FRI0434

PREDICTIVE FACTORS OF SERIOUS INFECTIONS IN PATIENTS WITH CHRONIC INFLAMMATORY ARTHRITIS ON TREATMENT WITH TNF INHIBITORS: REAL LIFE DATA

A. Ruiz Ochoa¹, J. Escudero Argaluz², B. Alvarez Rodríguez², M. Vasques Rocha², C. Stoye², O. Pompei Fernández², S. Gil², J. R. De Dios², J. Calvo-Alen². ¹Universidad del País Vasco, Vitoria, Spain; ²Hospital Universitario Araba, Rheumatology, Vitoria, Spain

Background: Serious infections are one of the most feared adverse events in patients treated with biologics. To this regard data coming from randomized and long-term extension studies may not totally applied to the usual clinical practice due to the different profile of the treated patients. To study the associated factors for serious infections in patients with inflammatory arthritis treated with TNF inhibitors

Objectives: To study the associated factors for serious infections in patients with inflammatory arthritis treated with TNF inhibitors

Methods: All the medical records of the patients with inflammatory arthritis being treated with TNF inhibitors at the beginning of 2016 were retrospectively reviewed. All serious infections suffered for these patients until the end of 2018 were recorded. Serious infections were defined as those which required to admitted at the hospital for intravenous treatment. Potential variables associated with the development of these infections including: demographic and clinical characteristics, concomitant treatments or comorbidity (by Charlson index) were studied. Standard statistical tests for descriptive and univariate analyses were used

and a multivariable logistic regression model was built to check independent associations.

Results: Overall 334 patients (50.3% women) with a mean age of 56.67 (± 12.853) were studied: 140 (41.92%) Rheumatoid arthritis (RA), 55 (16.46%) psoriatic arthritis (PsA) and 138 (41.62%) spondyloarthritis (Sp). Forty-five serious infections were observed in 30 patients, being respiratory (40%) and urinary (8.8%) the most frequent localizations. Only one patient died. By univariate analysis, disease duration, age, concomitant use of glucocorticoids (GC) (but not of synthetic DMARDs), Charlson index and specifically Diabetes Mellitus were associated with infection ($p<0.05$). The type of arthritis was not associated and the results in the subset of RA patients were overall similar. In the multivariate analysis the use of GC [OR: 5.31 (1.98,14.26)] and the Charlson index [OR:2.48 (1.70;3.60)] were found to be independently associated to infection.

Conclusion: In patients with inflammatory arthritis and treated with TNF inhibitors around a 10% developed any serious infection along three years of follow up. Use of GC and comorbidity emerged as the main risk factors for this complication.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.4347

FRI0435

THE CLINICAL SIGNIFICANCE OF SERUM AUTOANTIBODIES AND HLA-B27 MOLECULE TESTING IN UYGUR PATIENTS WITH HUMAN IMMUNODEFICIENCY VIRUS INFECTION

X. Chen¹, L. Wu², X. Wu², C. N. Luo², Y. M. Shi². ¹People's Hospital of Xinjiang Uygur Autonomous Region, Department of Rheumatology and Immunology, Urumqi, China; ²People's Hospital of Xinjiang Uygur Autonomous Region, Urumqi, China

Background: AIDS is a deadly infectious disease caused by the HIV. When HIV infects a host, it may induce production of autoantibodies due to the structural antigen similarity between viral proteins and selfantigens. The molecular mimicry between HIV protein and self-antigens could cause antibody cross-reactions and lead to development of autoimmune disease.

Objectives: To explore the clinical value of serum autoantibodies and human leukocyte antigen (HLA-B27) molecular testing in Uygur patients with human immunodeficiency virus (HIV) infection.

Methods: A total of 727 HIV-infected Uygur patients who visited Kuche Infectious Diseases Hospital during May 2016 to March 2017 were include in this study. The other 390 healthy people were enrolled as controls. Serum antinuclear antibodies (ANA) and ANA Profile, anti-cyclic citrullinated peptide (CCP) antibody, and HLA-B27 molecule were tested.

Results: Among 727 HIV-infected Uygur patients, 317 were males and 410 were females with mean age (35.52 ± 13.44) years old. The mean duration of disease was (6.34 ± 3.05)years. There were 697(95.87%) patients receiving Highly active antiretroviral therapy (HAART) with mean duration of treatment (6.34 ± 3.05)years. Rheumatic manifestations were recorded in 238 (32.74%) HIV-infected Uygur patients, mainly with dry mouth and dry eye (15.41%), alopecia (9.90%), arthralgia (8.94%), ect. Compared with the health controls, positive ANA was more common in HIV infected Uygur patients (33.42% vs. 17.43%, $P<0.001$) with low titers (ANA titer:1:100). HIV-infected Uygur patients had higher positive anti-u1-RNP antibodies positive rate (1.10%), but lower anti-SSA antibodies positive rate (0.14%) and anti-CCP antibodies positive rate (0.28%). Patients with positive ANA in HAART group were significantly less than that in non-treatment group (38.72% vs. 50.00%, $P=0.049$). Only one female patient was HLA-B27 positive (0.14%), which was significantly lower than that in healthy controls (3.08%) ($P<0.001$). Also, only one patient was diagnosed with rheumatoid arthritis (RA).

Conclusion: Rheumatic manifestations are common in HIV-infected Uygur patients. Several autoantibodies are positive, but the coincidence of rheumatic diseases is rare. It's noted that patients with Rheumatic manifestations and low titre positive ANA should be considered as a differential diagnosis of HIV infection.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.5155

FRI0436

ARTICULAR AND PERIARTICULAR INFECTIOUS DISEASES: REVIEW AND ANALYSIS OF A COHORT IN A 3RD LEVEL HOSPITAL

R. Dos Santos Sobrin¹, N. Pérez Gómez¹, E. Perez-Pampín¹, A. Souto Vilas¹, A. Mera Varela¹. ¹Clinical University Hospital in Santiago de Compostela, Rheumatology Department, Santiago de Compostela, Spain

Background: Infectious diseases in Rheumatology are still a prevalent health-care issue nowadays. Microbiological tests and early beginning of antibiotic