³Radoud University Medical Centre, Nijmegen, Netherlands; ⁴Bicocca University, Milan, Italy; ⁵IRCCS San Raffaele Scientific Institute, Milan, Italy

Background: Histiocytoses are disorders characterized by tissue infiltration by macrophages, dendritic cells, or monocyte-derived cells. These diseases are classified in five groups based on histologic, clinical, and molecular features: Langerhans-related, cutaneous/mucocutaneous and malignant histiocytoses, Rosai-Dorfman disease, and hemophagocytic lymphohistiocytosis (1). Langerhans-related histiocytoses comprise Langerhans cell histiocytosis and Erdheim-Chester disease, both inflammatory myeloid-driven diseases characterized by clonal activating mutations along the MAPK or related pathways, most commonly *BRAFV600E*, and by severe tissue and systemic inflammation (2). Here, we describe and characterize a novel, related histiocytosis chiefly manifested with severe synovial involvement.

Objectives: Here, we describe a novel histiocytosis whose histologic and clinical picture (severe synovial involvement, systemic inflammation, skin lesions, and diabetes insipidus) differed from known histiocytic disorder. In addition, we performed molecular studies aimed at identifying causative activating mutations. Finally, by means of a dynamic 3D tissue culture system we characterized immune-metabolic mechanisms underlying disease pathogenesis and clinical response to treatment.

Methods: The mutational status of oncogenes was determined with a mass spectrometry multiplexed genotyping approach (PentaPanel). Biospy samples were cultured in RCCSTM bioreactor (Synthecon) in the presence/absence of a MEK inhibitor (GSK1120212, 1nM), and then either processed for immunohistochemical analyses, or lysed for western blot analysis. Culture supernatants were collected for cytokine, chemokine and metabolome determination. The Bio-Plex Multiple-Cytokine Assay (Bio-Rad) and the Ella assay (ProteinSimple) were used to determine cytokine concentrations in supernatants and serum, respectively. Metabolomic studies were performed as described (3).

Results: We identified a causative mutation in the proto-oncogene *KRAS* (*KRASG12D*, not previously reported in related histiocytoses). In addition, 3D culture studies of patient's biopsies revealed *KRAS*-driven signaling, phenotypic, and immunometabolic features. These included constitutive ERK and AKT phosphorylation, up-regulated glucose metabolism with glycolysis and TCA cycle activation, and deregulated release of pro-inflammatory cytokines IL-1 β , IL-6 and TNF α . All these features reverted upon pharmacologic inhibition of the MAPK pathway. Characterization of this novel condition instructed effective treatment of the patient with the MEK inhibitor cobimetinib.

Conclusion: Genetic, clinical, and histopathology features differentiate this condition from known histiocytic disorders. Mechanistically, *KRASG12D* causes constitutive activation of the MAPK pathway in macrophages, which results in maladaptive changes in cell energy metabolism sustaining rampant production of pro-inflammatory cytokines. Besides instructing effective treatment of this patient, these studies revealed metabolic rewiring as key to pathologic inflammatory activation of macrophages in human disease.

References:

- [1] Emile JF, et al. Revised classification of histiocytoses. *Blood*. 2016
- [2] Cavalli G, et al. The multifaceted clinical presentations and manifestations of Erdheim-Chester disease. *Ann Rheum Dis*. 2013
- [3] Cavalli G, et al. Interleukin 37 reverses the metabolic cost of inflammation, increases oxidative respiration, and improves exercise tolerance. *Proc Natl Acad Sci U S A*. 2017

Disclosure of Interests: Giulio Cavalli Consultant of: SOBI, Pfizer, Sanofi, Novartis, Paid instructor for: SOBI, Novartis, Speakers bureau: SOBI, Novartis, Giacomo De Luca Speakers bureau: SOBI, Novartis, Celgene, Pfizer, MSD, Riccardo Biavasco Employee of: Bluebird, Travis Nemkov: None declared, Angelo D'Alessandro: None declared, Rob Arts: None declared, Antonello Villa: None declared, Daniela Belloni: None declared, Greta Grassini: None declared, Giulia Cangi: None declared, Claudio Doglioni: None declared, Elisabetta Ferrero: None declared, Marina Ferrarini: None declared, Lorenzo Dagna: None declared

DOI: 10.1136/annrheumdis-2020-eular.3170

FRI0480 SCHNITZLER'S SYNDROME: DESCRIPTION OF AN ITALIAN MULTICENTER COHORT

F. Crisafulli¹, P. Airò², F. Franceschini¹, A. Tincani¹, M. Frassi² on behalf of Working Group of Autoinflammatory Diseases of SIR (Italian Society of Rheumatology). ¹ASST Spedali Civili and University of Brescia, Rheumatology and Clinical Immunology Unit, Brescia, Italy; ²ASST Spedali Civili of Brescia, Rheumatology and Clinical Immunology Unit, Brescia, Italy

Background: Schnitzler's syndrome is an autoinflammatory disease characterized by monoclonal gammopathy and recurrent episodes of urticaria accompanied by clinical and laboratory signs of acute inflammation. Although the exact

pathogenic mechanisms have not been fully clarified, the role of Interleukin-1 seems to be central.

Objectives: To describe clinical features and therapeutic approach in patients with Schnitzler's Syndrome.

Methods: Retrospective analysis of an Italian multicenter cohort. Data are expressed as the median (IQR).

Results: The clinical data of 24 patients from 9 centers (median follow-up 6 years [2-10]; median age at diagnosis 56.5 years [51.25-64.25]) were collected. The median diagnostic delay was 2 years (0-10); the diagnosis was made consensually at the onset of symptoms in 4 cases. The main clinical and laboratory features are shown in Table 1. Therapeutic response was evaluable in 20 patients: all received corticosteroids (CS; 25mg/day [25-50]); in one case, a good clinical response was observed. Eight patients were initially treated with colchicine: in 3 cases it was effective in controlling symptoms and reducing the dose of CS; other 8 patients were treated with csDMARDs (n:1 [1-2]): only 1 patient had a good response to cyclosporin.

Table 1. Clinical and laboratory features

Chronic Urticarial Rash, n (%)	24/24 (100)
Pruritus, n (%)	17/24 (71)
Intermittent fever, n (%)	23/24 (96)
Arthralgia/Arthritis, n (%)	20/24 (83)
Bone pain, n (%)	8/24 (33)
Weight loss, n (%)	9/24 (38)
Angioedema, n (%)	4/24 (17)
Lymphadenopathy, n (%)	7/24 (29)
Hepatomegaly, n (%)	3/24 (12)
Splenomegaly, n (%)	3/24 (12)
Neuropathy, n (%)	4/24 (17)
Raised ESR or CRP, n (%)	24/24 (100)
Leukocytosis, n (%)	17/24 (71)
Anemia, n (%)	9/24 (38)
Monoclonal Gammopathy	
IgG λ , n (%)	5/22 (23)
IgG κ , n (%)	6/22 (27)
IgM λ , n (%)	1/22 (5)
IgM κ , n (%)	12/22 (55)
Bence Jones Protein, n (%)	6/23 (26)

A bDMARD was initiated in 15 patients. In 7 of the 14 patients initially treated with anakinra this therapy was continued with benefit whereas in the other 7 patients the treatment was discontinued for primary inefficacy (1 patient), secondary inefficacy (3 patients) and adverse events (3 patients; 2 injection site reaction, 1 severe allergic reaction). After anakinra discontinuation, 5 patients were treated with canakinumab with a good response in 3 cases and a partial response in 1 case (persistent arthritis); 1 patient died during the treatment. No response was observed in 3 patients treated with TNF inhibitors as a 2nd or 3rd line bDMARDs, as well as in 1 case initially treated with tocilizumab (in which a good response was afterwards obtained with canakinumab). bDMARDs were associated with a csDMARD in 2 patients (methotrexate and methotrexate + cyclosporine).

In one case monoclonal gammopathy evolved into Multiple Myeloma and the patient died 15 years after the onset of symptoms. Idiopathic myelofibrosis and myelodysplasia were found in one and in two patients, respectively.

Conclusion: In most cases csDMARDs and bDMARDs like anti-IL6 and anti-TNF α were not able to control the disease. In contrast, in some cases, a good response to colchicine was observed; refractory patients may be successfully treated with anti-IL1 agents. Patients should be supervised for possible evolution towards lymphoproliferative disease.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.3770

FRI0481 HEMOPHAGOCYTIC SECONDARY SYNDROME. DIFFERENCES BETWEEN AUTOIMMUNE AND HEMATO-ONCOLOGICAL ETIOLOGIES

C. A. Egües Dubuc¹, A. De Diego¹, P. Cabrera Miranda², N. Alcorta Lorenzo¹, J. A. Valero Jaimes¹, J. R. Furundarena Salsamendi³, L. M. Lopez Dominguez¹, J. J. Cancio Fanlo¹, O. Maiz-Alonso¹, E. Uriarte Isacelaya¹, J. Calvo⁴, J. M. Belzunegui Otano¹. ¹Donostia University Hospital, Rheumatology, Donostia, Spain; ²University Hospital October 12, Preventive Medicine and Public Health, Madrid, Spain; ³Donostia University Hospital, Haematology, San Sebastian, Spain; ⁴Araba University Hospital, Rheumatology, Vitoria, Spain

Background: The Hemophagocytic Syndrome (HPS) is has been classified into 2 groups: primary and secondary. The secondary form is mainly associated with hemato-oncological diseases (HOD) and autoimmune diseases (AID). The