including angiogenesis. Less is known about the role of YKL-40 in inflammatory diseases such as GCA.

Objectives: Our objective was to investigate the cellular source and the pro-angiogenic function of YKL-40 in GCA patients.

Methods: For this study we performed immunohistochemistry (IHC) and cell culture experiments. IHC for YKL-40 and CD206 was performed on GCA positive temporal artery biopsies (TABs; n=12) and GCA positive aortas (n=10) of treatment-naïve patients. Expression of YKL-40 by macrophages was confirmed by double staining with macrophage transcription factor PU.1. Additionally, the TABs were stained for IL-13R α 2, recently described as the receptor for YKL-40. The effect of skewing signals on YKL-40 production was assessed by cell culture of monocyte-derived macrophages of GCA patients with either M-CSF or GM-CSF (n=8). Subsequently, the supernatant was assayed by ELISA. Finally, the angiogenic potential of YKL-40 was investigated by tube formation experiments using human microvascular endothelial cells (HMVECs).

Results: YKL-40 is produced by a distinct subset of macrophages in GCA TABs and aortas, usually located in or near the media (Figure 1 shows representative stainings in consecutive slides of a GCA TAB). We here show YKL-40 to be expressed by CD206+/MMP-9+ macrophages in all GCA TABs and aortas. In vitro, macrophages were found to produce YKL-40 (Figure 2 shows an increasing YKL-40 production during the maturation of monocytes towards macrophages over 8 days of culture). GM-CSF stimulation, which is known to upregulate CD206 expression in macrophages, gave rise to higher YKL-40 production by GCA macrophages, when compared to M-CSF stimulated macrophages from GCA patients (p=0.038). In addition, YKL-40 stimulation of HMVECs induced more tube formation compared to unstimulated HMVECs. Finally, we showed, by IHC, abundant expression of the YKL-40 receptor IL-13R α 2 in TABs of GCA patients.

Figure 1

Conclusion: Taken together, we show here that a distinct subset of macrophages, skewed by GM-CSF and highly positive for CD206, is responsible for the production of YKL-40 in GCA. The results are in line with previous reports demonstrating that CD206 expression distinguishes YKL-40 positive macrophages from YKL-40 negative macrophages (1). Thus, YKL-40 production by CD206+ macrophages may be involved in angiogenesis in GCA tissues, a process important for the continuation of the inflammatory process.

Figure 2

References:

- [1] Bonneh-Barkay, D. et al., 2012. Astrocyte and macrophage regulation of YKL-40 expression and cellular response in neuroinflammation. *Brain Path.* 22: 530-546

Disclosure of Interests: Yannick van Sleen: None declared, William Febry Jiemy: None declared, Sarah A. Pringle: None declared, Wayel Abdulahad: None declared, Kornelis van der Geest Speakers bureau: Roche (2019), Maria Sandovici: None declared, Peter Heeringa: None declared, Elisabeth Brouwer Consultant of: Roche (consultancy fee 2017 and 2018 paid to the UMCG), Speakers bureau: Roche (2017 and 2018 paid to the UMCG), Annemieke Boots Consultant of: Grünenthal GmbH until 2017

DOI: 10.1136/annrheumdis-2020-eular.4455

Connective tissue diseases - genomics, proteomics and pathogenesis

OP0135

INHIBITION OF HSP90 REDUCES PROGRESSION OF DERMAL FIBROSIS AND INDUCES REGRESSION OF ESTABLISHED EXPERIMENTAL DERMAL FIBROSIS INDUCED BY BLEOMYCIN

H. Štorkánová^{1,2}, L. Štorkánová¹, S. Oreska^{1,2}, M. Špiritović³, B. Heřmánek³, R. Bečvář^{1,2}, K. Pavelka^{1,2}, J. Vencovský^{1,2}, J. Distler⁴, L. Šenolt^{1,2}, M. Tomčík^{1,2}. ¹Institute of Rheumatology, Prague, Czech Republic; ²Department of Rheumatology, 1st Faculty of Medicine, Charles University, Prague, Czech Republic, Prague, Czech Republic; ³Faculty of Physical Education and Sport, Charles University, Prague, Czech Republic, Prague, Czech Republic; ⁴Department of Internal Medicine III and Institute for Clinical Immunology, University of Erlangen-Nuremberg, Erlangen, Germany, Erlangen, Germany

Background: Our previous study demonstrated that Heat shock protein 90 (Hsp90) is overexpressed in the skin of patients with systemic sclerosis (SSc), in cultured SSc fibroblasts and preclinical models of SSc. HSP90 is a new regulator of canonical TGF- β signalling and its inhibition prevents the stimulatory effects of TGF- β on collagen synthesis and dermal fibrosis in three preclinical models of SSc.

Objectives: Herein, we aimed to evaluate the efficacy of Hsp90 inhibitor (17-DMAG) in the treatment of established experimental dermal fibrosis induced by bleomycin.

Methods: Design consisted of three control groups, I (NaCl-s.c./6 weeks), II (bleomycin-s.c./3w and NaCl-s.c./3w), III (bleomycin-s.c./6w), and 2 treatment groups (bleomycin-s.c./6w). During the last 3 weeks, one group was treated with 17-DMAG 0.5mg/kg-i.p. every third day, whereas one group (with nintedanib 50mg/kg-p.o. twice daily) served as a comparator with already published efficacy in this setting. Total of 40 BL6 mice were examined weekly for weight, activity and fur texture. The effects of 17-DMAG were determined by assessment of dermal thickness (HE-staining), collagen content (hydroxyproline assay), myofibroblast

counts (α -SMA staining) and of 23 serum inflammatory cytokines/chemokines (Mouse-Cytokine-23-plex, Bio-Rad-Laboratories).

Results: 17-DMAG decreased dermal thickening by $53\pm 3\%$ ($p<0.001$) (nintedanib by $46\pm 2\%$, $p<0.001$), collagen content by $48\pm 5\%$ ($p=0.004$) (nintedanib by $50\pm 4\%$, $p=0.003$), myofibroblast counts by $42\pm 9\%$ ($p<0.001$) (nintedanib by $44\pm 7\%$, $p<0.001$), and levels of IL-1 α , IL-6, IL-12(p40), CXCL1, MCP-1, MIP-1 α/β , RANTES (in all: $p<0.05$) compared to vehicle-treated mice injected with bleomycin for 6w. Moreover, 17-DMAG also induced regression of pre-established fibrosis to below the levels of vehicle-treated mice injected with bleomycin for 3w and NaCl for 3w (dermal thickness by $14\pm 3\%$, collagen content by $20\pm 5\%$, myofibroblast counts by $13\pm 9\%$; whereas in nintedanib by $10\pm 3\%$, $21\pm 4\%$, $17\pm 7\%$, respectively; in all: $p<0.05$), and levels of IL-12(p40), CXCL1, MCP-1, MIP-1 β , RANTES (in all: $p<0.05$). No significant weight loss, decrease in activity or changes in fur texture were observed upon 17-DMAG treatment.

Conclusion: This is the first study on effects of Hsp90 inhibitor 17-DMAG in the treatment of established dermal fibrosis. We demonstrate that 17-DMAG effectively prevents the progression and induces regression of established bleomycin-induced dermal fibrosis, in an extent that was comparable to nintedanib in this study (which was recently FDA approved for slowing the rate of decline in lung function in adults with SSc-ILD). 17-DMAG was well tolerated without obvious clinical signs of toxicity. These data suggest that Hsp90 could be a novel potential target in the treatment of SSc dermal fibrosis.

Acknowledgments: Supported by AZV-16-33542A, MHCR023728, SVV260373, Boehringer Ingelheim.

Disclosure of Interests: Hana Štorkánová: None declared, Lenka Štorkánová: None declared, Sabina Oreska: None declared, Maja Špiritović: None declared, Barbora Heřmánková: None declared, Radim Bečvář Consultant of: Actelion, Roche, Karel Pavelka Consultant of: Abbvie, MSD, BMS, Eisai, Roche, UCB, Medac, Pfizer, Biogen, Speakers bureau: Abbvie, MSD, BMS, Eisai, Roche, UCB, Medac, Pfizer, Biogen, Jiří Vencovský: None declared, Jörg Distler Grant/research support from: Boehringer Ingelheim, Consultant of: Boehringer Ingelheim, Paid instructor for: Boehringer Ingelheim, Speakers bureau: Boehringer Ingelheim, Ladislav Šenolt: None declared, Michal Tomcik: None declared

DOI: 10.1136/annrheumdis-2020-eular.3597

OP0136

THE INFLUENCE OF LONG-TERM EXERCISE AND IN VITRO EXERCISE-MIMICKING STIMULATION ON THE PRODUCTION OF MYOKINES AND CYTOKINES IN MYOTUBES OF PATIENTS WITH CHRONIC IDIOPATHIC INFLAMMATORY MYOPATHIES

M. Vokurková¹, L. Vernerová¹, M. Špiritović², H. Štorkánová³, S. Oreska³, M. Klein³, J. Ukropec⁴, B. Ukropcová⁵, M. Tomčík³, J. Vencovský³. ¹Institute of Rheumatology, Prague, Czech Republic; ²Institute of Rheumatology and Department of Physiotherapy, Faculty of Physical Education and Sport, Charles University, Prague, Czech Republic; ³Institute of Rheumatology and Department of Rheumatology, First Faculty of Medicine, Charles University in Prague, Prague, Czech Republic; ⁴Institute of Experimental Endocrinology, Biomedical Research Center, Slovak Academy of Sciences, Bratislava, Slovakia (Slovak Republic); ⁵Institute of Experimental Endocrinology, Biomedical Research Center, Slovak Academy of Sciences and Institute of Pathophysiology, Faculty of Medicine, Comenius University, Bratislava, Slovakia (Slovak Republic)

Background: It has been demonstrated several times that endurance exercise has beneficial effects on the condition of patients with idiopathic inflammatory myopathies (IIM). Muscle contraction during exercise is a major stimulus for the release of myokines that are supposed to take part in the beneficial adaption to exercise.

Objectives: The aim of this study was to find out how a six-month physiotherapy and *in vitro* exercise-mimicking treatment affect myokine and cytokine production in myotubes of IIM patients.

Methods: Seven patients with chronic IIM took part in a six-month physiotherapy (stretching and strengthening), which significantly improved their muscle strength and endurance. IIM patients ($n=7$) before and after the six months exercise and their respective healthy counterparts (HC, $n=9$) underwent a *musculus vastus lateralis* biopsy. Isolated skeletal muscle cells were grown, differentiated into myotubes, which were treated with a pharmacological cocktail: palmitate, forskolin and ionomycin (PFI) to mimic exercise-stimulated contractions *in vitro*. Myokine and cytokine concentrations produced by myotubes to the culture medium were analyzed with ELISA and the multiplex immunoassay, respectively. RT-PCR was used for the evaluation of myokine gene expression in the cultured myotubes.

Results: Compared to myotubes of healthy controls, myotubes of IIM patient released more myostatin and activin A into the medium. The myostatin gene was expressed significantly more in muscle cells of patients than in healthy controls' cells ($p<0.05$). After a six-month rehabilitation program, activin A secretion was four-fold reduced in myotubes of patients with IIM, while myostatin release

and gene expression remained unchanged. In myotubes of IIM patients, less follistatin and more follistatin like 3 were detected in the culture medium compared to HC myotubes. Myotubes derived from IIM patients after six months of rehabilitation secreted twice as much follistatin and half the amount of follistatin like 3 into the medium than myotubes derived from IIM patients prior to rehabilitation ($p<0.05$). There was no difference in secretion of interleukin (IL) 6, IL-17, tumor necrosis factor (TNF) and vascular endothelial growth factor (VEGF) between myotubes of IIM patients and myotubes of HC. However, six-month exercise significantly ($p<0.05$) reduced release of IL-6, TNF and VEGF in myotubes of IIM patients. Contrary to our expectation, stimulation of PFI had no effect on the release of myostatin, activin A, follistatin and follistatin like 3, or the expression of their genes. PFI treatment significantly ($p<0.05$) increased IL-6 secretion in myotubes from HC and IIM patients prior to six months of rehabilitation. On the other hand, it was observed that myotubes of HC and IIM patients exposed to the PFI cocktail secreted significantly less inflammatory cytokines IL-17, TNF and VEGF into the medium compared to unstimulated myotubes ($p<0.05$).

Conclusion: In conclusion, long-term exercise influenced the production of myokines and decreased release of inflammatory cytokines in myotubes of IIM patients. *In vitro* exercise-mimicking treatment increased the secretion of IL-6 and decreased the release of inflammatory cytokines as IL-17, TNF- α and VEGF in myotubes of patients with IIM and healthy individuals.

Acknowledgments: This work was supported by the Ministry of Health of the Czech Republic grants nr. 16-33746A and donation 140.0000008.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.5543

OP0137

GENOME-WIDE WHOLE-BLOOD TRANSCRIPTOME PROFILING IN A LARGE EUROPEAN COHORT OF SYSTEMIC SCLEROSIS PATIENTS

L. Beretta¹, G. Barturen², B. Vigone¹, C. Bellocchi^{1,3}, N. Hunzelmann⁴, E. Delanghe⁵, L. Kovács⁶, R. Cervera⁷, M. Gerosa³, R. Ortega Castro⁸, I. Almeida⁹, D. Cornec¹⁰, C. Chizzolini¹¹, J. O. Pers¹⁰, Z. Makowska¹², A. Buttgerit¹², R. Lesche¹², M. Kerick¹³, M. Alarcon-Riquelme², J. Martin Ibáñez¹³. ¹Scleroderma Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; ²GENYO, Centre for Genomics and Oncological Research Pfizer, University of Granada, Andalusian Regional Government, PTS GRANADA, Granada, Spain; ³Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy; ⁴Klinik und Poliklinik für Dermatologie und Venerologie, Köln, Germany; ⁵Division of Rheumatology, University Hospitals Leuven and Skeletal Biology and Engineering Research Center, Leuven, Belgium; ⁶Department of Rheumatology and Immunology, University of Szeged, Szeged, Hungary; ⁷Department of Autoimmune Diseases, Hospital Clinic, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain; ⁸IMIBIC, Reina Sofia Hospital, University of Cordoba, Cordoba, Spain; ⁹Serviço de Imunologia EX-CICAP, Centro Hospitalar e Universitário do Porto, Porto, Portugal; ¹⁰U1227, Université de Brest, Inserm, Labex IGO, CHU de Brest, Brest, France; ¹¹Immunology & Allergy, University Hospital and School of Medicine, Geneva, Geneva, Switzerland; ¹²Bayer AG, Berlin, Germany; ¹³Institute of Parasitology and Biomedicine López-Neyra, IPBLN-CSIC, Granada, Spain

Background: The analysis of annotated transcripts from genome-wide expression studies data is of paramount importance to understand the molecular phenomena underlying the occurrence of complex diseases, such as systemic sclerosis (SSc).

Objectives: To perform whole-blood transcriptome and pathway analysis on whole-blood (WB) RNA collected in two cohorts of European SSc patients. Via a discovery and validation strategy we aimed at characterizing the molecular pathways that differentiate SSc from controls and that are reproducible in geographically diverse populations.

Methods: WB samples from 252 controls and 162 SSc patients were collected in RNA stabilizers. Patients were divided into a discovery ($n=79$; Southern Europe) and validation cohort ($n=83$; Central-Western Europe). RNA sequencing was performed by an Illumina assay. Functional annotations of Reactome pathways were performed with the FAIME algorithm. In parallel, a immunophenotyping analysis on 28 circulating cell populations was assessed. We then tested: the presence of differentially expressed genes or pathways and the correlation between absolute cell counts and RNA transcripts/FAIME scores in regression models. Results significant in both populations were considered as replicated.

Results: A total of 15224 genes and 1277 related functional pathways were available for analysis. Among these, 99 genes and 225 pathways were significant in both sets. The heatmap in figure shows the relative expression of replicated pathways and the distribution of cases and controls (red and green bars). Among the significant pathways we found a deregulation in: type-I IFN, TLR-cascade and signalling, function of the tumor suppressor p53 protein, platelet degranulation and activation. Correlation analysis showed that the count of several cell