$p < 0.001$), moderate correlations were found with HADS-A and HADS-D (HADS-A ($r = 0.618$, $p < 0.001$), HADS-D ($r = 0.507$, $p < 0.001$)). Low correlation was found between BETY-BQ and HAQ ($r = 0.286$, $p = 0.003$) (Table 2).

Table 1. Descriptive Statistics

	Minimum	Maximum	Mean	Standard Deviation
Age (years)	20	60	44.98	11.04
Height (cm)	145.00	190.00	165.61	8.88
Weight (kg)	44	138	75.54	17.36
VKI (kg/m^2)	17.85	49.00	27.54	6.02

Table 2.

	PsAQoL	HADS-A	HADS-D	HAQ
BETY-BQ	$r = 0.826^{**}$ $p < 0.001$	$r = 0.618^{**}$ $p < 0.001$	$r = 0.507^{**}$ $p < 0.001$	$r = 0.286^{**}$ $p = 0.003$

** Correlation is significant at the 0.01 level (2-tailed)

Conclusion: In this study, it was found that BETY-BQ is a scale that can be used in total to evaluate the quality of life, emotion-status level and functionality of individuals diagnosed with PsA. It was thought that BETY-BQ could be considered and preferred by clinicians when individuals with PsA should be evaluated from a biopsychosocial point of view.

References:

- Unal E, Arin G, Karaca Nb, Kiraz S, Akdogan A, Kalyoncu U, et al. Development of a quality of life measurement for rheumatic patients: item pool construction Journal of Exercise Therapy and Rehabilitation. 2017; 4 (2): 67-75.

Disclosure of Interests: Senem Bulut: None declared, Edibe Ünal: None declared, Aykut Özçadırcı: None declared, Umut Kalyoncu Consultant of: Abbvie, Amgen, Janssen, Lilly, Novartis, UCB

DOI: 10.1136/annrheumdis-2020-eular.5609

FRI0620-HPR THE EFFECT OF THERMOFORMABLE FOOT ORTHOSES ON WALKING IN RHEUMATOID ARTHRITIS PATIENTS: PRELIMINARY RESULTS FROM AN OPEN CLINICAL TRIAL

S. Fellous¹, H. Rkain¹, A. Ilham¹, L. Benzakour², T. Latifa¹, N. Alami¹, R. Guieu³, Y. Jammes³, N. Hajjaj-Hassouni⁴, F. Allali¹. ¹Faculty of Medicine and Pharmacy of Rabat, Mohammed V University, Rabat, Department of Rheumatology B, El Ayachi Hospital, Rabat, Morocco; ²PROTEOR Morocco, Casablanca, Morocco; ³University Aix-Marseille, Marseille, UMR MD2 Faculty of Medicine, Marseille, France; ⁴International university of Rabat, Rabat, Morocco

Background: Foot pain is common in rheumatoid arthritis and appears to persist despite modern day medical management.

Objectives: To evaluate the impact of thermoformable foot orthoses on walking in rheumatoid arthritis (RA) patients.

Methods: This is an open clinical trial, that included 14 consecutive patients (85.7% female, mean age 54.8 ± 10 years) with RA (median duration of progression of 9 [5-12] years), the average DAS28 was 2.7 ± 1.2 and the functional impact objectified by the Health Assessment Questionnaire (HAQ) was on average 0.9 ± 0.7 . The foot problem was bilateral in 100% and inaugural in 85.7% of the cases.

The 14 rheumatoid subjects were examined and appropriate foot orthoses were prescribed according to each patient's needs. All the patients were evaluated at baseline and 8 weeks after use of orthoses. Gait pain, difficulty walking and the 10 Meter Walk test were noted at each appointment.

We used dynamic baropodometric analysis to assess postural evaluation. We calculated the lateral-medial index of each foot before and after the use of orthoses.

Table 1. Assessment of walking before and after the use of orthoses

	Before orthoses	After orthoses	p
Pain when walking* (EVA 0-10)	5 [3-5,2]	0 [0-2]	0,002
Difficulty walking* (0-10)			
- In house	4 [3-5,2]	2 [0-2]	0,002
- Outside	6 [4,7-7]	2 [1,5-2,5]	0,02
10 Meter Walk test** (Normal comfortable speed)			
- Number of steps	18,64 $\pm$ 3,7	16,9 $\pm$ 5	0,2
- Duration (sec)	11,9 $\pm$ 4,6	11,8 $\pm$ 5,2	0,9
- Walking speed (m/min)	56,4 $\pm$ 17,7	58,6 $\pm$ 20,3	0,6
latero-medial (L/M) index**			
- L/M index of the right foot	1,18 $\pm$ 0,17	1,23 $\pm$ 0,23	0,1
- L/M index of the left foot	1,25 $\pm$ 0,17	1,26 $\pm$ 0,19	0,9

*median and quartile

**average and standard deviation

p significant if $< 0,05$

Results: A significant decrease in walking pain ($p = 0.002$) and difficulty walking ($p = 0.02$) was found with the use of orthoses. The variations in 10 meter walk test and dynamic baropodometric parameters were not significant ($p > 0,05$).

There were no significant correlations between pain and difficulty walking, the progression of RA, the duration of foot damage and the functional impact measured by the HAQ.

Conclusion: Thermoformable foot orthoses significantly reduced pain and difficulty walking. The absence of factors associated with pain and difficulty walking could possibly be related to the small sample size.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.5698

FRI0621-HPR CHANGES IN LOCUS OF CONTROL LEVEL IN PATIENTS WITH RHEUMATOID ARTHRITIS AFTER THE COURSE OF BIOFEEDBACK TRAINING

R. Grekhoff¹. ¹Zborovsky' Research Institute for Clinical and Experimental Rheumatology, Volgograd, Russian Federation

Background:

Objectives: Our aim was to study the effect of biofeedback (BFB) training on the locus of control in patients suffering from rheumatoid arthritis (RA), and to justify the use of this method in the complex treatment of the disease.

Methods: 40 RA patients hospitalized in the rheumatology department were examined. The average age of patients was 48.6 ± 7.73 years (from 30 to 70 years), women accounted for the majority = 26 (86%), the average duration of the disease was 12 ± 3.44 years. We use J. Rotter's Locus of Control Scale in E.F.Bazhin adaptation. RA patients were divided into two groups: the main (20 patients) and control (20 patients). Patients of the main group received complex therapy with 12 sessions of BFB training, mainly based on the parameters of the brain's electrical activity — EEG relaxation using the Reakor™ psychophysiological rehabilitation complex manufactured by Medicom MTD (Taganrog).

Results: We revealed externality in RA patients in the general field (3.03 ± 0.3), as well as in the field of relation to the disease (3.86 ± 0.23) and in the field of production relations (3.43 ± 0.25). After BFB trainings, an increase in internality was observed on the scales of the general sphere ($p < 0.05$) and attitude to the disease ($p < 0.01$) in patients of the main group. In the group of patients receiving conventional treatment, the dynamics of the results was unreliable.

Conclusion: It should be noted that the locus of control (or subjective control) is a quality that characterizes a person's tendency to attribute responsibility for the results of his activity to external forces, or to his own abilities and efforts. Externality is manifested when people prefer to shift responsibility for important events of their life to external circumstances, and external forces (bosses, colleagues, etc.). In the field of attitude to the disease, externality is manifested when patient behaves passively, and believes that he cannot influence the course of the disease in any way, shifting all responsibility for the treatment results to medical staff, which can lead to non-compliance with the treatment regimen and an increase in the level of anxiety and depression, decreased self-esteem. The onset of the disease and its associated social consequences (disability, loss of social roles, etc.) can cause a negative mental state of learned helplessness. Learned helplessness is defined as a condition that occurs as a result of uncontrolled, mainly negative events, which manifests itself in violations of emotional, motivational and cognitive processes. In other words, RA patient suffering from this condition expects treatment failures and reduces control over compliance with the treatment regimen. BFB therapy can be used in order to correct and prevent the state of learned helplessness by increasing the level of internality.

It is assumed that increasing internality in the BFB process is associated with teaching the patient the skills of self-regulation of physiological processes. The mechanisms of BFB therapeutic effect are not only changes in physiological parameters (improvement of cerebral and peripheral blood flow, muscle relaxation, and improvement of sleep) but also in a shift in the locus of control from external to internal, which can increase compliance, reduce neurotic complaints, mobilize volitional potential and improve patient self-esteem.

As a result of BFB course, an increase in the internality was noted in patients on the scales of the general sphere and the sphere of attitude to the disease. It is advisable to use the BFB to increase the compliance and effectiveness of complex treatment of RA patients.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2020-eular.1571

FRI0622-HPR IMPACT OF A PHARMACIST'S INTERVENTION ON THE KNOWLEDGE OF BIOLOGICS AND ADHERENCE IN PATIENTS WITH SPONDYLOARTHRITIS: A RANDOMIZED, OPEN-LABEL, CONTROLLED TRIAL

L. Gutermann¹, S. Dumas², C. López-Medina³, L. Boissinot⁴, C. Cotteret⁵, V. Perut⁶, A. Moltó⁷, O. Conort⁸, M. Dougados⁹. ¹Hôpital Marie Lannelongue, Pharmacy, Le Plessis Robinson, France; ²Hôpital Cochin - APHP, Pharmacy,