SAT0110

SARCOPENIA IN PATIENTS WITH RHEUMATOID ARTHRITIS ON THE TREATMENT WITH BIOLOGIC DISEASE MODIFYING ANTI-RHEUMATIC DRUGS

E. Hasegawa^{1,2}, S. Ito¹, Y. Kurosawa^{1,2}, S. Taniguchi¹, D. Kobayashi^{1,2}, A. Abe¹, H. Otani¹, K. Nakazono¹, A. Murasawa¹, I. Narita², H. Ishikawa¹. ¹Niigata Rheumatic Center, Department of Rheumatology, Niigata, Japan; ²Niigata University Graduate School of Medical and Dental Sciences, Division of Clinical Nephrology and Rheumatology, Niigata, Japan

Background: Sarcopenia is characterized by loss of muscle mass and strength, which lead to lower physical ability, less quality of life (QoL), frailty and mortality. Rheumatoid arthritis (RA) is considered to be one of the causes of sarcopenia.

Objectives: To clarify the effectiveness of biologic disease modifying anti-rheumatic drugs (bDMARDs) on sarcopenia, including physical ability, body composition and nutritional status.

Methods: This is a prospective cohort study including consecutive 48 patients (male 11, female 37, age 64.2±15.1) with RA who started bDMARDs in Niigata Rheumatic Center. Diagnosis of sarcopenia was according to the diagnostic algorithm of European Working Group on Sarcopenia in Older People (EWGSOP). We monitored disease activity of RA, physical ability, body composition, nutritional status and QoL at baseline, 6 months and at 12 months. Disease activity was measured by disease activity score-28 joint count based on erythrocyte sedimentation rate (DAS28-ESR), clinical disease activity index (CDAI). Physical activity was measured by Health Assessment Questionnaire (HAQ), 10m walking test (10MWT). Nutritional status was measured by controlling nutrition status (CONUT) score, and prognostic nutritional index (PNI). Overall QoL was measured by Euro-QoL 5 dimensions (EQ5D).

Results: Among 48 patients who started bDMARDs, 21 patients were classified as having sarcopenia. The bDMARDs used were adalimumab in 10 cases, certolizumab pegol in 9 cases, abatacept in 9 cases, golimumab in 7 cases, tocilizumab in 5 cases, infliximab in 5 cases and etanercept in 3 cases. DAS28-ESR (4.7 ± 1.4 vs. 2.7 ± 1.0 , $p < 0.001$) and CDAI (18.4 ± 9.4 vs. 7.4 ± 5.5 , $p < 0.001$) were significantly decreased by 12 months of bDMARDs therapy. Physical activity was significantly ameliorated after 12 months of bDMARDs; HAQ (1.1 ± 0.9 vs. 0.6 ± 0.8 , $p < 0.001$), 10MWT (1.5 ± 0.7 m/s vs. 1.7 ± 0.6 , $p = 0.002$). EQ-5D was also ameliorated (0.63 ± 0.15 vs. 0.74 ± 0.19 , $p = 0.0002$). As for body composition analysis, there were significant increase in body weight (54.6 ± 12.4 kg vs. 55.8 ± 13.6 , $p = 0.006$), but there was no significant increase in skeletal muscle mass index (5.9 ± 1.1 kg/m² vs. 5.9 ± 1.1 , $p = 0.229$). Among 21 patients who were classified as sarcopenia when starting bDMARDs, the number of patients having sarcopenia significantly decreased after 12 months of bDMARDs (100% vs. 52.3%, $p = 0.0005$) and skeletal muscle index of these patients were significantly increased (5.1 ± 0.5 kg/m² vs. 5.3 ± 0.7 , $p = 0.046$).

Conclusion: Twelve months of bDMARDs therapy significantly ameliorated disease activity, nutritional status and physical activity. In RA patients with sarcopenia, bDMARDs significantly increased skeletal muscle and may be effective for treatment of sarcopenia.

Disclosure of Interests: Eriko Hasegawa: None declared, Satoshi Ito Speakers bureau: Abbvie, Eisai, Yoichi Kurosawa: None declared, Shinji Taniguchi: None declared, Daisuke Kobayashi: None declared, Asami Abe: None declared, Hiroshi Otani: None declared, Kiyoshi Nakazono: None declared, Akira Murasawa: None declared, Ichiei Narita: None declared, Hajime Ishikawa: None declared
DOI: 10.1136/annrheumdis-2020-eular.1036

SAT0111

THE RELATIONSHIP BETWEEN THE ADMINISTRATION OF IL-6 INHIBITORS AND INSULIN RESISTANCE IN PATIENTS WITH ACTIVE RHEUMATOID ARTHRITIS

A. Jitaru¹, C. Pomirleanu², M. M. Leon-Constantin³, F. Mitu³, C. Ancuta². ¹Grigore T. Popa University of Medicine and Pharmacy, Faculty of Medicine, Iasi, Romania; ²Clinical Rehabilitation Hospital, 2nd Rheumatology Clinic, Iasi, Romania; ³Clinical Rehabilitation Hospital, Cardiovascular Rehabilitation Clinic, Iasi, Romania

Background: Rheumatoid arthritis (RA) is associated with an increased cardiovascular (CV) risk, due not only to the traditional risk factors (hypertension, insulin resistance/diabetes, obesity, smoking), but to the inflammatory status as well. The blockade of interleukin-6 (IL-6) can regulate the glucose metabolism, reducing the glucose level and insulin resistance (IR). This beneficial effect is seen more in patients with normal values of body mass index (BMI), compared to the obese population.

Objectives: Given the mentioned existing data, we aim to demonstrate the positive effect of IL-6 inhibitors in active RA patients with normal or increased BMI.

Methods: We recruited 56 consecutive patients with definite and active RA, non-responders/partial responders to conventional synthetic Drug Modifying Anti-Rheumatic Drugs (csDMARDs)/biological therapy. For a period of 52 weeks, patients received subcutaneous Tocilizumab (TCZ) in a dose of 162mg once a week, according to European League Against Rheumatism (EULAR) recommendation

and National Protocol. We assessed demographics, RA-related parameters (clinical, inflammatory and immune) and metabolic markers, as well as the peripheral response to insulin, quantified by Homeostasis Model Assessment for insulin resistance (HOMA-IR) and the Quantitative Insulin Sensitivity Check Index (QUICKI). We did not include in the study the patients known with diabetes mellitus (DM) and those undergoing glucocorticoids.

Results: After 52 weeks of treatment, most of the patients showed a statistically significant reduction of HOMA-IR (3.61 ± 1.21 at the onset vs. 2.45 ± 1.46 at the end of the study, $p < 0.001$), while QUICKI registered a slight increase (0.32 ± 0.01 at the onset vs. 0.33 ± 0.01 at the end of the study, $p < 0.001$). Also, the decrease in insulin and glucose levels were more obvious in patients with normal BMI, strictly related to disease activity.

Conclusion: Long-term administration of TCZ in active RA is associated with a significant reduction of disease activity and IR, especially in normal weight patients. This confirms that obesity, as a CV risk factor, represents one of the main causes of IR.

References:

- [1] Castañeda S, Remuzgo-Martínez S, López-Mejías R et al. Rapid beneficial effect of the IL-6 receptor blockade on insulin resistance and insulin sensitivity in non-diabetic patients with rheumatoid arthritis. *Clin Exp Rheumatol*. 2019; 37(3):465-473.
- [2] Lehrskov LL, Christensen RH. The role of interleukin-6 in glucose homeostasis and lipid metabolism. *Semin Immunopathol*. 2019; 41(4):491-499.
- [3] Ursini F, Russo E, Ruscitti P, Giacomelli R, De Sarro G. The effect of non-TNF-targeted biologics and small molecules on insulin resistance in inflammatory arthritis. *Autoimmun Rev*. 2018 Apr;17(4):399-404.

Disclosure of Interests: Alexandra Jitaru: None declared, Cristina Pomirleanu: None declared, Maria-Magdalena Leon-Constantin: None declared, Florin Mitu: None declared, CODRINA ANCUTA Consultant of: AbbVie, Pfizer, Roche, Novartis, UCB, Ewopharma, Merck Sharpe and Dohme, and Eli Lilly, Speakers bureau: AbbVie, Pfizer, Roche, Novartis, UCB, Ewopharma, Merck Sharpe and Dohme, and Eli Lilly

DOI: 10.1136/annrheumdis-2020-eular.6509

SAT0112

EFFECT OF RITUXIMAB ON IMMUNOGLOBULIN LEVELS AND RISK OF ASSOCIATED INFECTION

S. Khalid¹, R. Smith², L. Yalakk¹. ¹Basingstoke and North Hampshire Hospital, Rheumatology, Basingstoke, United Kingdom; ²Salisbury District Hospital, Salisbury, United Kingdom

Background: Rituximab (RTX) is an anti-CD20 monoclonal antibody that suppresses B-lymphocytes and may induce hypogammaglobulinemia. Studies have shown that sustained low levels of immunoglobulins (Ig) are associated with significantly increased risks of infections.

Objectives: To determine the relationship between the serum Ig levels and risk of infection during (RTX) therapy for rheumatic diseases. We also aimed to identify the most common type of infections and pathogens associated with them.

Methods: A multi-centre retrospective observational study of patients with autoimmune diseases treated with RTX between 2009-2019. Serum Ig levels (IgM, IgG and IgA) were measured at baseline and 6-12 months after each cycle and note was made of any hypogammaglobulinemia. Infections, evidenced by a positive microbiology sample, radiograph or requirement of antibiotics and/or hospitalization, were recorded.

Results: 146 patients were included with a mean age of 61.5 years (standard deviation 13.8). 105 patients (71.9%) had Rheumatoid Arthritis. 51/146 had had 1 or more episodes of infections whilst on Rituximab which required treatment. Of these, 26 (50.9%) had recurrent infections. 33/51 had low immunoglobulins of at least 1 type. After receiving RTX treatment, 14 had low IgG, 29 had low IgM and only 4 had low IgA. 6 patients with low IgG, but 13 patients with low IgM suffered with recurrent infections.

There was a statistically significant higher proportion of patients with infection who had low Ig levels compared to those with normal levels, with a $p < 0.01$ on chi square testing.

Lower respiratory tract infections were the commonest infection seen with 28 cases, with most cases (15/28) caused by Haemophilus Influenza. 18 cases of urinary tract infections were seen, with 12/18 being caused by E. Coli. 3 cases of shingles were also seen.

Conclusion: Our study shows that an increased risk of infection was associated with hypogammaglobulinemia after rituximab therapy and highlights the importance of monitoring these patients. In our data set, more patients with recurrent infections had low IgM, supporting a need for better understanding of low IgM and its relation to infection. Respiratory tract infections were the most common infection with Haemophilus Influenza being the commonest pathogen. Recent studies have shown an increase in haemophilus influenza infections. Reasons for the increase may include a waning immunity to Haemophilus, changes in the organism and greater numbers of high-risk people- such as in our study. This supports a rationale