

# Immunogenicity and safety of the BNT162b2 mRNA COVID-19 vaccine in adult patients with autoimmune inflammatory rheumatic diseases and in the general population: a multicentre study

Victoria Furer <sup>1,2</sup>, Tali Eviatar,<sup>1,2</sup> Devy Zisman,<sup>3,4</sup> Hagit Peleg,<sup>5</sup> Daphna Paran,<sup>1,2</sup> David Levartovsky,<sup>1</sup> Michael Zisapel,<sup>1</sup> Ofir Elalouf,<sup>1,2</sup> Ilana Kaufman,<sup>1,2</sup> Roni Meidan,<sup>2,6</sup> Adi Broyde,<sup>1,2</sup> Ari Polachek,<sup>1,2</sup> Jonathan Wollman,<sup>1,2</sup> Ira Litinsky,<sup>1,2</sup> Katya Meridor,<sup>1,2</sup> Hila Nochomovitz,<sup>1,2</sup> Adi Silberman,<sup>1,2</sup> Dana Rosenberg,<sup>1,2</sup> Joy Feld,<sup>3</sup> Amir Haddad,<sup>3</sup> Tal Gazzit,<sup>3</sup> Muna Elias,<sup>3</sup> Nizar Higazi,<sup>3</sup> Fadi Kharouf,<sup>5,7</sup> Gabi Shefer,<sup>8</sup> Orly Sharon,<sup>8</sup> Sara Pel,<sup>1</sup> Sharon Nevo,<sup>1</sup> Ori Elkayam<sup>1,2</sup>

**Handling editor** Josef S Smolen

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2021-220647>).

For numbered affiliations see end of article.

## Correspondence to

Dr Victoria Furer, Rheumatology Department, Tel Aviv University, Tel Aviv, Israel; [furer.rheum@gmail.com](mailto:furer.rheum@gmail.com)

Received 26 April 2021  
Accepted 27 May 2021  
Published Online First  
14 June 2021



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Furer V, Eviatar T, Zisman D, et al. *Ann Rheum Dis* 2021;**80**:1330–1338.

## ABSTRACT

**Introduction** Vaccination represents a cornerstone in mastering the COVID-19 pandemic. Data on immunogenicity and safety of messenger RNA (mRNA) vaccines in patients with autoimmune inflammatory rheumatic diseases (AIIRD) are limited.

**Methods** A multicentre observational study evaluated the immunogenicity and safety of the two-dose regimen BNT162b2 mRNA vaccine in adult patients with AIIRD (n=686) compared with the general population (n=121). Serum IgG antibody levels against SARS-CoV-2 spike S1/S2 proteins were measured 2–6 weeks after the second vaccine dose. Seropositivity was defined as IgG  $\geq$ 15 binding antibody units (BAU)/mL. Vaccination efficacy, safety, and disease activity were assessed within 6 weeks after the second vaccine dose.

**Results** Following vaccination, the seropositivity rate and S1/S2 IgG levels were significantly lower among patients with AIIRD versus controls (86% (n=590) vs 100%,  $p<0.0001$  and  $132.9\pm 91.7$  vs  $218.6\pm 82.06$  BAU/mL,  $p<0.0001$ , respectively). Risk factors for reduced immunogenicity included older age and treatment with glucocorticoids, rituximab, mycophenolate mofetil (MMF), and abatacept. Rituximab was the main cause of a seronegative response (39% seropositivity). There were no postvaccination symptomatic cases of COVID-19 among patients with AIIRD and one mild case in the control group. Major adverse events in patients with AIIRD included death (n=2) several weeks after the second vaccine dose, non-disseminated herpes zoster (n=6), uveitis (n=2), and pericarditis (n=1). Postvaccination disease activity remained stable in the majority of patients.

**Conclusion** mRNA BNTb262 vaccine was immunogenic in the majority of patients with AIIRD, with an acceptable safety profile. Treatment with glucocorticoids, rituximab, MMF, and abatacept was associated with a significantly reduced BNT162b2-induced immunogenicity.

## INTRODUCTION

The prevention of COVID-19 pandemic has become of paramount importance. BNT162b2, a messenger

## Key messages

### What is already known about this subject?

- Data on efficacy and safety of the SARS-CoV-2 BNT162b2 messenger RNA (mRNA) vaccine in patients with autoimmune inflammatory rheumatic diseases (AIIRD) are limited.

### What does this study add?

- This is the largest observational prospective study conducted to confirm immunogenicity of the BNT162b2 mRNA vaccine in the majority of patients with AIIRD compared with controls.
- Immunogenicity was severely impaired by rituximab; moderately impaired by glucocorticoids, abatacept, and mycophenolate mofetil; and mildly impaired by methotrexate.
- The vaccine was generally safe in terms of adverse events.
- Postvaccination disease activity remained stable in the majority of patients with AIIRD.

### How might this impact on clinical practice or future developments?

- Most disease-modifying antirheumatic drugs, including methotrexate, anticytokine biologics and Janus kinase inhibitors, can be continued with relation to the administration of the BNT162b2 mRNA vaccine.
- Postponing treatment with rituximab, when feasible, should be considered to improve immunogenicity. Holding treatment with mycophenolate mofetil and abatacept, especially when combined with methotrexate, may be considered on an individual basis.

RNA (mRNA)-based vaccine, has demonstrated a high efficacy rate with an acceptable safety profile.<sup>1,2</sup> A mass BNT162b2 vaccination campaign has been launched in Israel, with high uptake of vaccination in about 55.5% of the country's population. Patients with autoimmune inflammatory rheumatic diseases

**Table 1** Demographic characteristics of patients with AIIRD and controls

	Age, median (range)	Female n (%)	Disease duration, years*	Influenza vaccine n (%)†
Controls, n=121	50 (18–90)†*	78 (65)	NA	89 (82.4)
<b>AIIRD diagnosis, n</b>				
All patients with AIIRD, n=686	59 (19–88)	475 (69.3)	10 (0–68)	542 (79.4)
RA, n=263	64 (20–88)	215 (81.75)	10 (0–50)	213 (82.88)
PsA, n=165	55 (20–86)	78 (47.56)	8 (0–68)	120 (74.07)
AxSpA, n=68	49.5 (21–83)	36 (52.94)	10 (1–51)	54 (80.6)
SLE, n=101	46 (22–80)	89 (88.12)	14 (0–44)	76 (77.55)
IIM, n=19	64 (34–76)	14 (73.68)	2 (1–21)	19 (100)
<b>Vasculitis, n=70</b>				
LVV, n=21	70 (26–85)	17 (80.95)	2.5 (0–12)	20 (95.24)
AAV, n=26	60.5 (26–85)	14 (53.85)	4 (0.75–28)	22 (84.62)
Other vasculitis, n=23	56 (19–77)	12 (52.17)	6 (0.5–35)	18 (78.26)

\*Data on disease duration were available for 683 patients with AIIRD (they were missing for two patients with PsA and one patient with SLE).

†Data on influenza vaccination were available for 781 participants: 673 AIIRD and 108 controls.

‡p<0.0001.

AAV, antineutrophil cytoplasmic antibody (ANCA)-associated vasculitis; AIIRD, autoimmune inflammatory rheumatic diseases; AxSpA, axial spondyloarthritis; IIM, idiopathic inflammatory myositis; LVV, large vessel vasculitis; PsA, psoriatic arthritis; RA, rheumatoid arthritis; SLE, systemic lupus erythematosus.

(AIIRD) have been prioritised for urgent vaccination to mitigate COVID-19 risk, consistent with the American College of Rheumatology (ACR) guidelines,<sup>3</sup> despite a paucity of data on the efficacy and safety of mRNA COVID-19 vaccines in this population. Recently, some encouraging data on mRNA vaccination in immunosuppressed patients have emerged based on two small studies with a limited follow-up.<sup>4–6</sup> Therefore, we conducted a large prospective observational multicentre study to evaluate immunogenicity, efficacy, and safety of the BNT162b2 mRNA

vaccine in patients with AIIRD compared with control subjects without rheumatic diseases or immunosuppressive therapies.

## METHODS

This prospective observational exploratory multicentre study was conducted at the Rheumatology Departments of Tel Aviv Sourasky, Carmel, and Hadassah Medical Center, Israel, between December 2020 and March 2021.

### End points of the study

The primary end point was immunogenicity of the BNT162b2 mRNA vaccine in adult patients with AIIRD compared with controls measured 2–6 weeks after the second vaccine dose.

### Secondary end points included

1. Effect of immunosuppressive treatments on vaccine's immunogenicity.
2. Efficacy of vaccination, defined as prevention of COVID-19 disease, confirmed by a PCR testing.
3. Safety of vaccination in patients with AIIRD compared with controls.
4. Effect of vaccination on clinical disease activity in patients with AIIRD.

### Study population

Consecutive adult patients (aged ≥18 years) were recruited into the study according to the following inclusion criteria: rheumatoid arthritis (RA)/ACR/European League Against Rheumatism (EULAR) 2010 classification criteria<sup>7</sup>; psoriatic arthritis (PsA)/Classification Criteria for PsA<sup>8</sup>; axial spondyloarthritis (axSpA)/Assessment of SpondyloArthritis International Society classification criteria<sup>9</sup>; systemic lupus erythematosus (SLE)/1997 ACR<sup>10</sup> or 2012 Systemic Lupus Erythematosus International Collaborating Clinics criteria<sup>11</sup>; systemic vasculitis: large vessel vasculitis (LVV), antineutrophil cytoplasmic antibody-associated vasculitis (AAV), including granulomatosis with polyangiitis (GPA), microscopic polyangiitis and eosinophilic GPA/Chapel Hill Consensus Conference definitions<sup>12</sup>; central nervous system (CNS) vasculitis, including primary CNS vasculitis, neuro-Behcet and Susac syndrome; and idiopathic inflammatory myositis (IIM)/EULAR/ACR classification criteria.<sup>13</sup>

**Table 2** Treatments used in patients with AIIRD

AIIRD diagnosis, n	Immunosuppressive treatments, n (%)								
	GC	MTX	TNFi	IL6i	Anti-CD20	ABA	JAKi	IL17i	MMF
All AIIRD, n=686	130 (18.95)	176 (25.66)	172 (25.07)	37 (5.39)	87 (12.68)	16 (2.33)	49 (6.9)	48 (7)	28 (4.08)
RA, n=263	55 (20.91)	116 (44.11)	47 (17.87)	29 (11.03)	43 (16.35)	15 (5.7)	46 (16.9)	0	0
PsA, n=165	3 (1.82)	36 (21.82)	74 (44.85)	0	0	1 (0.61)	2 (1.2)	40 (24.24)	0
AxSpA, n=68	1 (1.47)	9 (13.24)	48 (70.59)	0	1 (1.47)*	0	0	8 (11.76)	2 (2.94)
SLE, n=101	22 (21.78)	8 (7.92)	0	0	7 (6.93)	0	0	0	17 (16.83)
IIM, n=19	15 (78.95)	2 (10.53)	0	0	13 (68.42)	0	0	0	6 (31.58)
LVV, n=21	11 (52.38)	2 (9.52)	1 (4.76)	8 (38.1)	0	0	0	0	0
AAV, n=26	12 (46.15)	2 (7.69)	0	0	18 (69.23)	0	0	0	0
Other vasculitis, n=23	11 (47.83)	1 (4.35)	2 (8.7)	0	5 (21.74)	0	0	0	3 (13.04)

\*This patient had multiple sclerosis and was treated with ocrelizumab.

AAV, antineutrophil cytoplasmic antibody (ANCA)-associated vasculitis; ABA, abatacept; AIIRD, autoimmune inflammatory rheumatic diseases; anti-CD20, CD-20 inhibitors; AxSpA, axial spondyloarthritis; GC, glucocorticoids; IIM, idiopathic inflammatory myositis; IL6i, interleukin 6 inhibitors; IL17i, interleukin 17 inhibitors; JAKi, Janus kinase inhibitors; LVV, large vessel vasculitis; MMF, mycophenolate mofetil; MTX, methotrexate; PsA, psoriatic arthritis; RA, rheumatoid arthritis; SLE, systemic lupus erythematosus; TNFi, tumour necrosis factor inhibitors.

**Table 3** Immunogenicity of the BNT162b2 messenger RNA vaccine in patients with AIIRD and controls

Study participants, n	Seropositivity rate, n (% of total)	Serum anti-S1/S2 IgG titre, mean±SD, BAU/mL
Controls, n=121	121 (100)	218.6±82.06
Patients with AIIRD, n=686	590 (86.0)*	132.9±91.7*
RA, n=263	216 (82.1)	108.7±84.7
PsA, n=165	160 (96.9)	162.0±71.7
AxSpA, n=68	67 (98.5)	173.1±90.1
SLE, n=101	93 (92.1)	161.9±105.2
IIM, n=19	7 (36.8)	42.9±62.6
LVV, n=21	20 (95.2)	143.3±84.6
AAV, n=26	8 (30.8)	40.3±73.2
Other vasculitis, n=23	19 (86.6)	122.7±87.9

\*p&lt;0.0001.

AAV, antineutrophil cytoplasmic antibody (ANCA)-associated vasculitis; AIIRD, autoimmune inflammatory rheumatic diseases; AxSpA, axial spondyloarthritis; BAU, binding antibody units; IIM, idiopathic inflammatory myositis; LVV, large vessel vasculitis; PsA, psoriatic arthritis; RA, rheumatoid arthritis; SLE, systemic lupus erythematosus.

Patients were instructed to continue all medications during the vaccination period, except for rituximab treatment that was delayed after the vaccination in certain cases on a physician's discretion.

The control group included a sample of the general population, consisting mainly of healthcare personnel. Exclusion criteria for all groups were pregnancy, history of past vaccination allergy, and previous COVID-19 infection and for controls—history of AIIRD and immunosuppressive treatment.

### Vaccination procedure

All study participants were administered the two-dose regimen BNT162b2 mRNA vaccine (Pfizer-BioNTech), 30 µg per dose, by intramuscular injection in the deltoid muscle 3 weeks apart, as indicated by the national guidelines.

### Immunogenicity of the vaccine

The vaccine immunogenicity was evaluated by measuring the serum IgG neutralising antibody levels against SARS-CoV-2 trimeric spike S1/S2 glycoproteins, using the LIAISON (DiaSorin) quantitative assay, performed 2–6 weeks after the second vaccine dose. This Food and Drug Administration-authorized assay has a clinical sensitivity and specificity above 98%.<sup>14</sup> A value above 15 binding antibody units (BAU) was considered as positive, according to the manufacturer's instruction.

### Efficacy of the vaccine

The participants were questioned whether they contracted COVID-19 infection, confirmed by PCR, following each vaccine dose. In addition, up to the data cut-off, the patient files were reviewed for evidence of COVID-19 infection.

### Safety of the vaccine

The participants were contacted by phone within 2 weeks after the first vaccine dose and within 2–6 weeks after the second vaccine dose to complete a questionnaire regarding adverse events.

### Clinical assessment of AIIRD

Medical history and the use of medications were recorded. Data regarding disease activity before vaccination were retrieved from patients' medical records, within up to 3 months before vaccination. Postvaccination disease activity was assessed by an in-person clinical examination within 2–6 weeks after the second vaccine dose. The following disease activity indices were included: Clinical Disease Activity Index, Simplified Disease Activity Index, DAS-28-CRP for RA, Disease Activity in Psoriatic Arthritis, Leeds Enthesitis and Dactylitis Index, Psoriasis Area Severity Index for PsA, Bath Ankylosing Spondylitis Disease Activity Index and Ankylosing Spondylitis Disease Activity Score for axSpA, Systemic Lupus Disease Activity Index for SLE, and patients' and physician's global assessment, using a visual analogue scale of 0–10 mm, for vasculitis and inflammatory myositis.


### Patient and public involvement

The research question and outcome measures of this study were developed in collaboration with the representatives of patients with AIIRD based on a shared priority to investigate the efficacy and safety of the novel mRNA BNT162b2 vaccine. Patients with AIIRD under the care of the medical centres conducting the trial were actively informed regarding the study and offered to participate. In view of the ongoing COVID-19 pandemic and related

**Table 4** Immunogenicity of the BNT162b2 messenger RNA vaccine according to the use of immunosuppressive treatments in comparison with controls

Immunosuppressive treatments, n	Seropositivity rate, n (%)	P value
GC, n=130	86 (66)	<0.0001
GC monotherapy, n=13	10 (77)	<0.0001
MTX, n=176	148 (84)	<0.0001
MTX monotherapy, n=41	38 (92)	0.02
HCO, n=133	120 (90)	0.001
HCO monotherapy, n=50	49 (98)	0.65
LEF, n=28	25 (89)	0.004
LEF monotherapy, n=11	11 (100)	NA
TNFi, n=172	167 (97)	0.15
TNFi monotherapy, n=121	119 (98)	0.48
TNFi+MTX, n=29	27 (93)	0.04
IL6i, n=37	37 (100)	NA
IL6i monotherapy, n=19	19 (100)	NA
IL6i+MTX, n=7	7 (100)	NA
Anti-CD20, n=87	36 (41)	<0.0001
Anti-CD20 monotherapy, n=28	11 (39)	<0.0001
Rituximab+MTX, n=14	5 (36)	<0.0001
IL17i, n=48	47 (98)	0.63
IL17i monotherapy, n=37	37 (100)	NA
IL17i+MTX, n=7	6 (85)	0.05
Abatacept, n=16	10 (62)	<0.0001
Abatacept monotherapy, n=7	5 (71)	<0.0001
Abatacept+MTX, n=5	2 (40)	<0.0001
JAKi monotherapy, n=21	19 (90)	0.02
JAK+MTX, n=24	22 (92)	0.03
Belimumab, n=9	7 (77)	0.0001
MMF, n=28	18 (64)	<0.0001

anti-CD20, CD20 inhibitors; GC, glucocorticoids; HCO, hydroxychloroquine; IL6i, interleukin 6 inhibitors; IL17i, interleukin 17 inhibitors; JAKi, Janus kinase inhibitors; LEF, leflunomide; MMF, mycophenolate mofetil; MTX, methotrexate; TNFi, tumour necrosis factor inhibitors.



**Figure 1** Cumulative seropositive rate according to the interval (days) between the last course of rituximab administration and BNT162b2 vaccination. mRNA, messenger RNA.

stringent restrictions, patients were not involved in the conduct of the study. The main study results will be disseminated to the participants, and we will seek patient and public involvement in the development of an appropriate method of dissemination.

### Statistical analysis

Differences between continuous variables were tested for significance using the independent-samples t-test. Differences between categorical variables were tested for significance using the  $\chi^2$  test or Fisher's exact test (as appropriate). Multivariate models (linear and logistic) were adjusted for age, diagnosis, treatment with methotrexate and anti-CD20. All tests applied were two-tailed. The widths of the intervals have not been adjusted for multiplicity, and the inferences drawn from inferences may not be reproducible. Missing data were assumed as missing at random. No imputations were done. The data were analysed using R V.4.0.5 (R Development Core Team, Vienna, Austria).

## RESULTS

### Study population

A total of 710 patients with AIIRD and 124 controls vaccinated with the two-dose regimen BNT162b2 mRNA vaccine were enrolled in the study. The final analysis included 686 patients with AIIRD and 121 controls due to missing serology tests (table 1; online supplemental figure S1). RA was the most common disease (n=263), followed by PsA (n=165), SLE (n=101), systemic vasculitis (n=70), axSpA (n=68) and IIM (n=19). Patients with AIIRD included a subgroup of elderly patients aged  $\geq 65$  years (32.8%, n=225) and were significantly older than controls, mean age $\pm$ SD 56.76 $\pm$ 14.88 vs 50.76 $\pm$ 14.68, respectively;  $p < 0.0001$ .

A total of 95.2% (n=653) of patients with AIIRD were treated with immunomodulatory medications (table 2). Glucocorticoids (GC) were used in 18.95% (n=130), at a mean prednisone dose of 6.7 $\pm$ 6.25 mg/day. Conventional synthetic disease-modifying antirheumatic drug (csDMARD) monotherapy was used in 23.18% (n=159). Biologic DMARDs were used as a monotherapy or in combination

with csDMARDs in 38.19% (n=262) and 13.56% (n=93), respectively. Janus kinase inhibitors (JAKi) were used as a monotherapy or in combination with csDMARDs in 3.06% (n=21) and 3.79% (n=26), respectively. Eighty-seven (12.68%) patients were treated with CD20-depleting (anti-CD20) therapies, of whom 86 received rituximab at a mean dose of 1656.1 $\pm$ 623.6 mg. The mean interval between the last dose of rituximab and BNT162b2 vaccination was 51 $\pm$ 83 days. One patient received ocrelizumab. During the study period, changes in immunomodulatory drugs after the first vaccine dose were reported in 3% (n=20) of patients and after the second vaccine dose in 4.04% (n=27).

### Immunogenicity of the BNT162b2 vaccine

The seropositivity rate was 86% (n=590) in patients with AIIRD compared with 100% in controls ( $p < 0.0001$ ). The level of the S1/S2 antibodies was significantly reduced in patients with AIIRD compared with controls (mean $\pm$ SD, 132.9 $\pm$ 91.7 vs 218.6 $\pm$ 82.06;  $p < 0.0001$ ). In patients with PsA, axSpA, SLE and LVV, the seropositive rate was above 90%. In patients with RA, the seropositive rate was 82.1%, whereas the lowest seropositive rate (<40%) was observed in patients with AAV and IIM (table 3).

### Effect of immunosuppressive treatments on the immunogenicity of the BNT162b2 vaccine

More than 97% of patients treated with anticytokine therapies, including tumour necrosis factor inhibitors (TNFi), interleukin 17 inhibitors (IL-17i) and interleukin 6 inhibitors (IL-6i), had an appropriate immunogenic response when used as monotherapy (table 4). Anti-CD20 significantly impaired vaccine's immunogenicity, with the lowest seropositivity rate of 39%. The time interval between the prevaccination administration of rituximab and the BNT162b2 vaccination had a significant impact on the vaccine's immunogenicity, as shown in figure 1. The seropositivity rate in patients vaccinated within 6 months after rituximab treatment was below 20% but increased to about 50% in patients vaccinated 1 year after rituximab treatment. Similarly, the use of

**Table 5** Unadjusted and adjusted logistic regression models examining the factors associated with seropositivity

	Seropositivity rate, n (%)	Unadjusted OR (95% CI)	Adjusted OR (95% CI)	P value
Age >65 years, n=246	195 (79.27)	0.33 (0.22 to 0.52)	0.43 (0.25 to 0.75)	<b>0.002</b>
<b>AIIRD diagnosis</b>				
Psa, n=165	160 (96.97)	Reference	Reference	
RA, n=263	216 (82.13)	0.14 (0.06 to 0.37)	0.31 (0.11 to 0.82)	<b>0.02</b>
AxSpA, n=68	67 (98.53)	2.09 (0.24 to 18.26)	2.01 (0.23 to 17.72)	0.52
SLE, n=101	93 (92.08)	0.36 (0.12 to 1.14)	0.35 (0.11 to 1.16)	0.08
IIM, n=19	7 (36.84)	0.02 (0.01 to 0.07)	0.06 (0.02 to 0.27)	<b>&lt;0.001</b>
LVV, n=21	20 (95.24)	0.63 (0.07 to 5.63)	0.82 (0.09 to 7.54)	0.86
AAV, n=26	8 (30.77)	0.01 (0.004 to 0.05)	0.04 (0.01 to 0.17)	<b>&lt;0.001</b>
Other vasculitis, n=23	19 (82.61)	0.15 (0.04 to 0.6)	0.26 (0.06 to 1.22)	0.09
<b>AIIRD treatments</b>				
Anti-CD20, n=87	36 (41.38)	0.05 (0.03 to 0.08)	0.13 (0.07 to 0.24)	<b>&lt;0.001</b>
Anti-CD20 monotherapy, n=28	11 (39.29)	0.07 (0.03 to 0.16)	0.92 (0.33 to 2.57)	0.87
Anti-CD20 +MTX, n=14	5 (35.71)	0.07 (0.02 to 0.21)	0.94 (0.23 to 3.89)	0.93
MTX, n=176	148 (84.09)	0.64 (0.4 to 1.03)	0.58 (0.31 to 1.07)	0.08
MTX monotherapy, n=41	38 (92.68)	1.75 (0.53 to 5.79)	1.84 (0.5 to 6.74)	0.36
GC, n=130	86 (66.15%)	0.16 (0.1 to 0.29)	0.48 (0.26 to 0.87)	<b>0.02</b>
TNFi, n=172	167 (97.09)	5.6 (2.24 to 14.0)	1.89 (0.68 to 5.24)	0.22
TNFi monotherapy, n=121	119 (98.35)	9.46 (2.3 to 38.87)	2.58 (0.56 to 11.94)	0.22
TNFi +MTX, n=29	27 (93.1)	1.86 (0.44 to 7.94)	1.46 (0.31 to 6.91)	0.63
IL6i, n=37	37 (100)	NA	NA	NA
IL6i monotherapy, n=19	19 (100)	NA	NA	NA
IL6i+MTX, n=7	7 (100)	NA	NA	NA
IL17i, n=48	47 (97.92)	6.73 (0.92 to 49.32)	1.42 (0.16 to 12.83)	0.75
IL17 monotherapy, n=37	37 (100)	NA	NA	NA
IL17 +MTX, n=7	6 (85.71)	0.81 (0.1 to 6.8)	0.25 (0.02 to 2.7)	0.25
Abatacept, n=16	10 (62.5)	0.21 (0.08 to 0.6)	0.14 (0.04 to 0.43)	<b>&lt;0.001</b>
Abatacept monotherapy, n=7	5 (71.43)	0.33 (0.06 to 1.74)	0.2 (0.033 to 1.16)	0.073
Abatacept+MTX, n=5	2 (40)	0.09 (0.01 to 0.53)	0.07 (0.01 to 0.48)	<b>0.007</b>
JAKi monotherapy, n=21	19 (90.48)	1.29 (0.3 to 5.63)	0.72 (0.15 to 3.48)	0.68
JAKi+MTX, n=24	22 (91.67)	1.5 (0.35 to 6.48)	1.78 (0.38 to 8.35)	0.46
MMF, n=28	18 (64.29)	0.22 (0.1 to 0.5)	0.1 (0.03 to 0.34)	<b>0.0013</b>
MMF monotherapy, n=5	3 (60)	0.2 (0.03 to 1.21)	0.11 (0.02 to 0.83)	<b>0.03</b>

AAV, antineutrophil cytoplasmic antibody (ANCA)-associated vasculitis; AIIRD, autoimmune inflammatory rheumatic diseases; anti-CD20, CD20 inhibitors; AxSpA, axial spondyloarthritis; GC, glucocorticoids; IIM, idiopathic inflammatory myositis; IL6i, interleukin 6 inhibitors; IL17i, interleukin 17 inhibitors; JAKi, Janus kinase inhibitors; LVV, large vessel vasculitis; MMF, mycophenolate mofetil; MTX, methotrexate; PsA, psoriatic arthritis; RA, rheumatoid arthritis; SLE, systemic lupus erythematosus; TNFi, tumour necrosis factor inhibitors.

GC, MMF, and abatacept was significantly associated with a lack of humoral response. Seropositivity rate in patients treated with MTX monotherapy and in combination with other treatments was significantly reduced (92% and 84%, respectively), although at a lesser magnitude than with anti-CD20, MMF, and abatacept treatments. The combination of TNFi with MTX reduced the rate of seropositivity to 93% ( $p=0.04$ ).

In the univariate logistic regression model (table 5), age >65 years, diagnosis of RA, IIM, and AAV and treatment with GC, MMF, and anti-CD20, and abatacept were associated with a lack of humoral response to vaccination. Multivariate regression analysis (using PsA, the largest subgroup with the highest seropositivity, as a reference) accounting for age, AIIRD diagnosis, and treatment with MTX and anti-CD20 confirmed these associations (table 5). The impact of GC, MMF, anti-CD20 and abatacept on immunogenicity was independent from the concomitant use of other DMARDs (data not shown).

### Efficacy of the BNT162b2 vaccine

There were no COVID-19 symptomatic disease among AIIRD patients during the study follow-up, whereas one subject in the control group was diagnosed with mild COVID-19 after the second vaccine dose.

### Safety of the BNT162b2 vaccine

The prevalence of mild adverse events was similar in patients with AIIRD and controls. There were no serious or major adverse events in the control group. Two patients with AIIRD died after the second vaccine dose. The first patient had a history of AAV, in remission and without any immunosuppressive therapy for 3 years before the vaccination, apart from a low-dose prednisone. Three weeks after the second vaccine dose, she developed fulminant haemorrhagic cutaneous vasculitis with subsequent fatal sepsis. The second patient suffered from PsA which was in remission under treatment with secukinumab and had multiple comorbidities, including diabetes mellitus and ischaemic heart disease. He

**Table 6** Adverse events of the BNT162b2 vaccine in patients with AIIRD and controls

Adverse event	After the first vaccine dose		After the second vaccine dose					
	Controls n=121	AIIRD n=673	Controls n=121	AIIRD n=670				
<b>Local reactions, n (%)</b>								
Pain	69 (57.02)	377 (56.02)	51 (42.5)	314 (46.87)				
Erythema	4 (3.31)	12 (1.78)	6 (5)	10 (1.49)*				
Swelling	6 (4.69)	18 (2.68)	6 (5)	15 (2.24)				
Pruritus	3 (2.48)	8 (1.19)	2 (1.67)	4 (0.6)				
Tingling	7 (5.79)	3 (0.45)**	1 (0.83)	0				
<b>Systemic reactions, n (%)</b>								
Fever $\geq 38.0^{\circ}\text{C}$	1 (0.83)	8 (1.19)	6 (4.96)	35 (5.24)				
Nausea	0	7 (1.04)	2 (1.67)	14 (2.09)				
Vomiting	1 (0.83)	3 (0.45)	0	1 (0.15)				
Rhinorrhoea	3 (2.48)	0 *	0	1 (0.15)				
Cough	1 (0.83)	2 (0.3)	0	1 (0.15)				
Myalgia	5 (4.13)	25 (3.71)	21 (17.36)	63 (9.4)*				
Arthralgia	1 (0.83)	23 (3.42)	6 (4.96)	49 (7.32)				
Chills	2 (1.65)	13 (1.93)	21 (17.36)	60 (8.96)*				
Malaise	1 (0.83)	13 (1.93)	21 (17.36)	53 (7.91)*				
Headache	7 (5.97)	47 (6.98)	18 (14.88)	85 (12.69)				
Allergic reaction	0	0	0	1 (0.15)				
Lethargy	6 (4.96)	36 (5.35)	10 (8.26)	90 (13.49)				
Worsening of rheumatological symptoms	NA	17 (2.53)	NA	12 (1.79)				
Other symptoms, n	Dizziness	2	Throat pain	5	Weakness	3	Weakness	28
	Numbness	1	Arm numbness	5	Dizziness	1	Dizziness	11
	Throat pain	1	Dizziness	4	Numbness	3	Throat pain	6
	Chest pain	1	Weakness	4	Facial pain	1	Excessive sweating	3
			Rash	2	Chest pain	2	Pericarditis	1
			Flu-like	2			Chest pain	1
			Diarrhoea	2			Local lymphadenopathy	2
			Pruritus	2			Vaginal bleeding	2
			Palpitations	2			Lack of appetite	2
			Uveitis	1			Diarrhoea	2
			Herpes Labialis	1			High blood pressure	2
			Other	3			Herpes zoster	6
							Uveitis	2
							Other	7
	Death	0	0	0	2 (0.3)			

\* $p \leq 0.05$ ; \*\* $p \leq 0.0001$ .

AIIRD, autoimmune inflammatory rheumatic diseases.

died from a myocardial infarction 2 months after the second vaccine dose. Adverse events of special interest in patients with AIIRD included uveitis ( $n=2$ ), herpes labialis ( $n=1$ ), pericarditis ( $n=1$ ), and non-disseminated herpes zoster (HZ)<sup>15</sup> in five patients after the first vaccine dose and in one patient after the second vaccine dose. One case included HZ ophthalmicus, without corneal involvement (table 6).

### BNT162b2 vaccine effect on disease activity in patients with AIIRD


In patients with RA, PsA, axSpA and SLE, the postvaccination indices of disease activity remained stable (figure 2).

### DISCUSSION

The current approach to COVID-19 vaccination of patients with AIIRD is mainly based on data extrapolated from studies on other vaccines. Herein, we report the results of the first large multicentre prospective study conducted during the COVID-19 pandemic demonstrating that BNT162b2 mRNA vaccine was immunogenic in the majority of patients with AIIRD compared

with controls, with a seropositivity rate of 86% vs 100%, respectively. S1/S2 IgG levels were significantly lower among patients with AIIRD compared with controls. These findings confirm the results reported by Geisen *et al*, where considerable immunogenicity was induced by anti-SARS-CoV-2 mRNA vaccines in a small group of patients with chronic inflammatory diseases.<sup>4</sup> The mean level of the anti-spike S1/S2 IgG neutralising antibodies measured 2–6 weeks after the second vaccine dose was significantly lower in patients with AIIRD compared with controls in all age groups, consistent with the response to a single dose of mRNA vaccines in patients with rheumatic disease reported by Boyarsky *et al*,<sup>5</sup> raising concerns about the long-term protection of the vaccine in patients with AIIRD.

Our study provided detailed information regarding the impact of various immunosuppressive treatments on vaccine-induced immunogenicity. GC are essential for many patients with AIIRD. The seropositivity rate of patients with AIIRD treated with GC was only 66%. The data are scarce regarding the pure effect of GC on vaccination response in patients with AIIRD, as GC are commonly used in combination with other immunosuppressants



**Figure 2** Disease activity scores before and after completing two doses of BNT162b2 vaccine. Data on prevaccination and postvaccination disease activity measures were available for 165 patients with RA-SDAI, 182 patients with RA-CDAI, 164 patients with RA-DAS-28-CRP, 121 patients with PsA-CDAI, 117 patients with PsA-DAPSA, 131 patients with PsA-PASI, 43 patients with AxSpA-ASDAS, 47 patients with AxSpA-BASDAI and 85 patients with SLE-SLEDAI. ASDAS, Ankylosing Spondylitis Disease Activity Score; axSpA, axial spondyloarthritis; BASDAI, Bath Ankylosing Spondylitis Disease Activity Index; CDAI, Clinical Disease Activity Index; DAPSA, Disease Activity in Psoriatic Arthritis; PASI, Psoriasis Area Severity Index; PsA, psoriatic arthritis; RA, rheumatoid arthritis; SDAI, Simplified Disease Activity Index; SLE, systemic lupus erythematosus; SLEDAI, Systemic Lupus Disease Activity Index for SLE.

and their doses tend to vary over the disease course. A dose of  $\geq 10$  mg/day was associated with a reduced vaccine-induced humoral response to pneumococcal vaccine in patients with various inflammatory diseases.<sup>16</sup> The mean GC dose in our study population was relatively low (6.2 mg/day), precluding analysis on the dose-dependent effect of GC on vaccination response. MTX represents a cornerstone medication in a spectrum of rheumatic diseases. MTX may reduce humoral response to influenza and pneumococcal vaccines in patients with RA.<sup>17–20</sup> Temporary discontinuation of MTX for 2 weeks after vaccination improved the immunogenicity of influenza vaccination in patients with RA.<sup>21–22</sup> In our study, the use of MTX as monotherapy or in combination with other treatments was mainly associated with a slightly reduced seropositivity and lower levels of the S1/S2 IgG antibodies compared with controls, suggesting no need for treatment modification with MTX in most cases of anti-COVID-19 vaccination. Holding of MTX may be considered if combined with abatacept or rituximab, in view of a prevalent negative serological response under these regimens. Importantly, anticytokine biologics including TNFi, IL17i and IL6i did not interfere with the production of BNTb262-induced antibodies. This observation is in line with the studies demonstrating a considerable immunogenicity induced by influenza and pneumococcal vaccines in patients with treated with TNFi,<sup>20–23–26</sup> IL-6i<sup>27–29</sup> and IL-17i.<sup>30–31</sup> JAKi, representing a smaller fraction of treatments in our study, demonstrated a minor non-significantly negative effect on the production of BNTb262-induced antibodies. Limited data regarding other vaccines have shown that patients with RA treated with tofacitinib achieved a considerable response to influenza vaccine but an impaired response to a pneumococcal vaccine, especially when combined with MTX.<sup>32</sup> Holding of tofacitinib for 1 week prevaccination and postvaccination had little impact on the immunogenicity of either vaccine.<sup>32</sup> A considerable pneumococcal humoral response was achieved in patients with RA treated with baricitinib.<sup>33</sup>

Treatment with anti-CD20 therapies, mainly represented by rituximab in this study, significantly reduced vaccine-induced humoral response, with seropositivity of 41.3% when administered as monotherapy and 36% when administered in

combination with MTX. The interval between the administration of rituximab and vaccination had a critical role in predicting the response to the vaccine. Our findings are in line with the previously published data regarding the negative impact of anti-CD20 therapy on the humoral response to various vaccines, although seroprotection could be still achieved after vaccination in rituximab-treated<sup>20–26–34–36</sup> and ocrelizumab-treated<sup>37</sup> patients. The degree of B-cell recovery at the time of vaccination correlated with the extent of the humoral response to vaccination, as reported for influenza vaccine in patients with RA treated with rituximab.<sup>38</sup> In a retrospective analysis of 30 patients with rheumatic diseases treated with rituximab, only 10 patients (33.3%) developed a serological response to anti-SARS-CoV-2 vaccination.<sup>39</sup> B-cell depletion was associated with a lack of serological response, based on data available for 11 patients.<sup>39</sup> Unfortunately, these data were not available for patients treated with anti-CD20 therapies in our study. As protection from SARS-CoV-2 relies on both humoral and T-cell-mediated immunity,<sup>40–41</sup> patients with a deficient humoral response may be still protected by the latter. Treatment with abatacept reduced vaccine-induced humoral response, with a seropositive rate of 71% as monotherapy, reduced to 40% when combined with MTX. Previous data regarding the impact of abatacept on other vaccination-related immunity are conflicting.<sup>29–42</sup> Treatment with MMF reduced humoral response to a seropositivity rate of 64% in 28 patients. Consistently, solid organ transplant recipients treated with regimens including MMF were at risk of a negative humoral response to mRNA SARS-CoV-2 BNT162b2 vaccine.<sup>43–44</sup>

From the standpoint of particular AIIRD at risk of low immunogenic response to vaccine, RA, AAV and IIM were associated with a low humoral response to the vaccine. This finding seems to be at least partially explained by the underlying treatment.

Regarding the safety of vaccination, our study provides a reassurance for a good safety profile of the vaccine, with most adverse events being transient and mild, consistent with two other studies.<sup>4–6</sup> No causal link between the two deaths of the patients with AIIRD and vaccination could be established. Other rare adverse events in patients with AIIRD were limited

in number and did not seem to cause long-term complications. Occurrence of HZ in six patients with AIIRD merits a special attention as reported by our group.<sup>15</sup> In this study, the affected patients were women within the age range of 36–61 years (mean, 49±11 years) with a mild or stable rheumatic disease. Two patients were treated with JAKi and one with rituximab and MMF, indicating a baseline increased risk for HZ, whereas three others had a low level of immunosuppression, calling for a potential causal link between the events.<sup>15</sup> As the occurrence of HZ was not specifically captured in the mRNA vaccine clinical trials, no data are available on the postvaccination HZ prevalence in the general population. Case reports on HZ following the BNT162b2 mRNA vaccination in subjects without immunosuppressive treatment<sup>45,46</sup> and in one patient with AAV in remission<sup>47</sup> were recently published. Further epidemiological studies and surveillance programmes are needed to investigate the prevalence of HZ in vaccinated subjects.

There was no evidence of significant disease flares across different AIIRD. Yet, this should be interpreted with a certain caution due to an exploratory analysis of disease activity assessment performed within a variable prevaccination and postvaccination time frame.

The limitations of our study include a non-randomised design, a lack of matching between patients and controls by age, and the absence of long-term follow-up data. Neither data on B-cell repopulation at the time of vaccination for patients under anti-CD20 therapy nor data on cellular immunity were available.

In summary, the data presented in this study have important implications for the management of anti-COVID-19 vaccination in patients with a wide spectrum of AIIRD. Most immunosuppressive treatments, including csDMARDs, anticytokine biologics and JAKi, can be safely continued without significantly attenuating vaccine-induced immunogenicity. The results of our study do not support withholding MTX and JAKi in relation to COVID-19 vaccination as recommended by the ACR.<sup>3</sup>

Treatment with GC, rituximab, abatacept in combination with MTX, and MMF was associated with significantly decreased vaccine-induced immunogenicity. Therefore, timing of vaccination has a critical role in these cases. Postponing administration of rituximab and abatacept, especially when combined with methotrexate, when clinically feasible, seems to be reasonable to improve vaccine-induced immunogenicity. Yet, the absence of a humoral response does not preclude T-cell-mediated vaccine-induced immunity, and if shown effective, may serve as a rationale for anti-COVID-19 vaccination of these patients. Importantly, this study provides evidence of overall good tolerance of the BNT162b2 mRNA vaccine in adult patients with AIIRD. Further studies are needed to assess the durability of the humoral vaccination response, T-cell-mediated immunity in patients with a poor humoral response and long-term efficacy and safety of vaccination in patients with AIIRD.

#### Author affiliations

<sup>1</sup>Rheumatology Department, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel

<sup>2</sup>Tel Aviv University Sackler Faculty of Medicine, Tel Aviv, Israel

<sup>3</sup>Rheumatology Unit, Carmel Hospital, Haifa, Israel

<sup>4</sup>Medicine, Technion Israel Institute of Technology, Haifa, Israel

<sup>5</sup>Rheumatology Unit, Hadassah University Hospital, Jerusalem, Israel

<sup>6</sup>Internal Medicine, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel

<sup>7</sup>The Hebrew University of Jerusalem Faculty of Medicine, Jerusalem, Israel

<sup>8</sup>Endocrinology Department, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel

**Correction notice** This article has been corrected since it published Online First. The author affiliations have been amended.

**Acknowledgements** We thank Mr Yoram Neufeld and Ms Maayan Pel for their assistance with data management and Mr Yishai Friedlander, MPH, for performing the statistical analysis.

**Contributors** The study was designed, directed and coordinated by OE, the principle investigator. VF, TE, DZ and HP, the sub-investigators, were in charge of the study conducted at all stages. All the MD co-authors recruited participants into the study and evaluated predisease and postdisease activity measures in patients. GS and OS performed the serology tests. SP and SN served as main study coordinators and questioned the study participants regarding the adverse events of vaccination. OE, VF and TE had full access to the study's data and wrote the article, which was critically reviewed by DP, DZ and HP.

**Funding** The study was funded by the department's fund of each medical centre participating in the study.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** The study was performed in accordance with the principles of the Declaration of Helsinki and approved by the research ethics committees of the three medical centres: TLV-1055-20, CMC-0238-20 and HMO-0025-21, respectively. The participants signed an informed consent on recruitment into the study.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article or uploaded as supplementary information. The data used to support the findings of this study are included within the article.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

#### ORCID iD

Victoria Furer <http://orcid.org/0000-0001-5193-4207>

#### REFERENCES

- Polack FP, Thomas SJ, Kitchin N, *et al.* Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N Engl J Med* 2020;383:2603–15.
- Dagan N, Barda N, Kepten E, *et al.* BNT162b2 mRNA Covid-19 vaccine in a nationwide mass vaccination setting. *N Engl J Med* 2021;384:1412–23.
- Force D. COVID-19 vaccine clinical guidance summary for patients with rheumatic and musculoskeletal diseases.
- Geisen UM, Berner DK, Tran F. Immunogenicity and safety of anti-SARS-CoV-2 mRNA vaccines in patients with chronic inflammatory conditions and immunosuppressive therapy in a monocentric cohort. *Ann Rheum Dis* 2021;80:1302–7.
- Boyersky BJ, Ruddy JA, Connolly CM, *et al.* Antibody response to a single dose of SARS-CoV-2 mRNA vaccine in patients with rheumatic and musculoskeletal diseases. *Ann Rheum Dis* 2021;80:1098–9.
- Connolly CM, Ruddy JA, Boyarsky BJ, *et al.* Safety of the first dose of mRNA SARS-CoV-2 vaccines in patients with rheumatic and musculoskeletal diseases. *Ann Rheum Dis* 2021;80:1100–1.
- Aletaha D, Neogi T, Silman AJ. Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League against rheumatism collaborative initiative. *Arthritis Rheum* 2010;2010:2569–81.
- Taylor W, Gladman D, Helliwell P, *et al.* Classification criteria for psoriatic arthritis: development of new criteria from a large international study. *Arthritis Rheum* 2006;54:2665–73.
- Rudwaleit M, van der Heijde D, Landewé R, *et al.* The development of assessment of spondyloarthritis International Society classification criteria for axial spondyloarthritis (Part II): validation and final selection. *Ann Rheum Dis* 2009;68:777–83.
- Hochberg MC. Updating the American College of rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1997;40:1725.
- Petri M, Orbai A-M, Alarcón GS, *et al.* Derivation and validation of the systemic lupus international collaborating clinics classification criteria for systemic lupus erythematosus. *Arthritis Rheum* 2012;64:2677–86.
- Jennette JC, Falk RJ, Andrasz K, *et al.* Nomenclature of systemic vasculitides. proposal of an international consensus conference. *Arthritis Rheum* 1994;37:187–92.
- Lundberg IE, Tjærnlund A, Bottai M. European League against Rheumatism/ American College of rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies and their major subgroups. *Ann Rheum Dis* 2017;2017:1955–64.



- 14 Criscuolo E, Diotti RA, Strollo M, *et al.* Weak correlation between antibody titers and neutralizing activity in sera from SARS-CoV-2 infected subjects. *J Med Virol* 2021;93:2160–7.
- 15 Furer V, Zisman D, Kibari A. Herpes zoster following BNT162b2 mRNA Covid-19 vaccination in patients with autoimmune inflammatory rheumatic diseases: a case series. *Rheumatology*.
- 16 Fischer L, Gerstel PF, Poncet A, *et al.* Pneumococcal polysaccharide vaccination in adults undergoing immunosuppressive treatment for inflammatory diseases—a longitudinal study. *Arthritis Res Ther* 2015;17:151.
- 17 Kapetanovic MC, Saxne T, Sjöholm A, *et al.* Influence of methotrexate, TNF blockers and prednisolone on antibody responses to pneumococcal polysaccharide vaccine in patients with rheumatoid arthritis. *Rheumatology* 2006;45:106–11.
- 18 Kapetanovic MC, Roseman C, Jönsson G, *et al.* Antibody response is reduced following vaccination with 7-valent conjugate pneumococcal vaccine in adult methotrexate-treated patients with established arthritis, but not those treated with tumor necrosis factor inhibitors. *Arthritis Rheum* 2011;63:3723–32.
- 19 Ribeiro ACM, Guedes LKN, Moraes JCB, *et al.* Reduced seroprotection after pandemic H1N1 influenza adjuvant-free vaccination in patients with rheumatoid arthritis: implications for clinical practice. *Ann Rheum Dis* 2011;70:2144–7.
- 20 Hua C, Barnette T, Combe B, *et al.* Effect of methotrexate, anti-tumor necrosis factor  $\alpha$ , and rituximab on the immune response to influenza and pneumococcal vaccines in patients with rheumatoid arthritis: a systematic review and meta-analysis. *Arthritis Care Res* 2014;66:1016–26.
- 21 Park JK, Lee MA, Lee EY, *et al.* Effect of methotrexate discontinuation on efficacy of seasonal influenza vaccination in patients with rheumatoid arthritis: a randomised clinical trial. *Ann Rheum Dis* 2017;76:1559–65.
- 22 Park JK, Lee YJ, Shin K, *et al.* Impact of temporary methotrexate discontinuation for 2 weeks on immunogenicity of seasonal influenza vaccination in patients with rheumatoid arthritis: a randomised clinical trial. *Ann Rheum Dis* 2018;77:898–904.
- 23 Fomin I, Caspi D, Levy V, *et al.* Vaccination against influenza in rheumatoid arthritis: the effect of disease modifying drugs, including TNF alpha blockers. *Ann Rheum Dis* 2006;65:191–4.
- 24 Elkayam O, Bashkin A, Mandelboim M, *et al.* The effect of infliximab and timing of vaccination on the humoral response to influenza vaccination in patients with rheumatoid arthritis and ankylosing spondylitis. *Semin Arthritis Rheum* 2010;39:442–7.
- 25 Kivitz AJ, Schechtman J, Texter M, *et al.* Vaccine responses in patients with rheumatoid arthritis treated with certolizumab pegol: results from a single-blind randomized phase IV trial. *J Rheumatol* 2014;41:648–57.
- 26 Rondaan C, Furer V, Heijstek MW, *et al.* Efficacy, immunogenicity and safety of vaccination in adult patients with autoimmune inflammatory rheumatic diseases: a systematic literature review for the 2019 update of EULAR recommendations. *RMD Open* 2019;5:e001035.
- 27 Mori S, Ueki Y, Hirakata N, *et al.* Impact of tocilizumab therapy on antibody response to influenza vaccine in patients with rheumatoid arthritis. *Ann Rheum Dis* 2012;71:2006–10.
- 28 Tsuru T, Terao K, Murakami M, *et al.* Immune response to influenza vaccine and pneumococcal polysaccharide vaccine under IL-6 signal inhibition therapy with tocilizumab. *Mod Rheumatol* 2014;24:511–6.
- 29 Crnkic Kapetanovic M, Saxne T, Jönsson G, *et al.* Rituximab and abatacept but not tocilizumab impair antibody response to pneumococcal conjugate vaccine in patients with rheumatoid arthritis. *Arthritis Res Ther* 2013;15:R171.
- 30 Richi P, Martin MD, de Ory F, *et al.* Secukinumab does not impair the immunogenic response to the influenza vaccine in patients. *RMD Open* 2019;5:e001018.
- 31 Furer V, Zisman D, Kaufman I, *et al.* Immunogenicity and safety of vaccination against seasonal influenza vaccine in patients with psoriatic arthritis treated with secukinumab. *Vaccine* 2020;38:847–51.
- 32 Winthrop KL, Silverfield J, Racewicz A, *et al.* The effect of tofacitinib on pneumococcal and influenza vaccine responses in rheumatoid arthritis. *Ann Rheum Dis* 2016;75:687–95.
- 33 Winthrop KL, Bingham CO, Komocsar WJ, *et al.* Evaluation of pneumococcal and tetanus vaccine responses in patients with rheumatoid arthritis receiving baricitinib: results from a long-term extension trial substudy. *Arthritis Res Ther* 2019;21:102.
- 34 Bingham CO, Looney RJ, Deodhar A, *et al.* Immunization responses in rheumatoid arthritis patients treated with rituximab: results from a controlled clinical trial. *Arthritis Rheum* 2010;62:64–74.
- 35 van Assen S, Holvast A, Benne CA, *et al.* Humoral responses after influenza vaccination are severely reduced in patients with rheumatoid arthritis treated with rituximab. *Arthritis Rheum* 2010;62:75–81.
- 36 Westra J, van Assen S, Wilting KR, *et al.* Rituximab impairs immunoglobulin (Ig)M and IgG (subclass) responses after influenza vaccination in rheumatoid arthritis patients. *Clin Exp Immunol* 2014;178:40–7.
- 37 Bar-Or A, Calkwood JC, Chognot C, *et al.* Effect of ocrelizumab on vaccine responses in patients with multiple sclerosis: the VELOCE study. *Neurology* 2020;95:e1999–2008.
- 38 Eisenberg RA, Jawad AF, Boyer J, *et al.* Rituximab-treated patients have a poor response to influenza vaccination. *J Clin Immunol* 2013;33:388–96.
- 39 Spiera J, Jinich S, Jannat-Khah D. Rituximab, but not other antirheumatic therapies, is associated with impaired serological response to SARS-CoV-2 vaccination in patients with rheumatic diseases. *Ann Rheum Dis* 2021;80:1352–3.
- 40 Grifoni A, Weiskopf D, Ramirez SI, *et al.* Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell* 2020;181:1489–501.
- 41 Jeyanathan M, Afkhami S, Smaili F, *et al.* Immunological considerations for COVID-19 vaccine strategies. *Nat Rev Immunol* 2020;20:615–32.
- 42 Alten R, Bingham CO, Cohen SB, *et al.* Antibody response to pneumococcal and influenza vaccination in patients with rheumatoid arthritis receiving abatacept. *BMC Musculoskelet Disord* 2016;17:231.
- 43 Grupper A, Rabinowich L, Schwartz D, *et al.* Reduced humoral response to mRNA SARS-CoV-2 BNT162b2 vaccine in kidney transplant recipients without prior exposure to the virus. *Am J Transplant* 2021. doi:10.1111/ajt.16615. [Epub ahead of print: 18 Apr 2021] (Published Online First: 18 April 2021).
- 44 Peled Y, Ram E, Lavee J, *et al.* BNT162b2 vaccination in heart transplant recipients: clinical experience and antibody response. *J Heart Lung Transplant* 2021. doi:10.1016/j.healun.2021.04.003. [Epub ahead of print: 21 Apr 2021].
- 45 Rodríguez-Jiménez P, Chicharro P, Cabrera L-M, *et al.* Varicella-Zoster virus reactivation after SARS-CoV-2 BNT162b2 mRNA vaccination: report of 5 cases. *JAAD Case Rep* 2021;12:58–9.
- 46 Bostan E, Yalici-Armagan B. Herpes zoster following inactivated COVID-19 vaccine: a coexistence or coincidence? *J Cosmet Dermatol* 2021;20:1566–7.
- 47 Eid E, Abdullah L, Kurban M, *et al.* Herpes zoster emergence following mRNA COVID-19 vaccine. *J Med Virol* 2021. doi:10.1002/jmv.27036. [Epub ahead of print: 29 Apr 2021] (Published Online First: 29 April 2021).