OPEN ACCESS

TRANSLATIONAL SCIENCE

Local steroid activation is a critical mediator of the anti-inflammatory actions of therapeutic glucocorticoids

Chloe Fenton,^{1,2} Claire Martin,¹ Rachel Jones,^{1,3} Adam Croft,² Joana Campos,² Amy J Naylor,^{2,4} Angela E Taylor,^{1,5} Myriam Chimen,^{2,6} Mark Cooper,⁷ Gareth G Lavery,^{1,3,5} Karim Raza ,^{2,8} Rowan S Hardy ^{2,3,6}

Handling editor Josef S Smolen

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2020-218493>).

For numbered affiliations see end of article.

Correspondence to

Dr Rowan S Hardy, IMSR, University of Birmingham, Birmingham B15 2TT, UK; R.Hardy@bham.ac.uk

Received 2 July 2020

Revised 5 October 2020

Accepted 6 October 2020

Published Online First

8 November 2020

ABSTRACT

Objectives The enzyme 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1) plays a well-characterised role in the metabolism and activation of endogenous glucocorticoids (GCs). However, despite its potent upregulation at sites of inflammation, its role in peripheral metabolism and action of therapeutic GCs remains poorly understood. We investigated the contribution of 11 β -HSD1 to the anti-inflammatory properties of the active GC corticosterone, administered at therapeutic doses in murine models of polyarthritis. **Methods** Using the tumour necrosis factor-tg and K/BxN serum-induced models of polyarthritis, we examined the anti-inflammatory properties of oral administration of corticosterone in animals with global, myeloid and mesenchymal targeted transgenic deletion of 11 β -HSD1. Disease activity and joint inflammation were scored daily. Joint destruction and measures of local and systemic inflammation were determined by histology, micro-CT, quantitative RT-PCR, fluorescence activated cell sorting and ELISA.

Results Global deletion of 11 β -HSD1 resulted in a profound GC resistance in animals receiving corticosterone, characterised by persistent synovitis, joint destruction and inflammatory leucocyte infiltration. This was partially reproduced with myeloid, but not mesenchymal 11 β -HSD1 deletion, where paracrine GC signalling between cell populations was shown to overcome targeted deletion of 11 β -HSD1.

Conclusions We identify an entirely novel component of therapeutic GC action, whereby following their systemic metabolism, they require peripheral reactivation and amplification by 11 β -HSD1 at sites of inflammation to deliver their anti-inflammatory therapeutic effects. This study provides a novel mechanistic understanding of the anti-inflammatory properties of therapeutic GCs and their targeting to sites of inflammation in polyarthritis.

INTRODUCTION

Due to their anti-inflammatory actions, therapeutic glucocorticoids (GCs) have been widely used in the management of inflammation. However, despite their continuing widespread use, several critical aspects of their therapeutic action remain unclear.¹ The enzyme 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1) plays a well characterised role in the hepatic activation of structurally inactive GCs (such as cortisone and prednisone), converting

Key messages

What is already known about this subject?

- Potent anti-inflammatory glucocorticoids such as prednisolone are rapidly metabolised, and circulate in both their active and inactive (prednisone) forms.
- Enzymes such as 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1), which is potentially upregulated at sites of inflammation, reactivates inactive glucocorticoids such as prednisone.

What does this study add?

- This study demonstrates that following their oral delivery and systemic metabolism, the anti-inflammatory properties of active glucocorticoids are completely dependent on their peripheral reactivation by 11 β -HSD type 1.
- The global deletion of 11 β -HSD type 1 results in profound therapeutic glucocorticoid resistance.

How might this impact on clinical practice or future developments?

- This study provides a novel mechanistic understanding of the anti-inflammatory properties of therapeutic glucocorticoids and their targeting to sites of inflammation.

them to their active counterparts (such as hydrocortisone and prednisolone).^{2,3} However, the role of 11 β -HSD1 in mediating the anti-inflammatory, disease-modifying actions of therapeutic GCs remains poorly understood. This represents a significant barrier to our understanding of the mechanisms of action of therapeutic GC action in vivo and to the development of GCs with an enhanced benefit:risk ratio.

We explore the contribution of pre-receptor steroid metabolism by the enzyme 11 β -HSD1 to the anti-inflammatory actions of GCs using in vivo models of chronic polyarthritis. We demonstrate a fundamental role for the peripheral re-activation of GCs in mediating their anti-inflammatory properties, with mice with global 11 β -HSD1 deletion showing a complete resistance to their therapeutic effects of orally administered GCs in their active form. These findings change our understanding of

► <http://dx.doi.org/10.1136/annrheumdis-2020-218798>

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

To cite: Fenton C, Martin C, Jones R, et al. *Ann Rheum Dis* 2021;**80**:250–260.

how many structurally active therapeutic GCs elicit their anti-inflammatory effects, requiring peripheral reactivation by the enzyme 11 β -HSD1, after their initial systemic inactivation, to mediate their beneficial immune-modulatory effects.

MATERIALS AND METHODS

Models of polyarthritis

The tumour necrosis factor (TNF)-tg model of chronic inflammatory polyarthritis, obtained courtesy of Professor George Kollias (BSRC Fleming, Athens), was maintained on a C57BL/6 background and compared with WT littermates.⁴ At day 32 of age, at the first onset of measurable polyarthritis, male TNF-tg mice received drinking water supplemented with either corticosterone (Cort) (100 μ g/mL, 0.66% ethanol), or vehicle (0.66% ethanol) for 3 weeks. Mice were scored as previously described.^{5,6} At day 53, serum was collected by cardiac puncture and tissues excised for analysis. Serum transfer-induced arthritis (STIA) was induced by intravenous injection of 100 μ L arthritogenic serum from KRN mice (K/BxN).⁷ Ankle or wrist joint thickness was monitored using callipers and reported as the change from baseline.

Targeted deletion of 11 β -HSD1

11 β -HSD1 KO animals with global 11 β -HSD1 deletion were crossed with TNF-tg animals to generate TNF-tg^{11 β KO} animals as previously described.⁸ Mesenchymal 11 β -HSD1 KO animals were generated by crossing flx/flx-HSD11B1 mice with Twist2-cre mice to generate 11 β HSD1flx/flx/Twist2cre animals, which were paired with TNF-tg animals to produce TNF-tg^{11 β HSD1flx/flx/Twist2cre} (TNF-tg^{11 β flx/tw2cre}).^{9–11} Myeloid targeted 11 β -HSD1 KO animals were generated by crossing flx/flx-HSD11B1 mice with LysM-cre mice to generate 11 β HSD1flx/flx/LysMcre animals, which were paired with TNF-tg animals to produce TNF-tg^{11 β HSD1flx/flx/LysMcre} (TNF-tg^{11 β flx/LysMcre}).¹²

11 β -HSD1 activity

11 β -HSD1 activity was determined by thin-layer chromatography as previously reported.^{8,13} Briefly, ex vivo tissue biopsies and in vitro cultures were incubated with 100 nmol/L of 11-dehydrocorticosterone (11-DHC) and tritiated [³H] tracer. Steroid conversion was measured using a Bioscan imager (Bioscan, Washington, District of Columbia, USA) and fractional conversion calculated.

Primary fibroblast-like synoviocytes and macrophage culture

Primary fibroblast-like synoviocytes (FLS) were isolated from combined hind legs and front paws from mice following dissection and cleaning of tissue as previously reported.¹³ Briefly, joints were digested in RPMI containing 2% fetal calf serum (FCS), 2.5 mg/mL collagenase D (Roche) and 20 μ g/mL DNase (Sigma-Aldrich) for 45 min at 37°C with agitation. After filtering, cells were cultured in RPMI containing 10% FCS and 1% pen-strep and cultured to passage 3 before use. Primary murine peritoneal macrophages were isolated by CD11b+ve selection with CD11b MicroBeads (Miltenyi Biotec, Surrey, UK) following peritoneal lavage in phosphate-buffered saline, and maintained in Dulbecco's Modified Eagle Medium containing 10% FCS and 1% pen-strep and maintained for up to 48 hours.

Gene expression analysis

Gene expression was assessed by TaqMan Gene Expression Assays (ThermoFisher Scientific) following mRNA isolation by innuPREP RNA Mini Kit (Analytikjena, Cambridge) and reverse transcription (Multiscribe, ThermoFisher Scientific) as

per the manufacturer's guidelines. Ccl2, cxcl2, Cxcl10, Tnf α , il1 β , Il6 and gilz were determined using species-specific probe sets by real-time PCR on an ABI7500 system (Applied Biosystems, Warrington, UK). mRNA abundance was normalised to either 18S or Gapdh. Data, obtained as Ct values and Δ Ct determined (Ct target–Ct 18S/GAPDH), were expressed as arbitrary units (AU) using the following transformation: (arbitrary units (AU)=1000 $\times$ (2^{– Δ Ct})).

ELISA analysis

Serum interleukin (IL)-6 and corticosterone (R&D Systems, Abingdon, UK) were determined using commercially available ELISA assays in accordance with the manufacturer's instructions.

Histological analysis of joints

Histochemistry was performed on paraffin-embedded 10 μ m sections. Pannus size at the humerus/ulna joint interface and osteoclast numbers on the bone surface pannus (following tartrate-resistant acid phosphatase (TRAP) staining) were determined using ImageJ software as previously reported.^{5,14} For quantification, the mean of three adjacent 10 μ m sections cut from the centre of the joint from six animals were assessed.

MicroCT morphometry analysis

Front paws from mice were imaged using a Skyscan 1172 micro-CT scanner (Bruker) using X-ray beam settings of 60 kV/167 μ A with a 0.5 mm aluminium filter. Projections were taken every 0.45° at 580 ms exposure. Image volumes were reconstructed using the Feldkamp algorithm (NRecon V1.6.1.5, Bruker) having applied beam hardening correction. Front paws were reconstructed and MeshLab V1.3.2 was used to generate meshes which could then be scored for bone erosions as described previously.⁵

Serum steroid measurements

Serum samples were collected by cardiac bleeds to assess systemic metabolism between groups; 200 μ L of serum was spiked with 0.2 ng of internal standard (corticosterone-d8 and cortisol-d4; purchased from Sigma-Aldrich, UK). Steroids were extracted via liquid-liquid extraction with 2 mL of tert-methyl butyl ether (MTBE). MTBE was evaporated to dryness under nitrogen at 55°C. Samples were reconstituted in 125 μ L of 50/50 methanol/water for liquid chromatography tandem mass spectrometry analysis.^{15,16} Samples were measured on a Waters Xevo-XS mass spectrometer coupled to an Acquity uPLC with an electrospray ionisation source in positive ionisation mode. Steroids were identified by comparison to authentic reference standards, (Sigma-Aldrich), with matching retention time and identical mass transitions and quantified relative to a calibration series. Concentrations were calculated relative to internal standard corticosterone to corticosterone-d8 and 11-DHC and its isomer metabolite to cortisol-d4.

Tissue digestion and flow cytometric analysis of synoviocytes

One hind leg and one front paw per mouse was dissected and cleaned of tissue as previously reported.⁸ Briefly, joints were digested in RPMI containing 2% FCS, 2.5 mg/mL collagenase D (Roche) and 20 μ g/mL DNase (Sigma-Aldrich) for 45 min at 37°C with agitation. After filtering, cells were centrifuged, red cells lysed and cells counted before being filtered through 40 μ m cell strainer, incubated with anti-CD16/CD32 blocking antibody (1:200; eBioscience) for 10 min at RT, followed by staining with antibody cocktail at 4°C. Antibodies for membrane

Therapy

staining are outlined in online supplemental table 1). Data were acquired using a BD LSR Fortessa X20 and analysed using FlowJo software (FlowJo). The following gating strategy was used for myeloid cells: live cells were gated on CD45+CD11b+ cells. Neutrophils identified as Ly6g+, macrophages were Ly6g-, F4/80+ and inflammatory activated M1-like macrophages were F4/80+MHC class II+. T cells were identified as live CD45+CD3+. CD3+ cells were then stratified as CD4+

or CD8+ T cells. B cells were identified as CD45+CD3 and CD19+.

Statistical analysis

Statistical significance was defined as $p < 0.05$ using either an unpaired Student's *t*-test or two-way analysis of variance with Tukey post hoc analysis where a Gaussian distribution was identified.

Figure 1 (A) Total clinical scores (arbitrary units (AU)), (B) arthritic paw scores (AU) and (C) representative images of synovitis at the ulna/humerus joint interface of wild-type (WT), tumour necrosis factor (TNF)-tg, 11βKO and TNF-tg^{11βKO} animals receiving either vehicle or corticosterone (100 µg/mL) in the drinking water for 3 weeks. (D) of TNF-tg and TNF-tg^{11βKO} animals receiving vehicle or corticosterone (100 µg/mL) in the drinking water for 3 weeks. (E) Serum levels of interleukin (IL)-6 determined by ELISA, (F) histological scoring of synovitis (AU), (G) representative images of TRAP stained osteoclasts at the ulna/humerus joint interface, (H) osteoclast number (AU) at the ulna/humerus joint interface, (I) quantification of bone erosion (AU) in the wrist, metacarpals and phalanges of WT, TNF-tg, 11βKO and TNF-tg^{11βKO} animals receiving either vehicle or corticosterone (100 µg/mL) in the drinking water for 3 weeks. (J) representative images of three-dimensional reconstructions of front paws of TNF-tg and TNF-tg^{11βKO} animals receiving vehicle or corticosterone (100 µg/mL) in the drinking water for 3 weeks, red arrows indicate erosions. (K) Total clinical scores (AU), (L) arthritic paw scores (AU) and swelling (mm) of (M) back paws and (N) ankles of WT and 11βKO animals after induction of arthritis with K/BxN serum receiving either vehicle or corticosterone (100 µg/mL) in the drinking water for 1 week. Values are expressed as mean±SE, n=6 per group for all TNF-tg experiments and n=5 (K/BxN), n=6 (K/BxN/Cort), n=6 (K/BxN^{11βKO}) and n=5 (K/BxN^{11βKO}/Cort). Statistical significance was determined using two-way analysis of variance with Tukey post hoc analysis. * $P < 0.05$, ** $p < 0.005$, *** $p < 0.001$.

RESULTS

11 β -HSD1 KO animals are resistant to therapeutic GCs

We crossed the TNF-tg murine model of chronic polyarthritis onto the 11 β KO background to generate TNF-tg animals with deletion of 11 β -HSD1 (TNF-tg^{11 β KO}). Wild-type (WT), TNF-tg, 11 β KO and TNF-tg^{11 β KO} animals received either vehicle or corticosterone in drinking water at 50 and 100 μ g/mL as previously reported.^{2 14 17} At 50 μ g/mL, no significant change in disease activity or joint inflammation were apparent in TNF-tg animals and was discontinued from the study (online supplemental figure 1A–C). At 100 μ g/mL corticosterone resulted in a significant reduction in clinical scores and joint inflammation in TNF-tg animals (figure 1A,B). In contrast, anti-inflammatory effects of corticosterone were absent in TNF-tg^{11 β KO} animals. Similarly, serum IL-6 was reduced in TNF-tg mice receiving corticosterone ($p<0.05$), which was absent in TNF-tg^{11 β KO} counterparts (figure 1D). TNF-tg mice receiving corticosterone showed a marked reduction in pannus invasion and osteoclast numbers at the pannus bone interface, which was entirely absent in TNF-tg^{11 β KO} animals (figure 1C–G). Micro-CT analysis of juxta-articular erosions confirmed that corticosterone significantly reduced joint destruction in TNF-tg mice, but not TNF-tg^{11 β KO} animals (figure 1H,I). In the KBxN serum induction model of polyarthritis, similar patterns were observed with total clinical scores and joint inflammation scores being reduced by corticosterone in WT, but not in 11 β KO animals at day 6 (figure 1J–M). Here, at early time points (days 3–4) GCs were able to partially suppress clinical scores of disease, including weight loss and lethargy in 11 β KO animals, without impacting on measures of joint inflammation and swelling. These data demonstrate that GC activation by the enzyme 11 β -HSD1 is a necessary step in mediating the anti-inflammatory actions of the GC corticosterone in the joints of animals with polyarthritis.

Oral corticosterone generates circulating 11-DHC substrate for 11 β -HSD1 activation

Metabolism and inactivation of therapeutic GCs by renal 11 β -HSD2 creates a circulating pool of inactive GC (corticosterone to 11-DHC in mice) available for peripheral activation by 11 β -HSD1. To assess the systemic metabolism of oral administered corticosterone, we measured serum levels of the

corticosterone and its inactive derivative 11-DHC. No differences were observed in daily intake of corticosterone between groups, determined by quantifying daily drinking water intake per mouse, with an average exposure of 22.5 ± 1.44 μ g/g of body weight. Here, serum corticosterone and 11-DHC were detected and significantly increased following administration of corticosterone, with exposure comparable across groups (figure 2A,B). In all groups, exposure to corticosterone resulted in a significant reduction in adrenal weights relative to vehicle (figure 2C). Analysis of corticosterone inactivation by the 11 β -HSD enzymes within the synovium was assessed by thin layer chromatography in WT, TNF-tg, 11 β KO, TNF-tg anNFtg^{11 β KO} animals and showed no significant variation between groups (online supplemental figure 2). These data confirm a comparable increase in serum corticosterone and inactive 11-DHC in TNF-tg and TNF-tg^{11 β KO} animals receiving oral corticosterone.

Effects of oral corticosterone on leucocyte recruitment are dependent on 11 β -HSD1

We examined infiltrating leucocytes and inflammatory mediators in synovial tissue digests to assess their regulation by corticosterone. Here, while corticosterone resulted in a significant decrease in total leucocytes, neutrophils, macrophages, CD8+ and CD19+, but not CD3+ and CD4+ populations in TNF-tg mice, TNF-tg^{11 β KO} animals were resistant to the actions of corticosterone on many of these parameters, with no apparent reduction in total leucocytes, macrophages and neutrophils (figure 3A–D). TNF-tg^{11 β KO} animals receiving corticosterone also possessed significantly higher numbers of total leucocytes, neutrophils, macrophages, CD3+ ($p<0.01$) and CD4+ cell populations relative to TNF-tg counterparts receiving corticosterone (figure 3A–E). Here, corticosterone skewed macrophage polarisation, with reduced numbers of inflammatory activated M1-like polarised or macrophages relative to total macrophages, TNF-tg animals, while TNF-tg^{11 β KO} animals showed complete resistance to this effect (figure 3H). However, TNF-tg^{11 β KO} animals retained an effective suppression of both CD8+ and CD19+ cell populations in response to corticosterone (figure 3F–H). Analysis of gene expression in synovial tissue digests revealed a significant reduction in the chemokines *Ccl2*, *Cxcl10* and cytokine *Il-1b*, and an increased expression of

Figure 2 (A, B) Serum corticosterone and 11-dehydrocorticosterone (11-DHC) determined by liquid chromatography tandem mass spectrometry (LCMS) from tumour necrosis factor (TNF)-tg, 11 β KO and TNF-tg11 β KO animals receiving either vehicle or corticosterone (100 μ g/mL) in the drinking water for 3 weeks. (C) Adrenal weights in wild-type (WT), TNF-tg, 11 β KO and TNF-tg11 β KO animals receiving either vehicle or corticosterone (100 μ g/mL) in the drinking water for 3 weeks. Values are expressed as mean $\pm$ SE, n=6 per group for ELISA and adrenal weights and n=3 per group for LCMS. Statistical significance was determined using two-way analysis of variance with Tukey post hoc analysis. * $P<0.05$, ** $p<0.005$, *** $p<0.001$. NS, not significant.

Figure 3 Cell numbers of (A) total leukocytes, (B) neutrophils, (C) macrophages, (D) CD3+ populations, (E) CD4+ populations, (F) CD8+ populations, (G) CD19+ populations and (H) the M1-like/total macrophage ratio determined by flow cytometry in tumour necrosis factor (TNF)-tg and TNF-tg^{11βKO} animals receiving vehicle or corticosterone (100 µg/mL) in the drinking water for 3 weeks. Gene expression (AU) of (I) *Ccl2*, (J) *Cxcl2*, (K) *Cxcl10*, (L) *Il-1*, (M) *mTnf* and (N) *Gilz* determined by quantitative PCR in tibia isolated from TNF-tg and TNF-tg^{11βKO} animals receiving vehicle or corticosterone (100 µg/mL) in the drinking water for 3 weeks. Values are expressed as mean±SE, n=6 per group. Statistical significance was determined using two-way analysis of variance with Tukey post hoc analysis. *P<0.05, **p<0.005, ***p<0.001.

anti-inflammatory *Gilz* expression in TNF-tg animals receiving corticosterone, which was entirely absent in TNF-tg^{11βKO} animals (figure 3I-N). These data reveal that TNF-tg^{11βKO} animals show marked resistance to the anti-inflammatory properties of therapeutic GCs on leukocyte recruitment and on regulation of local inflammatory mediators.

Mice with stromal deletion of 11β-HSD1 retain anti-inflammatory responses to GCs

Given the stromal upregulation of stromal 11β-HSD1 sites of inflammation, we wished to delineate its specific contribution to GC resistance in the TNF-tg^{11βflx/tw2cre} mouse relative to TNF-tg littermates, where we have previously reported effective

mesenchymal deletion.³⁸ A significant reduction in 11β-HSD1 activity was apparent in primary fibroblasts and osteoblasts isolated from TNF-tg^{11βflx/tw2cre} animals, while activity was retained in non mesenchyme derived tissues such as livers and spleen (figure 4A,B). Suppression of adrenal weights was apparent across all groups in response to corticosterone (figure 4C). Corticosterone significantly reduced clinical scores and measures of joint inflammation in both TNF-tg^{11βflx/tw2cre} and TNF-tg littermates (figure 4D,E). While circulating levels of the acute response cytokine IL-6, remained elevated in TNF-tg^{11βflx/tw2cre} receiving corticosterone, analysis of pannus invasion, osteoclast numbers and joint destruction by micro-CT, indicated that corticosterone was equally effective at suppressing disease activity

Figure 4 Corticosterone production (pmol/mg tissue/hour) in (A) fibroblast-like synoviocytes (FLS) and osteoblasts (OBs) cultures and (B) liver and spleen ex vivo biopsies isolated from tumour necrosis factor (TNF)-tg and TNF-tg ^{11βflx/tw2cre} mice determined by scanning thin layer chromatography. (C) Adrenal weights (mg), (D) total clinical scores (AU), (E) arthritic paw scores (arbitrary units (AU)), (F) serum IL-6 determined by ELISA, (G) histological scoring (AU) and (H) representative images of synovitis at the ulna/humerus joint interface, (I) histological scoring (AU) and (J) representative images of TRAP stained osteoclast numbers at the ulna/humerus joint interface, (K) quantification of bone erosion (AU) in the wrist, metacarpals and phalanges and (L) representative images of three-dimensional reconstructions of front paws in TNF-tg and TNF-tg ^{11βflx/tw2cre} animals receiving either vehicle or corticosterone (100 µg/mL) in the drinking water for 3 weeks. Values are expressed as mean±SE, n=6 per group. Statistical significance was determined using two-way analysis of variance with Tukey post hoc analysis. *P<0.05, **p<0.005, ***p<0.001.

in both TNF-tg and TNF-tg ^{11βflx/tw2cre} animals (figure 4C,G-L). Consequently, despite effective deletion of 11β-HSD1 in the mesenchymal compartment, TNF-tg ^{11βflx/tw2cre} animals retain a robust anti-inflammatory response to corticosterone.

Partial GCs resistance with myeloid deletion of 11β-HSD1

At sites of inflammation, 11β-HSD1 is highly expressed in macrophages and is implicated in regulating their anti-inflammatory properties.^{3 18 19} We used the LysMCre mouse (targeted towards neutrophils, macrophages and granulocytes) to generate tg ^{11βflx/LysMCre} animals with a deletion of 11β-HSD1 in the myeloid compartment.¹² 11β-HSD1 activity was significantly reduced in both peripheral blood mononuclear cell and peritoneal macrophages relative to WT counterparts (figure 5A). In contrast, normal 11β-HSD1 activity was apparent in tissues such as muscle, fat and liver (figure 5A,B). Corticosterone significantly

reduced adrenal weights in both TNF-tg ^{11βflx/LysMCre} animals and TNF-tg littermate controls of (figure 5C). TNF-tg ^{11βflx/LysMCre} mice receiving corticosterone showed a significant reduction in joint inflammation scores but not in total clinical scores, while serum IL-6 levels were similarly decreased in both TNF-tg and TNF-tg ^{11βflx/LysMCre} (figure 4D-F). A, significant reductions in both pannus size and osteoclast numbers were apparent in both TNF-tg and TNF-tg ^{11βflx/LysMCre} animals receiving corticosterone (figure 5G-J). However, evidence of residual pannus and osteoclast numbers in TNF-tg ^{11βflx/LysMCre} animals was supported by a greater incidence of juxta-articular joint destruction determined by micro-CT relative to TNF-tg counterparts (p<0.05) (figure 5K,L). These data demonstrate that mice with a myeloid targeted deletion of 11β-HSD1 retain the capacity to respond to therapeutic GCs.

Figure 5 Corticosterone production (pmol/mg tissue/hour) in (A) tibialis anterior (TA) muscle, fat and liver ex vivo biopsies and (B) monocyte, peripheral blood mononuclear cell (PBMC)-derived macrophages and resident macrophages isolated from tumour necrosis factor (TNF)-tg and TNF-tg^{11βHSD1^{LysMCre}} mice determined by scanning thin layer chromatography. (C) Adrenal weights (mg), (D) total clinical scores (arbitrary units (AU)), (E) arthritic paw scores (AU), (F) serum interleukin (IL)-6 determined by ELISA, (G) histological scoring (AU) and (H) representative images of synovitis at the ulna/humerus joint interface, (I) histological scoring (AU) and (J) representative images of TRAP stained osteoclast numbers at the ulna/humerus joint interface, (K) quantification of bone erosion (AU) in the wrist, metacarpals and phalanges and (L) representative images of three-dimensional reconstructions of front paws in TNF-tg and TNF-tg^{11βHSD1^{LysMCre}} animals receiving either vehicle or corticosterone (100 µg/mL) in the drinking water for 3 weeks. Values are expressed as mean±SE, n=6 per group. Statistical significance was determined using two-way analysis of variance with Tukey post hoc analysis. *P<0.05, **p<0.005, ***p<0.001.

Paracrine GC signalling compensates for cell-specific 11β-HSD1 deletion

Given our findings in the stromal and myeloid targeted models, we performed co-culture experiments in FLS and macrophage to determine if GCs activated in one cell population could influence the other by paracrine signalling. We generated conditioned media by exposing WT and 11β-HSD1 KO FLS to the inactive GC 11-DHC for 24 hours, which was then placed on 11β-HSD1 KO macrophages for a further 24 hours prior to measuring GC response genes (figure 6A). Here, 11β-HSD1 KO macrophages responded to conditioned media from WT FLS exposed to 11-DHC (increasing *Gilz* and suppressing *IL-6*), but not conditioned media from 11β-HSD1 KO FLS (figure 6B,C). Conditioned media from WT and 11β-HSD1 KO macrophages conditioned with 11-DHC were then placed on 11β-HSD1 KO FLS for 24 hours and GC responsive gene analysed (figure 6D). 11β-HSD1 KO FLS responded to conditioned media from WT macrophages exposed to 11-DHC (increasing *Gilz* and

suppressing *IL-6*), but did not respond to conditioned media generated in 11β-HSD1 KO macrophages (figure 6E,F). Similarly, *IL-6* production in 11β-HSD1 KO FLS was suppressed in response to media from WT macrophages exposed to 11-DHC, but not from 11β-HSD1 KO macrophages exposed to 11-DHC (figure 6G). These data confirm that 11β-HSD1 can mediate paracrine GC signalling between distinct cell populations present at site of inflammation, including macrophages and FLS.

DISCUSSION

Despite the potent upregulation of 11β-HSD1 at sites of inflammation, its roles in mediating the effects of active therapeutic GCs have remained poorly understood.^{19–22} Here, studies by Schmidt *et al* and Hardy *et al* reported increasing levels of 11β-HSD1 within FLS and synovial macrophages that correlated with inflammation. Using murine models of polyarthritis, we have identified an entirely novel and, until now, unrecognised

Figure 6 (A) Schematic representation of conditioned media experiments in which media from wild-type (WT) and 11 β KO fibroblast-like synoviocytes (FLS) treated with either vehicle or dehydrocorticosterone (DHC) is used to treat 11 β KO macrophages. Gene expression (arbitrary units (AU)) of (B) *Gilz* and (C) *Il-6* in 11 β KO macrophages treated with conditioned media from WT and 11 β KO FLS determined by quantitative PCR (qPCR). (D) Schematic representation of conditioned media experiments in which media from WT and 11 β KO macrophages treated with either vehicle or DHC is used to treat 11 β KO FLS. Gene expression (AU) of (E) *Gilz* and (F) *Il-6* in 11 β KO FLS treated with conditioned media from WT and 11 β KO macrophages determined by qPCR. (G) Protein levels of IL-6 (ng/mL) in the media of 11 β KO FLS treated with conditioned media from WT and 11 β KO macrophages determined by ELISA. Values are expressed as mean $\pm$ SE, n=3 per group. Statistical significance was determined using one-way analysis of variance with Tukey post hoc analysis. *P<0.05, **p<0.005, ***p<0.001.

component of therapeutic GC action, whereby they require peripheral reactivation by 11 β -HSD1 at sites of inflammation to deliver anti-inflammatory effects (figure 7). Here, the global transgenic deletion of 11 β -HSD1 prevents this critical step, resulting in severe GC resistance in both TNF-tg and K/BxN models of polyarthritis.

The importance of endogenous GC metabolism by 11 β -HSD1 in the pathophysiology of inflammatory polyarthritis are well established.^{8 18 23} We used oral corticosterone to suppress disease activity and joint inflammation in murine models of polyarthritis.^{2 14 24} We observed effective suppression at 100 μ g/mL in the drinking water, where daily intake of corticosterone

was 22.5 μ g/g in the mice, and would be estimated to equate to administration of 40mg hydrocortisone or 10mg of prednisolone per day in an adult. Here, total active corticosterone and inactive 11-DHC, increased in TNF-tg and TNF-tg^{11 β KO} mice, with the reduced transcortin binding affinity of 11-DC predicted to further elevate its circulating free levels.²⁵ In the models of polyarthritis, disease activity, synovitis and joint destruction were markedly suppressed in animals receiving corticosterone. However, we observed a profound GC resistance in 11 β -HSD1 KO animals, despite equivalent serum exposure to corticosterone and 11-DHC. While these animals with deletion of 11 β -HSD1 retained a capacity to respond to

therapeutic GC in the inflamed synovium is the suppression of leucocyte recruitment and reduction in pro-inflammatory cytokines and chemokines through suppression of pro-inflammatory pathways.^{26–28} In this feature of anti-inflammatory GC action, there remains ongoing debate in relation to the relative contributions of transactivation and transrepression as mechanism underlying the anti-inflammatory effects of GCs.²⁹ In this study, we were unable to assess whether the anti-inflammatory properties of GC metabolism by 11 β -HSD1 were predominantly mediated by transrepression or transactivation, which remains a prominent area of interest and the focus of prominent reviews in the field.¹ In TNF-tg^{11 β KO} animals, expression of pro-inflammatory mediators persisted at sites of inflammation in response to corticosterone, with increased inflammatory activated M1-like polarisation, revealing a critical role for therapeutic GC metabolism by 11 β -HSD1 in this process. These data suggest that in response to therapeutic GCs, 11 β -HSD1 may mediate a shift from inflammatory activated M1-like macrophages to M2-like polarisation. However, the precise nature of these changes in this setting remains complex and is the subject, requiring more detailed characterisation, which in itself is the feature of

a notable systematic review by Tardito *et al.*³⁰ Of interest, we observed that several leucocyte population, including CD8 T cells and CD19 B cells within the synovium of 11 β -HSD1 KO animals retained responsiveness to oral corticosterone and were suppressed to a similar degree as WT counterparts. These data indicate that certain leucocyte populations retain the capacity to respond to circulating levels of active corticosterone, present even in the absence of 11 β -HSD1, suggesting they possess a lower GC receptor activation threshold that is independent of 11 β -HSD1.

Given that FLS and macrophages highly express 11 β -HSD1 in inflammatory environments, we examined whether targeted deletion of 11 β -HSD1 within mesenchymal derived FLS and myeloid-derived macrophages could recapitulate global steroid resistance. Here, despite effective 11 β -HSD1 deletion in FLS, TNF-tg^{11 β flx/tw2cre} animals showed an entirely normal anti-inflammatory response to oral corticosterone, suggesting that GC reactivation within this subset alone was not critical to the anti-inflammatory properties of corticosterone. Similar findings were evident in myeloid-targeted TNF-tg^{11 β flx/LysMcre}. Here despite effective deletion of 11 β -HSD1 within macrophages, TNF-tg^{11 β flx/LysMcre} animals retained the capacity to respond to corticosterone. However, their response did appear to be muted, with disease activity scores being greater than TNF-tg counterparts, and with evidence of persistent joint destruction despite exposure to therapeutic GCs. The contribution of 11 β -HSD1 within further leucocyte populations such as T cells to corticosterone resistance in the global KO deserve further scrutiny in this context but went beyond the scope of this study.

However, these data may suggest that the autocrine amplification of GCs by 11 β -HSD1 within fibroblasts or macrophages alone is insufficient to mediate the anti-inflammatory actions of therapeutic GCs in vivo, as occurs with cell-targeted GC receptor KO studies.^{31–33} Instead, we explored whether paracrine signalling between macrophages and FLS might overcome cell-targeted deletion of 11 β -HSD1 using in vitro models. These experiments revealed that metabolism of 11-DHC by 11 β -HSD1 could mediate paracrine GC signalling between FLS and macrophages, compensating for cell-targeted deletion of 11 β -HSD1 and reversing steroid resistance. Consequently, it may be that targeting any one cell population is insufficient to reproduce the phenotype observed with global 11 β -HSD1 deletion. Ultimately, it is important to note that results observed in animal models are not always replicated in human disease, and these findings now require robust validation in patients with inflammatory disease.

In this study, we demonstrate a profound and previously unrecognised role for pre-receptor metabolism and activation of GCs by the enzyme 11 β -HSD1 in mediating the anti-inflammatory therapeutic actions of oral GCs. Consequently, this study adds significant insight into our mechanistic understanding of therapeutic GC action. Here, a greater awareness of the how 11 β -HSD1 targets the anti-inflammatory actions of therapeutic GCs at sites of inflammation may be able to inform the development of better-tolerated steroids that possess enhanced kinetics and activation efficiency by 11 β -HSD1 to improve targeting and dosing efficacy, as well as informing ongoing studies examining the application of therapeutic 11 β -HSD1 inhibitors.

Author affiliations

¹Institute for Metabolism and Systems Research, University of Birmingham, Birmingham, UK

²Research into Inflammatory Arthritis Centre, Versus Arthritis, Institute of Inflammation and Ageing, University of Birmingham, Birmingham, UK

³MRC Arthritis Research UK Centre for Musculoskeletal Ageing Research, University of Birmingham Edgbaston Campus, Birmingham, UK

⁴Centre for Translational Inflammation Research, University of Birmingham, Birmingham, UK

⁵Centre for Endocrinology, Diabetes and Metabolism, University of Birmingham, Birmingham, UK

⁶Institute for Clinical Sciences, University of Birmingham, Birmingham, UK

⁷ANZAC Research Institute, The University of Sydney, Sydney, New South Wales, Australia

⁸Rheumatology, Sandwell and West Birmingham Hospitals NHS Trust, Birmingham, UK

Correction notice This article has been corrected since it published Online First. Figures 3 and 4 have been corrected.

Twitter Joana Campos @joanadcampos

Acknowledgements The authors would like to thank the Versus Arthritis funded Research into Inflammatory Arthritis Centre and the Medical Research Council and Versus Arthritis Centre of Excellence in Musculoskeletal (MSK) Ageing Research. The authors would like to thank Professor George Kollias (BSRC Fleming, Athens) for access to the TNF-tg model of polyarthritis, the Biomedical Services Unit (University of Birmingham) for supporting animal experiments and the Department of Musculoskeletal Pathology (Robert Aitken Institute, University of Birmingham) for embedding and cutting tissue for histology.

Contributors CF carried out experiments with support from CM, RJ, AC, AJN, AET, MC, JC and RH. RH, KR, GGL and MC conceived and supervised the project. RH wrote the manuscript. All authors discussed the results and contributed to the final manuscript.

Funding This research was supported by Versus Arthritis grants (reference: 19859 and 20843).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Procedures were performed under guidelines by the Animal (Scientific Procedures) Act 1986 in accordance with the project licence (P51102987) and approved by the Birmingham Ethical Review Subcommittee (BERSC).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. All data relevant to the study are included in the article or uploaded as online supplementary information. Data are available on reasonable request from Dr Hardy. Reuse of data is not permitted by a third party without authorisation.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Karim Raza <http://orcid.org/0000-0003-1570-1232>

Rowan S Hardy <http://orcid.org/0000-0001-6938-6739>

REFERENCES

- Hardy RS, Raza K, Cooper MS. Therapeutic glucocorticoids: mechanisms of actions in rheumatic diseases. *Nat Rev Rheumatol* 2020;16:133–44.
- Fenton CG, Webster JM, Martin CS, *et al.* Therapeutic glucocorticoids prevent bone loss but drive muscle wasting when administered in chronic polyarthritis. *Arthritis Res Ther* 2019;21:182.
- Hardy R, Rabbitt EH, Filer A, *et al.* Local and systemic glucocorticoid metabolism in inflammatory arthritis. *Ann Rheum Dis* 2008;67:1204–10.
- Keffer J, Probert L, Cazaris H, *et al.* Transgenic mice expressing human tumour necrosis factor: a predictive genetic model of arthritis. *Embo J* 1991;10:4025–31.

- 5 Naylor AJ, Desanti G, Saghir AN, *et al.* TNF α depleting therapy improves fertility and animal welfare in TNF α -driven transgenic models of polyarthritis when administered in their routine breeding. *Lab Anim* 2018;52:59–68.
- 6 Hardy RS, Doig CL, Hussain Z, *et al.* 11 β -Hydroxysteroid dehydrogenase type 1 within muscle protects against the adverse effects of local inflammation. *J Pathol* 2016;240:472–83.
- 7 Kollias G, Papadaki P, Apparailly F, *et al.* Animal models for arthritis: innovative tools for prevention and treatment. *Ann Rheum Dis* 2011;70:1357–62.
- 8 Hardy RS, Fenton C, Croft AP, *et al.* 11 beta-hydroxysteroid dehydrogenase type 1 regulates synovitis, joint destruction, and systemic bone loss in chronic polyarthritis. *J Autoimmun* 2018;92:104–13.
- 9 Yu K, Xu J, Liu Z, *et al.* Conditional inactivation of FGF receptor 2 reveals an essential role for FGF signaling in the regulation of osteoblast function and bone growth. *Development* 2003;130:3063–74.
- 10 Li A, Hardy R, Stoner S, *et al.* Deletion of mesenchymal glucocorticoid receptor attenuates embryonic lung development and abdominal wall closure. *PLoS One* 2013;8:e63578.
- 11 Semjonous NM, Sherlock M, Jeyasuria P, *et al.* Hexose-6-Phosphate dehydrogenase contributes to skeletal muscle homeostasis independent of 11 β -hydroxysteroid dehydrogenase type 1. *Endocrinology* 2011;152:93–102.
- 12 Abram CL, Roberge GL, Hu Y, *et al.* Comparative analysis of the efficiency and specificity of myeloid-Cre deleting strains using ROSA-EYFP reporter mice. *J Immunol Methods* 2014;408:89–100.
- 13 Hardy RS, Hülso C, Liu Y, *et al.* Characterisation of fibroblast-like synoviocytes from a murine model of joint inflammation. *Arthritis Res Ther* 2013;15:R24.
- 14 Fenton CG, Doig CL, Fareed S, *et al.* 11 β -HSD1 plays a critical role in trabecular bone loss associated with systemic glucocorticoid therapy. *Arthritis Res Ther* 2019;21:188.
- 15 Haring R, Wallaschofski H, Teumer A, *et al.* A SULT2A1 genetic variant identified by GWAS as associated with low serum DHEAS does not impact on the actual DHEA/DHEAS ratio. *J Mol Endocrinol* 2013;50:73–7.
- 16 O'Reilly MW, Westgate CS, Hornby C, *et al.* A unique androgen excess signature in idiopathic intracranial hypertension is linked to cerebrospinal fluid dynamics. *JCI Insight* 2019;4. doi:10.1172/jci.insight.125348. [Epub ahead of print: 21 Mar 2019].
- 17 Morgan SA, McCabe EL, Gathercole LL, *et al.* 11 β -HSD1 is the major regulator of the tissue-specific effects of circulating glucocorticoid excess. *Proc Natl Acad Sci U S A* 2014;111:E2482–91.
- 18 Coutinho AE, Gray M, Brownstein DG, *et al.* 11 β -Hydroxysteroid dehydrogenase type 1, but not type 2, deficiency worsens acute inflammation and experimental arthritis in mice. *Endocrinology* 2012;153:234–40.
- 19 Gilmour JS, Coutinho AE, Cailhier J-F, *et al.* Local amplification of glucocorticoids by 11 beta-hydroxysteroid dehydrogenase type 1 promotes macrophage phagocytosis of apoptotic leukocytes. *J Immunol* 2006;176:7605–11.
- 20 Schmidt M, Weidler C, Naumann H, *et al.* Reduced capacity for the reactivation of glucocorticoids in rheumatoid arthritis synovial cells: possible role of the sympathetic nervous system? *Arthritis Rheum* 2005;52:1711–20.
- 21 Hardy R, Rabbitt EH, Filer A, *et al.* Local and systemic glucocorticoid metabolism in inflammatory arthritis. *Ann Rheum Dis* 2008;67:1204–10.
- 22 Stegk JP, Ebert B, Martin H-J, *et al.* Expression profiles of human 11beta-hydroxysteroid dehydrogenases type 1 and type 2 in inflammatory bowel diseases. *Mol Cell Endocrinol* 2009;301:104–8.
- 23 Hardy RS, Doig CL, Hussain Z, *et al.* 11 β -Hydroxysteroid dehydrogenase type 1 within muscle protects against the adverse effects of local inflammation. *J Pathol* 2016;240:472–83.
- 24 Gasparini SJ, Weber M-C, Henneicke H, *et al.* Continuous corticosterone delivery via the drinking water or pellet implantation: a comparative study in mice. *Steroids* 2016;116:76–82.
- 25 Dunn JF, Nisula BC, Rodbard D. Transport of steroid hormones: binding of 21 endogenous steroids to both testosterone-binding globulin and corticosteroid-binding globulin in human plasma. *J Clin Endocrinol Metab* 1981;53:58–68.
- 26 King EM, Chivers JE, Rider CF, *et al.* Glucocorticoid repression of inflammatory gene expression shows differential responsiveness by transactivation- and transrepression-dependent mechanisms. *PLoS One* 2013;8:e53936.
- 27 Miyata M, Lee J-Y, Susuki-Miyata S, *et al.* Glucocorticoids suppress inflammation via the upregulation of negative regulator IRAK-M. *Nat Commun* 2015;6:6062.
- 28 Eddleston J, Herschbach J, Wagelie-Steffen AL, *et al.* The anti-inflammatory effect of glucocorticoids is mediated by glucocorticoid-induced leucine zipper in epithelial cells. *J Allergy Clin Immunol* 2007;119:115–22.
- 29 Hua G, Zein N, Paulen L, *et al.* The glucocorticoid receptor agonistic modulators CpdX and CpdX-D3 do not generate the debilitating effects of synthetic glucocorticoids. *Proc Natl Acad Sci U S A* 2019;116:14200–9.
- 30 Tardito S, Martinelli G, Soldano S, *et al.* Macrophage M1/M2 polarization and rheumatoid arthritis: a systematic review. *Autoimmun Rev* 2019;18:102397.
- 31 Koenen M, Culemann S, Vettorazzi S, *et al.* Glucocorticoid receptor in stromal cells is essential for glucocorticoid-mediated suppression of inflammation in arthritis. *Ann Rheum Dis* 2018;77:1610–8.
- 32 Baschant U, Frappart L, Rauchhaus U, *et al.* Glucocorticoid therapy of antigen-induced arthritis depends on the dimerized glucocorticoid receptor in T cells. *Proc Natl Acad Sci U S A* 2011;108:19317–22.
- 33 Tuckermann JP, Kleiman A, Moriggl R, *et al.* Macrophages and neutrophils are the targets for immune suppression by glucocorticoids in contact allergy. *J Clin Invest* 2007;117:1381–90.