Functional genomics of autoimmune diseases

Akari Suzuki, Matteo Maurizio Guerrini, Kazuhiko Yamamoto

Handling editor Josef S Smolen

Laboratory for Autoimmune Diseases, RIKEN Center for Integrative Medical Sciences, Yokohama, Kanagawa, Japan

Correspondence to

Dr Kazuhiko Yamamoto, Laboratory for Autoimmune Diseases, RIKEN Center for Integrative Medical Sciences, Yokohama 230-0045, Kanagawa, Japan; kazuhiko.yamamoto@riken.jp

Received 30 October 2020
Accepted 6 December 2020
Published Online First
6 January 2021

ABSTRACT

For more than a decade, genome-wide association studies have been applied to autoimmune diseases and have expanded our understanding on the pathogenesis. Genetic risk factors associated with diseases and traits are essentially causative. However, elucidation of the biological mechanism of disease from genetic factors is challenging. In fact, it is difficult to identify the causal variant among multiple variants located on the same haplotype or linkage disequilibrium block and thus the responsible biological genes remain elusive. Recently, multiple studies have revealed that the majority of risk variants locate in the non-coding region of the genome and they are the most likely to regulate gene expression such as quantitative trait loci. Enhancer, promoter and long non-coding RNA appear to be the main target mechanisms of the risk variants. In this review, we discuss functional genetics to challenge these puzzles.

INTRODUCTION

Autoimmune diseases, such as rheumatoid arthritis, are thought to arise from the interaction of genetic and environmental factors. The contribution of genetic factors to autoimmune diseases has been examined for more than 40 years. Initial investigations focused on major histocompatibility complex genes and were followed by those that focused on several other candidate genes. Improvements in commercial arrays since 2007 has led to the development of the current form of genome-wide association studies (GWASs). GWASs have been applied in many studies investigating autoimmune diseases and have expanded our understanding of their underlying genetic factors. GWASs rely on single-nucleotide polymorphisms (SNPs) and haplotype blocks, which are distributed across all chromosomes. A number of representative tag SNPs—500 000 to several millions—are selected based on the haplotype information and combined into a single array for typing. The loci associated with disease susceptibility can then be determined by comparing variants represented by tag SNPs between patients and controls and identifying those that are significantly different.

GWASs have helped identify thousands of loci as being statistically associated with a risk for various diseases and traits.¹ The same loci have also been identified in other independent analyses of the same diseases, suggesting that the associations determined using GWASs hold true.² Genetic risk factors are essentially causative for specific diseases and traits; however, understanding the biological mechanism of risk from a genetic factor is challenging. Although there are only three possible patterns, with corresponding intermediate phenotypes, in which risk variants are involved in the development of a disease (figure 1), the types of change brought about by risk variants are complex. Furthermore,

many variants significantly associated with each other are often in strong linkage disequilibrium (LD) and form a haplotype.³ Thus, although associations between variants and diseases can be identified, it is difficult to identify the causal variant among multiple variants located on the same haplotype,⁴ and experimental validations are needed to determine the functional variants.⁵ Furthermore, understanding the biological functions of risk variants is challenging, since more than 80% of disease-associated variants are located in non-coding regions of the genome.⁶ Transcriptome analysis can offer some clues regarding how these variants alter gene regulation and the expression of splice variants of target genes,⁷ but the detailed relationship between risk variants and genes is largely unknown.

Analyses that integrate various ‘omics’ studies and GWAS data have emerged as powerful tools for understanding the functions of risk variants identified in a GWAS. The capabilities of next-generation sequencing techniques for analysing the functions of non-coding regions have advanced dramatically in recent years, enabling the comprehensive analysis of enhancers, promoters, histone modifications and chromatin structures. Analysis of the expression of quantitative trait loci (QTL) is used to investigate how particular variants lead to intermediate phenotypes (eg, epigenetic statuses). QTL are DNA markers on a chromosome that indicate genes involved in a quantitative trait.⁸ Recently, QTL analysis has been used to identify loci that are associated with a particular quantitative phenotypic trait or disease and can be caused to polygenic effects.⁹ Therefore, QTL analysis has been used for assessing human polygenic diseases and phenotypes as well as intermediate phenotypes, such as gene expression.

In this review, we discuss the possible functional genomic strategies for integrating GWAS results with the current understanding of specific diseases. In particular, we focus on promoters, enhancers and long non-coding RNAs. We also list a suite of analytical methods for the functional dissection of disease-associated risk loci. We believe that functional genomics will enhance our understanding of the immunological and biological functions of genes.

FUNCTIONS OF GWAS-IDENTIFIED RISK VARIANTS

The ever-increasing number of genome-wide genotyping arrays has enabled the imputation of variants and the compilation of larger and more accurate reference datasets.^{10–12} Identifying the function of risk variants is important for understanding the mechanism of disease onset. Indeed, several risk variants affect the structures of proteins, for example, by inducing amino acid alterations or

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Suzuki A, Guerrini MM, Yamamoto K. *Ann Rheum Dis* 2021;**80**:689–697.

Figure 1 Risk patterns of causative variants and intermediate phenotypes. Disease development involves a variety of characteristic intracellular and internal events. These are collectively referred to as intermediate phenotypes when we consider a causal genotype and the disease outcome. Intermediate phenotypes involve gene expression, protein expression and epigenetic effects and others. There are three possible patterns via which causative variants are associated with disease. First, a risk-associated genotype acts on the disease via the intermediate phenotype, which is assigned as the causal intermediate phenotype. Second, the intermediate phenotype is not causal but the consequence of the disease. Third, intermediate phenotypes are independent from the disease.

splicing variations. However, the functions of the majority (more than 80%) of risk variants are unknown because many of them are located in non-coding genomic regions. To elucidate the functions of risk variants, more detailed GWAS and omics data regarding intermediate phenotypes of the whole genome are necessary.

Post-GWAS omics analyses can improve our understanding of the molecular events orchestrating the development of diseases and traits. In particular, high-throughput transcriptomic^{3,7} and genome-wide epigenomic³ analyses are commonly used omics analyses that can be exploited to identify QTL. QTL identified through omics analyses overlapping with GWAS-identified risk variants hint at the pathogenetic mechanism of a disease. There are various types of QTL, which we have described below. The most common QTL type is expression QTL (eQTL), which is defined as a variant directly related to the expression level.

Expression QTL

The majority of disease-associated variants reside in non-coding regions of the genome, suggesting that they regulate gene expression. Indeed, a combination of statistical analysis, imputed genotypes in LD, transcriptomics and epigenomics has helped identify a number of eQTL that overlap with variants implicated in autoimmune diseases.³ Since eQTL are strongly influenced by epigenetic changes, they are often observed in conjunction with epigenetic QTL (figure 2). In many cases, eQTL function as cis-eQTL, which directly affect the expression of nearby genes (usually within 250 kb to 1 Mb) in an allele-specific manner.^{7,13,14} In contrast, eQTL that are located at distance from their gene-of-origin, often on different chromosomes, are referred to as trans-eQTL. Some cis-eQTL associated to house-keeping genes, are observed in many different tissue types; however, the majority of eQTL are highly tissue specific.¹² The 3D map of chromatin interaction can be used to contextualise eQTLs. Hi-C and other chromosome conformation capture technologies provide a quantitative and qualitative description of the 3D arrangement of the genome and further our understanding of gene regulation.¹⁵

Splicing QTL

Splicing QTL (sQTL), a subtype of eQTL, enable the quantitation of genetic variation due to the expression of RNA isoforms from alternative splicing events. sQTL are thought to contribute significantly to disease development because splicing is a major post-transcriptional modification and the primary mechanism

governing protein diversity. Alternative splicing affects multiexonic genes, often in a cell type-specific fashion, and modulates phenotypic changes in a flexible and dynamic manner.¹⁶ RNA-sequencing (RNA-seq) is used for analysing both eQTL and sQTL.¹⁶ In multiple sclerosis, the estimated effect size of sQTL is larger than that of eQTL, supporting the well-known importance of RNA splicing in the brain.³ Thus, RNA splicing is a promising area of investigation to understand the effect of genetic variation in a complex disease.

Epigenetic QTL, including DNA methylation QTL and histone modification QTL

Epigenetic events are classically defined as post-translational modifications that determine chromatin status, including DNA methylation and multiple types of histone modifications (eg, H3K4me1, H3K4me3 and H3K27ac; figure 2). DNA methylation QTL (meQTL) and histone modification QTL are directly affected by their different genotypes. Epigenetic QTL are cell type and context specific.¹⁷ Indeed, epigenetic QTLs are highly specific for disease conditions, such as malignancy, or the effects of specific drugs. It is tempting to speculate that the high degree of specificity inherent to epigenetic QTL can be exploited to identify novel patient-specific therapeutic targets.

Assay for transposase-accessible chromatin QTL and DNase I hypersensitive site QTL

Gene expression and epigenetic events involve chromatin status, which is affected by both genetic variants and environmental factors. DNase I hypersensitive sites (DHS) indicate an open chromatin structure influence and are influenced by transcriptional regulatory elements and histone modifications (figure 2).⁹ Assay for transposase-accessible chromatin sequencing (ATAC-seq) also detects open chromatin. Genetic variants that are correlated with the phenotypes of DHS and ATAC are called DHS-QTL and ATAC-QTL, respectively. These QTL contribute to differences in gene regulation between individuals.¹⁸

Transcription factor-binding site QTL

Sequence-specific transcription factors (TFs) regulate gene expression by binding to cis-regulatory elements in the promoter and enhancer regions. TFs typically recognise binding-site motifs with particular DNA sequences (figure 2). Variants in DNA TF binding sites are designated as TF-binding site QTL (bQTL). For

Figure 2 Genetic variants and quantitative trait loci (QTL). (A) Genetic variants can affect different types of transcriptional and post-transcriptional regulatory factors, including (a) gene expression level, (b) open chromatin status, (c) histone modifications, (d) promoter usage and enhancer activity, (e) chromatin conformation, (f) transcription factor binding sites, and (g) DNA methylation. (B) Actually, genome-wide association studies (GWAS) results integrate a variety of omics layer data constructed using the latest technologies (eg, the eQTL study involves RNA-seq for exploring gene expression profiles). In the upper row, the regional association plot of GWAS is presented. The purple dot indicates the top single-nucleotide polymorphism (SNP) (SNP with the smallest p value). SNPs highly correlated with the top SNP are equally likely to be causal in the LD block (pink box), demonstrating how omics data can be combined with GWAS findings to provide insights into causal variants and genes. These SNPs are usually outside the gene body (pink peaks, a) and are located in the regulatory region of the gene. The region is open (assay for transposase-accessible chromatin (ATAC)-peaks (blue), b) and flanked by modified histones (light blue and pink peaks, c). The non-coding RNA-expressed region (d) indicates active promoter and enhancer positions, which interact (Micro-C, brown bars, e) with each other. Integration of these data for each cell provides evidence for genes and cell types implicated in the disease GWAS findings. CAGE, cap analysis of gene expression; ChIP, chromatin immunoprecipitation.

example, the binding site of transcription factor 21 has been reported to have bQTL effects.¹⁹

Promoter usage QTL and enhancer activity QTL

eQTL variants are typically enriched in regulatory regions. In particular, variants located in promoter regions are associated with alternative transcript usage and include variations in the mRNA 5'-end. These are termed promoter usage QTL (puQTL; figure 2).¹⁸ On average, there are more than four promoters per human gene, indicating that studies of puQTL are important for understanding eQTL effects.¹⁸

Enhancers are located remotely from promoters, and, unlike promoters, they regulate gene expression in a

distance-independent manner by impacting transcriptional timing and cell-specificity of gene expression. Variants related to enhancer activity are referred to as enhancer activity QTL (eaQTL; figure 2). Methods that capture the 5'-end of a transcript are often used to identify puQTL and eaQTL.²⁰

Non-coding RNA QTL

Micro RNAs (miRNAs) and long non-coding RNAs (lncRNAs) represent a class of non-coding RNAs that act as post-transcriptional regulators of gene expression and mRNA translation.²¹ miRNAs and lncRNAs are transcripts composed of 21–25 bp and >200 bp, respectively, and are extensively reported to be involved in transcriptional regulation (figure 1).

We describe the relationship between promoters, enhancers and long non-coding RNAs in detail below.

PROMOTERS AND RISK VARIANTS

Promoters are regulatory modules located upstream of the transcription start site (TSS). They are approximately 100–1000bp long and contain a core region and one or more proximal regions.²² The core promoter supports the assembly of the transcription preinitiation complex, including general TFs and RNA polymerase (Pol) II.²³ Core promoters control the magnitude of transcription, but not its induction, by regulating the binding of RNA Pol II at the TSS. Once the RNA Pol II reaches a few bases downstream of the TSS, further transcript elongation requires the binding of TFs and cofactors to the proximal region of the promoter.^{24–26}

Genetic variants located in promoter regions, originally designated as eQTL, influence the strength and/or stability of the interaction between TFs and the promoter of an immune system-related gene, likely modifying its expression pattern and thus facilitating the onset of autoimmune diseases.²⁷ For example, a variant altering the binding affinity for the transcription factor C/EBP β in the promoter of *FasL* increases the risk of developing systemic lupus erythematosus.²⁸ Similarly, an eQTL variant in the promoter of *FCRL3*, highly expressed in B cells, increases its affinity for NF- κ B binding, rendering it a risk factor for rheumatoid arthritis.²⁹

Promoters are widespread; thus, promoter risk variants that alter gene expression cannot always be determined by eQTL analysis but are revealed by puQTL analysis. Cap analysis of gene expression (CAGE), which detects the 5'-end of target transcripts, can help detect puQTL and eaQTL.^{20 30} In some cases, the results of CAGE are not linked to eQTL results. For example, *TTC23* has two promoter regions 1.6kb apart; rs8028374 was mapped as a puQTL with significant opposite effects on the two promoter regions. However, *TTC23* has not been identified as an eQTL.¹⁸

Risk variants are found to be enriched in regulatory elements and are therefore thought to regulate gene expression levels, often in a cell type- and context-specific manner (eg, stimulation and disease conditions). Overall, the relative participation of alternative promoters in the transcriptional output of genes, a previously underappreciated factor affecting disease development and complex trait associations, but the integration of puQTL and eQTL in various cell types and cell conditions provides new insights into the mechanisms underlying eQTL and helps explain the prevalence of specific transcript isoforms.¹⁸

ENHANCERS AND RISK VARIANTS

Enhancers are distal regulatory modules that boost the transcription of associated promoters, irrespective of their orientation. The first enhancer was identified in vitro decades ago as a distal sequence that increased the transcription of a reporter gene under the control of the SV40 promoter.²⁴ Enhancers typically span 50–1500bp and contain multiple TF binding sites, enabling them to orchestrate proper spatiotemporal gene expression during development³¹ or in response to signalling molecules.³² Enhancers are structurally and functionally similar to the proximal regions of promoters. Indeed, both promoters and enhancers are bound by multiple TFs, support the formation of the preinitiation complex, and recruit RNA Pol II.³³ Functionally, both promoters and enhancers support bidirectional transcription and activate transcript elongation.³³ In promoters, the sense strand expresses mRNA and the antisense strand

expresses promoter upstream transcripts (PROMPT/uaRNA). In enhancers, both the sense strand and the antisense strand express non-coding enhancer RNAs (eRNA) which are short, typically unspliced and non-adenylated transcripts.

Most GWAS variants have been found to overlap with cell type-specific enhancers that have been newly annotated through epigenomic profiling (table 1).^{34 35} Enhancer variants probably play an important role in fine-tuning the transcriptional output of each cell type and therefore affect susceptibility to common diseases. A number of classical methods to identify enhancers rely on the evolutionary conservation of non-coding sequences.^{35 36} In a landmark study, the candidate causal variants for 21 autoimmune diseases were integrated with RNA-seq data and chromatin features of control and stimulated human CD4⁺ T-cell subsets, regulatory T cells, CD8⁺ T cells, B cells and monocytes.³ Importantly, 90% of the causal variants were located in the non-coding regions of the genome. Approximately 60% of the variants were mapped to enhancers specific to immune cells, which showed increased histone acetylation and transcriptional activity on immune stimulation.³ The same study demonstrated that causal disease variants had a high probability of being context-specific immune enhancers and suggested that most non-coding causal variants alter atypical regulatory motifs rather than recognisable consensus motifs.³ Validation of enhancers and promoters identified through GWASs has historically been highly challenging.

More recently, next-generation sequencing technologies used in large-scale epigenomics projects have enabled the mapping and characterisation of regulatory regions in the human genome. Several methods have been developed to efficiently detect enhancers, such as chromatin immunoprecipitation-sequencing (ChIP-seq), which is used to identify TF binding or a histone modification in the enhancer sequence, and the DHS-seq and ATAC-seq methods, which are used to identify the open chromatin structure of activated enhancers. These methods were initially used in combination to detect enhancers.³⁷ As direct detection methods of regulatory regions, self-transcribing active regulatory region sequencing (STARR-seq) and CAGE-sequencing have been developed, which are used for assaying the enhancer activity of millions of enhancer candidates from arbitrary sources of DNA.³⁸ In combination with next-generation sequencing, CAGE-sequencing enables the genome-wide determination of coding and non-coding TSSs.^{20 33 39} The resulting database is instrumental to the identification of promoters and enhancers, both of which are characterised by the bidirectional transcripts. Enhancer activation leads to the bidirectional production of low levels of eRNAs.⁴⁰

EFFECTS OF LNCRNAs AND OTHER TRANSCRIPTS FROM NON-CODING REGIONS

The cumulative expression of non-coding genes is approximately four times greater than the expression of all mRNAs, making lncRNAs one of the most abundant RNA species within the average cell.⁹ The genomic and epigenomic characterisation of lncRNA-coding regions revealed that lncRNAs preferentially originate from enhancers rather than promoters.²⁰ lncRNAs do not encode proteins; however, lncRNAs longer than 200 nucleotides fold into complex structures and have biological functions. Functional lncRNAs can specifically interact with proteins, DNA and other RNAs to exert their molecular functions (figure 3). Folded lncRNAs have a variety of functions, such as transcriptional regulation (figure 3A), recruitment of TFs and proteins (figure 3B), degradation of transcript and splicing variants (figure 3C), and chromatin remodelling (figure 3D).^{20 41 42}

Table 1 Genome-wide association studies (GWAS)-associated expression quantitative trait loci (eQTL) affecting gene regulatory elements

#	GWAS-associated QTL variant	Chr	Variant genomic position	Associated disease	Affected DNA regulatory element	Affected gene(s)	Gene TSS genomic position (Refseq)	Genomic distance variant to TSS (bp)	Reference paper
1	rs34481144	11	320 836	Influenza infection	Core promoter	IFITM3	320 813	23	65
2	rs71542466	6	32 666 728	/	Proximal promoter	HLA-DQB1	32 666 607	121	66
3	rs115662534(T)	20	46 117 814	/	Proximal enhancer	CD40	46 118 344	−530	67
	rs548231435(C)	20	46 118 145	/	Proximal promoter	CD40	46 118 344	−199	
4	rs4794067	17	47 731 462	Autoimmune hepatitis—related to SLE and periodontitis	Proximal enhancer (?)	TBX21 (Tbet)	47 733 455	−1993	68
5	rs10750836	11	69 048 055	SLE	Proximal promoter/proximal enhancer	TPCN2	69 048 998	−943	69
6	rs2275675	1	183 470 348	SLE	Proximal promoter/proximal enhancer	SMG7	183 472 621	−2273	70
7	rs6426749	1	22 384 980	Osteoporosis	Enhancer	LINC00339	22 025 504	359 476	71
8	rs174575	11	61 830 531	Colorectal cancer	Enhancer	FADS2	61 828 391	2140	72
						AP002754.2	61 655 702	174 829	
9	rs56069439	19	17 283 116	Breast cancer and ovarian cancer	Enhancer	ABHD8	17 301 616	−18500	73
10	rs4748812	10	22 550 154	Childhood acute lymphoblastic leukaemia	Enhancer	PIP4K2A	22 714 326	−164172	74
	rs11591377	10	22 124 373	Childhood acute lymphoblastic leukaemia	Enhancer	BMI1	22 326 450	−202077	
11	rs62510556	8	117 152 033	Type 2 diabetes	Enhancer	SLC30A8	116 950 217	201 816	75
12	rs7864322 rs12352658 rs7847449 and rs10759944 all in LD with the GWAS variant rs965513	9	97 786 652	Thyroid cancer	All variants in the same enhancer	FOXE1	97 853 915	−67263	76
			97 789 486			FOXE1	97 853 915	−64429	
			97 789 616			FOXE1	97 853 915	−64299	
			97 794 690			FOXE1	97 853 915	−59225	
			97 793 827			FOXE1	97 853 915	−60088	
			97 786 652			PTCSC2	97 853 080	−66428	
			97 789 486			PTCSC2	97 853 080	−63594	
			97 789 616			PTCSC2	97 853 080	−63464	
			97 794 690			PTCSC2	97 853 080	−58390	
			97 793 827			PTCSC2	97 853 080	−59253	
13	rs1335532	1	116 558 335	Protection from multiple sclerosis	Enhancer	CDS8	116 570 972	−12637	77
14	rs2159100	12	2 237 227	Schizophrenia	Enhancer	CACNA1C	2 052 987	184 240	78
15	rs3181077	3	46 204 161	Narcolepsy	Enhancer	CCR1	46 208 313	−4152	79
						CCR3	46 163 604	−46163604	
16	rs7163757	15	62 099 409	Type 2 diabetes	Enhancer	VPS13C	61 854 457	244 952	80
17	rs148314165	6	137 908 854	SLE	Enhancer	TNFAIP3	137 871 228	37 626	81
	rs200820567		137 908 903			TNFAIP3	137 871 228	37 675	
18	rs139767239, rs115133228, rs12722502, rs12722635, in LD with rs706778 and rs706779	10	6 058 827	T1D	Intronic enhancers	IL2RA	6 062 151	−3324	82
			6 054 123				6 062 151	−8028	
			6 051 176				6 062 151	−10975	
			6 051 177				6 062 151	−10974	
			6 056 986				6 062 151	−5165	
			6 056 861				6 062 151	−5290	
19	rs17810546	3	159 947 262	Coeliac disease	Silencer	IL12A	159 989 057	−41 795	83

bp, base pair; Chr, chromosome; SLE, systemic lupus erythematosus; T1D, type 1 diabetes; TSS, transcription start site.

lncRNAs are a key component of the repertoire of regulatory elements that control cell differentiation and maintain tissue homeostasis. Recently, lncRNAs have been clearly implicated in

immune cell lineage commitment and immune responses (table 2). They have also emerged as an important class of molecules implicated in various human cancers as well as cardiovascular,

Figure 3 Genetic variants and long non-coding RNAs (lncRNAs). Genetic variants can affect the quantification and function of lncRNAs. (A) lncR-QTL, (B) regulation of transcription by lncRNAs, (C) binding of RNAs and lncRNAs, resulting in the formation of alternative splicing isoforms, and (D) interaction of lncRNAs with chromatin-modifying enzymes and nucleosome-remodelling factor. QTL, quantitative trait loci.

neurodegenerative, and autoimmune diseases.^{43 44} If variants with eQTL effects are located on lncRNAs, they may affect transcriptional regulators.²⁰ Therefore, the relationship between lncRNAs and risk variants in diseases and traits has received a significant amount of attention and should be the focus of future research.

Genome-wide lncR-eQTL analysis is expected to uncover how GWAS variants relate with lncRNA expression in common diseases and phenotypes. A comprehensive eQTL analysis was previously performed using genome-wide lncRNA expression and genotype data from human peripheral blood mononuclear cells of 43 unrelated individuals. Subsequently, various omics

Table 2 Examples of lncRNAs involved in immune cell functions			
Cell type	lncRNA	Function	Reference
Myeloid cells	Morrbid	Cis regulation of the Bcl2l11 gene	84
	NTT	Facilitation of cell differentiation and upregulation of IL10 and CXCL10	85
	PACER	Suppression of Cox2	86
	TCON_00019715	Promotion of macrophages to the M1 phenotype	87
	HOTAIR	IkB degradation	88
	Lnc-MC	Facilitation of monocyte differentiation and upregulation of ACVR1B	89
	LincRNA-COX2	Transcriptional suppression of target genes, for example, Ccl5 gene suppression via hnRNP A2/B and hnRNP A2/B1 binding	90
Dendritic cells	Lnc-DC	Suppression of IL6 and IL8 expression via STAT3 binding	47
	HOTAIRM1	monocyte/DC differentiation with miR-3960	91
	NEAT1	Induction of tolerogenic phenotype in DC	92
	Malat1	Induction of tolerogenic phenotype in DC	93 94
T cell	NeST	Upregulation IFNG gene expression	95
	NRON	Regulation of NFAT	96
	TH2-LCR	Regulation of cytokine expression in Th2 cells	97
	LncRNA-CD244	Inhibition of IFNG and TNF	98
	Linc-MAF-4	Regulation of MAF expression	42
	LicR-Ccr2-5'AS	Upreguration of Ccr gene expression	99
	NKILA	Inhibition of NF-kB activity	100

integrative network analyses were applied to construct variant-lncRNA-mRNA interaction networks, resulting in the detection of 707 pairs of cis-effect associations ($p < 5.64E-06$) and 6657 trans-effect associations ($p < 3.51E-08$).⁴⁵

The integration of genetic data sets from 1829 expression profiles in the FANTOM5 project uncovered the existence of nearly 20 000 functional lncRNAs. Approximately 2000 lncRNAs were expressed in significantly associated tissue/trait pairs. For example, the association between the middle temporal gyrus and autism spectrum disorder involves 18 lncRNAs from 49 genes.²⁰ On the other hand, a genetic variant over-represented in patients with coeliac disease lies within *lnc-13*, reduces its binding affinity for HDAC1, and weakens the epigenetic repression of inflammatory genes.⁴⁶ lncRNAs regulate multiple processes in disparate subsets of immune cells, such as the intensity of STAT3 signalling in human dendritic cells,⁴⁷ Th1 versus Th2 differentiation of human lymphocytes,⁴² and perhaps not surprisingly, the epigenetics-mediated training of immune responses in human monocytes.⁴¹ Thus, lncRNAs play an important role in gene regulation, and their profiling can reveal associations with specific tissues and cells.

FURTHER ANALYSES FOR FUNCTIONAL GENOMICS

Several different analyses have been used to annotate putative causal variants in GWASs, such as risk-associated SNPs in the promoter and enhancer regions (figure 2B; table 1).⁴⁸ Some of the functions of risk-associated enhancers and promoter variants have been experimentally proven, while others have been inferred from positional information; however, the functions of enhancers are particularly difficult to prove. In fact, the functions of many risk variants that exist outside of gene bodies remain unclear. Enhancers may be far away from the TSS, even further than 1 Mb. Transcriptional regulation from distal locations is facilitated by the folding of the genome, which places regulatory elements in three-dimensional proximity of the TSS. Advances in analytical techniques and next-generation sequencing techniques, like Hi-C-sequencing and micro-C-sequencing, allow the identification of trans-acting regulatory elements as well as cis-acting elements^{49–51} (figure 2B). Integrated analysis of omics data is also currently undergoing various statistical advances.

Biological analysis of regions regulated by TFs that were previously analysed by reporter assays can now be comprehensively performed through massive parallel reporter assays (MPRAs). In MPRAs, a library of barcoded candidate regulatory sequences is generated, cloned in the backbone of a reporter plasmid, and then transfected into an appropriate cell line. RNA-seq then allows the precise identification and quantitation of the transcriptional output associated with each regulatory variant. MPRAs facilitates the high-resolution, quantitative dissection of sequence-activity relationships of transcriptional regulatory elements with high reproducibility. MPRAs were developed to test the function of enhancer sequences in combination with targeted mutagenesis^{52 53} and have subsequently been exploited to test human promoters, enhancers⁵⁴ and eQTL.^{55 56} However, episomal reporter plasmids do not allow the testing of candidate regulatory sequences in their original context, a limitation that may affect the validation of epigenetic QTL. Chromatinised adenoviral and lentiviral vectors have therefore been developed to circumvent this limitation.^{57 58}

A new type of screening assay based on CRISPR/Cas9 technology examines the functions of regulatory regions in their original genomic context. CRISPR/Cas9 is used to test the function of candidate regulatory regions by targeted genomic deletion.⁵⁹ Other

CRISPR-based screening approaches, such as CRISPR/Cas9-mediated activation and inactivation, have been coupled with RNA-seq-based transcriptional quantification. However, these methods have not yet been scaled to the genome-wide level.⁶⁰ Furthermore, new approaches that combine CRISPR-based genome engineering with patient-specific induced pluripotent stem cell-based models are powerful tools for identifying putative causal loci that regulate gene expression and cellular functions.⁶¹

Advances in next-generation sequencing have improved single-cell gene expression analysis. Bulk RNA-seq experiments, commonly used in transcriptomic studies, examine the average gene expression among sampled cells and thus mask cell heterogeneity. However, single-cell RNA sequencing (scRNA-seq) can be used to reveal the heterogeneity of gene expression between various cells. Large-scale scRNA-seq can reveal patterns of gene, isoform, and allelic expression across different cell types and cell conditions.⁶² Most scRNA-seq methods use unique molecular identifiers (UMIs) of specific allelic origin.⁶² Utilisation of UMIs confers an important advantage to eQTL analysis, since they are useful to precisely understand allele imbalance. However, scRNA-seq analysis methods have low detection sensitivity and may not be able to detect many genes with low levels of expression, suggesting a need for improvement.⁶³ New statistical analyses have been developed to enable cell type identification, such as deconvolution analysis, which uses scRNA-seq and reference panel data. In addition to the transcriptomic analysis, scRNA-seq is also used in epigenomic analyses (eg, single-cell ChIP-sequencing). Additionally, spatial transcriptomics methods are also improving markedly.⁶⁴ In the future, spatial transcriptomics methods may enable the visualisation, while retaining the spatial information of the tissue.

CONCLUSIONS

As described above, considerable progress has been made in understanding the genetics and pathogenesis of autoimmune diseases. However, our understanding of the causality of these diseases is inadequate, and the development of new therapies based on genomic information for these diseases is required. Therefore, we need to apply a variety of approaches to identify the missing links.

GWASs are based on the hypothesis that commonly occurring diseases involve commonly occurring variants, with a large number of variants being involved in the development of a disease. Accordingly, functional analyses of risk variants are complicated. Statistical associations between genes and diseases alone cannot identify the risk variants that contribute to pathogenesis. Integration of GWAS data and omics data has the potential to elucidate the function of each risk variant while also identifying disease and tissue specificity. These findings will help reveal a more detailed mechanism of pathogenesis of autoimmune diseases and thus lead to their precise diagnosis and treatment.

Contributors AS and MMG carried out the experiment. AS wrote the manuscript with support from KY and MMG. KY supervised the project.

Funding This study was funded by Japan Society for the Promotion of Science (grant number 18H05285) and Chugai Pharmaceutical Co.

Competing interests KY has received grants from Chugai Pharmaceutical Co.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval Our study is performed in accordance with the International Conference on Harmonisation Guidelines for Good Clinical Practice and the Declaration of Helsinki, and the study protocols were approved by ethics review boards of RIKEN.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Kazuhiko Yamamoto <http://orcid.org/0000-0001-9037-3625>

REFERENCES

- Gallagher MD, Chen-Plotkin AS. The Post-GWAS era: from association to function. *Am J Hum Genet* 2018;102:717–30.
- Welter D, MacArthur J, Morales J, et al. The NHGRI GWAS catalog, a curated resource of SNP-trait associations. *Nucleic Acids Res* 2014;42:D1001–6.
- Farh KK-H, Marson A, Zhu J, et al. Genetic and epigenetic fine mapping of causal autoimmune disease variants. *Nature* 2015;518:337–43.
- Gabriel SB, Schaffner SF, Nguyen H, et al. The structure of haplotype blocks in the human genome. *Science* 2002;296:2225–9.
- Edwards SL, Beesley J, French JD, et al. Beyond GWASs: illuminating the dark road from association to function. *Am J Hum Genet* 2013;93:779–97.
- Maurano MT, Humbert R, Rynes S, et al. Systematic localization of common disease-associated variation in regulatory DNA. *Science* 2012;337:1190–5.
- Ishigaki K, Kochi Y, Yamamoto K. Genetics of human autoimmunity: from genetic information to functional insights. *Clin Immunol* 2018;186:9–13.
- Garrido-Cardenas JA, Mesa-Valle C, Manzano-Agugliaro F. Trends in plant research using molecular markers. *Planta* 2018;247:543–57.
- Ye Y, Zhang Z, Liu Y, et al. A multi-omics perspective of quantitative trait loci in precision medicine. *Trends Genet* 2020;36:318–36.
- Akiyama M, Ishigaki K, Sakaue S, et al. Characterizing rare and low-frequency height-associated variants in the Japanese population. *Nat Commun* 2019;10:4393.
- Kowalski MH, Qian H, Hou Z, et al. Use of >100,000 NHLBI Trans-Omics for Precision Medicine (TOPMed) Consortium whole genome sequences improves imputation quality and detection of rare variant associations in admixed African and Hispanic/Latino populations. *PLoS Genet* 2019;15:e1008500.
- Gerrits A, Li Y, Tesson BM, et al. Expression quantitative trait loci are highly sensitive to cellular differentiation state. *PLoS Genet* 2009;5:e1000692.
- Albert FW, Kruglyak L. The role of regulatory variation in complex traits and disease. *Nat Rev Genet* 2015;16:197–212.
- Gibson G, Powell JE, Marigorta UM. Expression quantitative trait locus analysis for translational medicine. *Genome Med* 2015;7:60.
- Patel L, Kang R, Rosenberg SC, et al. Dynamic reorganization of the genome shapes the recombination landscape in meiotic prophase. *Nat Struct Mol Biol* 2019;26:164–74.
- Braunschweig U, Gueroussov S, Plocik AM, et al. Dynamic integration of splicing within gene regulatory pathways. *Cell* 2013;152:1252–69.
- Chen L, Ge B, Casale FP, et al. Genetic drivers of epigenetic and transcriptional variation in human immune cells. *Cell* 2016;167:e24:1398–414.
- Garieri M, Delaneau O, Santoni F, et al. The effect of genetic variation on promoter usage and enhancer activity. *Nat Commun* 2017;8:1358.
- Zhao Q, Dacre M, Nguyen T, et al. Molecular mechanisms of coronary disease revealed using quantitative trait loci for TCF21 binding, chromatin accessibility, and chromosomal looping. *Genome Biol* 2020;21:135.
- Hon C-C, Ramiłowski JA, Harshbarger J, et al. An atlas of human long non-coding RNAs with accurate 5' ends. *Nature* 2017;543:199–204.
- Budach S, Heinig M, Marsico A. Principles of microRNA regulation revealed through modeling microRNA expression quantitative trait loci. *Genetics* 2016;203:1629–40.
- Haberle V, Stark A. Eukaryotic core promoters and the functional basis of transcription initiation. *Nat Rev Mol Cell Biol* 2018;19:621–37.
- Hampsey M. Molecular genetics of the RNA polymerase II general transcriptional machinery. *Microbiol Mol Biol Rev* 1998;62:465–503.
- Banerji J, Rusconi S, Schaffner W. Expression of a beta-globin gene is enhanced by remote SV40 DNA sequences. *Cell* 1981;27:299–308.
- Shlyueva D, Stampfel G, Stark A. Transcriptional enhancers: from properties to genome-wide predictions. *Nat Rev Genet* 2014;15:272–86.
- Spitz F, Furlong EEM. Transcription factors: from enhancer binding to developmental control. *Nat Rev Genet* 2012;13:613–26.
- Kumar A, Bansal M. Unveiling DNA structural features of promoters associated with various types of TSSs in prokaryotic transcriptomes and their role in gene expression. *DNA Res* 2017;24:25–35.
- Wu J, Metz C, Xu X, et al. A novel polymorphic CAAT/enhancer-binding protein beta element in the FasL gene promoter alters Fas ligand expression: a candidate background gene in African American systemic lupus erythematosus patients. *J Immunol* 2003;170:132–8.
- Kochi Y, Yamada R, Suzuki A, et al. A functional variant in FCRL3, encoding Fc receptor-like 3, is associated with rheumatoid arthritis and several autoimmunities. *Nat Genet* 2005;37:478–85.
- Hirabayashi S, Bhagat S, Matsuki Y, et al. NET-CAGE characterizes the dynamics and topology of human transcribed cis-regulatory elements. *Nat Genet* 2019;51:1369–79.
- Clyde DE, Corado MSG, Wu X, et al. A self-organizing system of repressor gradients establishes segmental complexity in Drosophila. *Nature* 2003;426:849–53.
- Kaikkonen MU, Spann NJ, Heinz S, et al. Remodeling of the enhancer landscape during macrophage activation is coupled to enhancer transcription. *Mol Cell* 2013;51:310–25.
- Andersson R, Sandelin A. Determinants of enhancer and promoter activities of regulatory elements. *Nat Rev Genet* 2020;21:71–87.
- Ernst J, Kheradpour P, Mikkelsen TS, et al. Mapping and analysis of chromatin state dynamics in nine human cell types. *Nature* 2011;473:43–9.
- Corradin O, Scacheri PC. Enhancer variants: evaluating functions in common disease. *Genome Med* 2014;6:85.
- Woolfe A, Goodson M, Goode DK, et al. Highly conserved non-coding sequences are associated with vertebrate development. *PLoS Biol* 2005;3:e7.
- Yañez-Cuna JO, Kvon EZ, Stark A. Deciphering the transcriptional cis-regulatory code. *Trends Genet* 2013;29:11–22.
- Jiang C, Li Y, Zhao Z, et al. Identifying and functionally characterizing tissue-specific and ubiquitously expressed human lncRNAs. *Oncotarget* 2016;7:7120–33.
- Shiraki T, Kondo S, Katayama S, et al. Cap analysis gene expression for high-throughput analysis of transcriptional starting point and identification of promoter usage. *Proc Natl Acad Sci U S A* 2003;100:15776–81.
- De Santa F, Barozzi I, Mietton F, et al. A large fraction of extragenic RNA Pol II transcription sites overlap enhancers. *PLoS Biol* 2010;8:e1000384.
- Fanucchi S, Fok ET, Dalla E, et al. Immune genes are primed for robust transcription by proximal long noncoding RNAs located in nuclear compartments. *Nat Genet* 2019;51:138–50.
- Ranzani V, Rossetti G, Panzeri I, et al. The long intergenic noncoding RNA landscape of human lymphocytes highlights the regulation of T cell differentiation by linc-MAF-4. *Nat Immunol* 2015;16:318–25.
- Esteller M. Non-Coding RNAs in human disease. *Nat Rev Genet* 2011;12:861–74.
- Han P, Chang C-P. Long non-coding RNA and chromatin remodeling. *RNA Biol* 2015;12:1094–8.
- Xia W, Zhu X-W, Mo X-B, et al. Integrative multi-omics analysis revealed SNP-lncRNA-mRNA (SLM) networks in human peripheral blood mononuclear cells. *Hum Genet* 2017;136:451–62.
- Castellanos-Rubio A, Fernandez-Jimenez N, Kratchmarov R, et al. A long noncoding RNA associated with susceptibility to celiac disease. *Science* 2016;352:91–5.
- Wang P, Xue Y, Han Y, et al. The STAT3-binding long noncoding RNA lnc-DC controls human dendritic cell differentiation. *Science* 2014;344:310–3.
- Ishigaki K, Kochi Y, Suzuki A, et al. Polygenic burdens on cell-specific pathways underlie the risk of rheumatoid arthritis. *Nat Genet* 2017;49:1120–5.
- Stevens TJ, Lando D, Basu S, et al. 3D structures of individual mammalian genomes studied by single-cell Hi-C. *Nature* 2017;544:59–64.
- de Souza N, Genomics deSN. Genomics. Micro-C maps of genome structure. *Nat Methods* 2015;12:812.
- Lu L, Liu X, Huang W-K, et al. Robust Hi-C maps of enhancer-promoter interactions reveal the function of non-coding genome in neural development and diseases. *Mol Cell* 2020;79:e15:521–34.
- Melnikov A, Murugan A, Zhang X, et al. Systematic dissection and optimization of inducible enhancers in human cells using a massively parallel reporter assay. *Nat Biotechnol* 2012;30:271–7.
- Smith RP, Taher L, Patwardhan RP, et al. Massively parallel decoding of mammalian regulatory sequences supports a flexible organizational model. *Nat Genet* 2013;45:1021–8.
- Nguyen TA, Jones RD, Snaveley AR, et al. High-Throughput functional comparison of promoter and enhancer activities. *Genome Res* 2016;26:1023–33.
- Tewhey R, Kotliar D, Park DS, et al. Direct identification of hundreds of Expression-Modulating variants using a multiplexed reporter assay. *Cell* 2016;165:1519–29.
- Ulirsch JC, Nandakumar SK, Wang L, et al. Systematic functional dissection of common genetic variation affecting red blood cell traits. *Cell* 2016;165:1530–45.
- Inoue F, Kircher M, Martin B, et al. A systematic comparison reveals substantial differences in chromosomal versus episomal encoding of enhancer activity. *Genome Res* 2017;27:38–52.
- Shen SQ, Myers CA, Hughes AEO, et al. Massively parallel cis-regulatory analysis in the mammalian central nervous system. *Genome Res* 2016;26:238–55.
- Rajagopal N, Srinivasan S, Kooshesh K, et al. High-Throughput mapping of regulatory DNA. *Nat Biotechnol* 2016;34:167–74.
- Xie S, Duan J, Li B, et al. Multiplexed engineering and analysis of combinatorial enhancer activity in single cells. *Mol Cell* 2017;66:285–99.
- Bassett AR. Editing the genome of hiPSC with CRISPR/Cas9: disease models. *Mamm Genome* 2017;28:348–64.
- Hagemann-Jensen M, Ziegenhain C, Chen P, et al. Single-Cell RNA counting at allele and isoform resolution using Smart-seq3. *Nat Biotechnol* 2020;38:708–14.
- Bagnoli JW, Ziegenhain C, Janjic A, et al. Sensitive and powerful single-cell RNA sequencing using mSCR-seq. *Nat Commun* 2018;9:2937.
- Burgess DJ. Spatial transcriptomics coming of age. *Nat Rev Genet* 2019;20:317.
- Allen EK, Randolph AG, Bhargale T, et al. SNP-mediated disruption of CTCF binding at the IFITM3 promoter is associated with risk of severe influenza in humans. *Nat Med* 2017;23:975–83.

- 66 Gutierrez-Arcelus M, Baglaenko Y, Arora J, *et al.* Allele-Specific expression changes dynamically during T cell activation in HLA and other autoimmune loci. *Nat Genet* 2020;52:247–53.
- 67 Putlyayeva LV, Demin DE, Korneev KV, *et al.* Potential Markers of Autoimmune Diseases, Alleles rs115662534(T) and rs548231435(C), Disrupt the Binding of Transcription Factors STAT1 and EBF1 to the Regulatory Elements of Human CD40 Gene. *Biochemistry* 2018;83:1534–42.
- 68 Sun W, Wu H-Y, Chen S. Influence of TBX21 T-1993C variant on autoimmune hepatitis development by yin-yang 1 binding. *World J Gastroenterol* 2017;23:8500–11.
- 69 Wen L, Zhu C, Zhu Z, *et al.* Exome-wide association study identifies four novel loci for systemic lupus erythematosus in Han Chinese population. *Ann Rheum Dis* 2018;77:417.
- 70 Deng Y, Zhao J, Sakurai D, *et al.* Decreased SMG7 expression associates with lupus-risk variants and elevated antinuclear antibody production. *Ann Rheum Dis* 2016;75:2007–13.
- 71 Chen X-F, Zhu D-L, Yang M, *et al.* An osteoporosis risk SNP at 1p36.12 acts as an allele-specific enhancer to modulate LINC00339 expression via long-range loop formation. *Am J Hum Genet* 2018;102:776–93.
- 72 Tian J, Lou J, Cai Y, *et al.* Risk SNP-Mediated Enhancer-Promoter Interaction Drives Colorectal Cancer through Both *FADS2* and *AP002754.2*. *Cancer Res* 2020;80:1804–18.
- 73 Lawrenson K, Kar S, McCue K, *et al.* Functional mechanisms underlying pleiotropic risk alleles at the 19p13.1 breast-ovarian cancer susceptibility locus. *Nat Commun* 2016;7:12675.
- 74 de Smith AJ, Walsh KM, Francis SS, *et al.* Bmi1 enhancer polymorphism underlies chromosome 10p12.31 association with childhood acute lymphoblastic leukemia. *Int J Cancer* 2018;143:2647–58.
- 75 Pound LD, Sarkar SA, Cauchi S, *et al.* Characterization of the human *Slc30a8* promoter and intronic enhancer. *J Mol Endocrinol* 2011;47:251–9.
- 76 He H, Li W, Liyanarachchi S, *et al.* Multiple functional variants in long-range enhancer elements contribute to the risk of SNP rs965513 in thyroid cancer. *Proc Natl Acad Sci U S A* 2015;112:6128–33.
- 77 Mitkin NA, Muratova AM, Korneev KV, *et al.* Protective C allele of the single-nucleotide polymorphism rs1335532 is associated with strong binding of ASCL2 transcription factor and elevated CD58 expression in B-cells. *Biochim Biophys Acta Mol Basis Dis* 2018;1864:3211–20.
- 78 Roussos P, Mitchell AC, Voloudakis G, *et al.* A role for noncoding variation in schizophrenia. *Cell Rep* 2014;9:1417–29.
- 79 Toyoda H, Miyagawa T, Koike A, *et al.* A polymorphism in *CCR1/CCR3* is associated with narcolepsy. *Brain Behav Immun* 2015;49:148–55.
- 80 Mehta ZB, Fine N, Pullen TJ, *et al.* Changes in the expression of the type 2 diabetes-associated gene *VPS13C* in the β -cell are associated with glucose intolerance in humans and mice. *Am J Physiol Endocrinol Metab* 2016;311:E488–507.
- 81 Wang S, Wen F, Tessner KL, *et al.* Talen-Mediated enhancer knockout influences *TNFAIP3* gene expression and mimics a molecular phenotype associated with systemic lupus erythematosus. *Genes Immun* 2016;17:165–70.
- 82 Schwartz AM, Demin DE, Vorontsov IE, *et al.* Multiple single nucleotide polymorphisms in the first intron of the *IL2RA* gene affect transcription factor binding and enhancer activity. *Gene* 2017;602:50–6.
- 83 Zwiers A, van Wanrooij RLJ, Dieckman T, *et al.* Celiac disease associated SNP rs17810546 is located in a gene silencing region. *Gene* 2020;726:144165.
- 84 Mowel WK, Kotzin JJ, McCright SJ, *et al.* Control of immune cell homeostasis and function by lncRNAs. *Trends Immunol* 2018;39:55–69.
- 85 Yang C-A, Li J-P, Yen J-C, *et al.* lncRNA *NTT/PBOV1* axis promotes monocyte differentiation and is elevated in rheumatoid arthritis. *Int J Mol Sci* 2018;19. doi:10.3390/ijms19092806. [Epub ahead of print: 18 Sep 2018].
- 86 Krawczyk M, Emerson BM. p50-associated COX-2 extragenic RNA (PACER) activates COX-2 gene expression by occluding repressive NF- κ B complexes. *Elife* 2014;3:e01776.
- 87 Xie Y, Wang M, Tian J, *et al.* Long non-coding RNA expressed in macrophage co-varies with the inflammatory phenotype during macrophage development and polarization. *J Cell Mol Med* 2019;23:6530–42.
- 88 Meredith EK, Balas MM, Sindy K, *et al.* An RNA matchmaker protein regulates the activity of the long noncoding RNA *HOTAIR*. *RNA* 2016;22:995–1010.
- 89 Chen M-T, Lin H-S, Shen C, *et al.* PU.1-Regulated long noncoding RNA *lnc-MC* controls human monocyte/macrophage differentiation through interaction with microRNA 199a-5p. *Mol Cell Biol* 2015;35:MCB.00429-15–24.
- 90 Basavappa M, Cherry S, Henao-Mejia J. Long noncoding RNAs and the regulation of innate immunity and host-virus interactions. *J Leukoc Biol* 2019;106:83–93.
- 91 Xin J, Li J, Feng Y, *et al.* Downregulation of long noncoding RNA *HOTAIRM1* promotes monocyte/dendritic cell differentiation through competitively binding to endogenous miR-3960. *Oncotargets Ther* 2017;10:1307–15.
- 92 Shui X, Chen S, Lin J, *et al.* Knockdown of lncRNA *NEAT1* inhibits Th17/CD4⁺ T cell differentiation through reducing the STAT3 protein level. *J Cell Physiol* 2019;234:22477–84.
- 93 Zhang M, Zheng Y, Sun Y, *et al.* Knockdown of *NEAT1* induces tolerogenic phenotype in dendritic cells by inhibiting activation of NLRP3 inflammasome. *Theranostics* 2019;9:3425–42.
- 94 Zhang Y, Zhang G, Liu Y, *et al.* GDF15 regulates MALAT-1 circular RNA and inactivates NF κ B signaling leading to immune tolerogenic DCs for preventing alloimmune rejection in heart transplantation. *Front Immunol* 2018;9:2407.
- 95 Gomez JA, Wapinski OL, Yang YW, *et al.* The nest long ncRNA controls microbial susceptibility and epigenetic activation of the interferon- γ locus. *Cell* 2013;152:743–54.
- 96 Imam H, Bano AS, Patel P, *et al.* The lncRNA *NRON* modulates HIV-1 replication in a NFAT-dependent manner and is differentially regulated by early and late viral proteins. *Sci Rep* 2015;5:8639.
- 97 Spurlock CF, 3Rd, Tossberg JT, Guo Y, Collier SP, Crooke PS, 3rd, Aune TM. expression and functions of long noncoding RNAs during human T helper cell differentiation. *Nat Commun* 2015;6:6932.
- 98 Wang Y, Zhong H, Xie X, *et al.* Long noncoding RNA derived from CD244 signaling epigenetically controls CD8⁺ T-cell immune responses in tuberculosis infection. *Proc Natl Acad Sci U S A* 2015;112:E3883–92.
- 99 Chen J, Ao L, Yang J. Long non-coding RNAs in diseases related to inflammation and immunity. *Ann Transl Med* 2019;7:494.
- 100 Zhang W, Guo Q, Liu G, *et al.* NKILA represses nasopharyngeal carcinoma carcinogenesis and metastasis by NF- κ B pathway inhibition. *PLoS Genet* 2019;15:e1008325.