

Supplementary Table 3. Multivariable Cox Proportional-hazard models, regressions adjusted for age and race. With remission as the outcome (DORIS definitions).

	All patients			Patients without Lupus Nephritis		
	Hazard Ratio	95% CI	P	Hazard Ratio	95% CI	p
Complete remission off treatment						
EULAR/ACR score	0.928	0.907-0-950	<.0001	0.926	0.897-0.955	<.0001
Age at first visit	1.004	0.995-1.013	0.3842	1.002	0.992-1.012	0.6484
Caucasian	1.773	1.205-2.609	0.0037	2.024	1.242-3.299	0.0047
Clinical remission off treatment						
EULAR/ACR score	0.052	0.935-0.969	<.0001	0.972	0.949-0.996	0.022
Age at first visit	0.997	0.989-1.005	0.5238	0.997	0.988-1.006	0.5168
Caucasian	1.435	1.085-1.898	0.0113	1.546	1.118-2.138	0.0084
Complete remission on treatment						
EULAR/ACR score	0.954	0.938-0.970	<.0001	0.949	0.927-0.972	<.0001
Age at first visit	1.01	1.003-1.017	0.0033	1.007	0.999-1.016	0.0683
Caucasian	1.352	1.062-1.722	0.0145	1.396	1.021-1.909	0.0364
Clinical remission on treatment						
EULAR/ACR score	0.982	0.971-0.992	0.0009	0.987	0.971-1.003	0.1003
Age at first visit	1.004	0.999-1.009	0.1371	1.002	0.996-1.008	0.5508
Caucasian	0.000	0.963-1.301	0.1417	1.161	0.961-1.403	0.1214