($p = 0.026$; OR=0.16; 95% CI=[0.03–0.79] and $p = 0.046$; OR=0.2; 95% CI=[0.04–1.00], while for *DRB1*03:01* negative association was revealed only for systemic JIA ($p = 0.03$; OR=0.11; 95% CI=[0.01–0.88]). At the same time the *DQB1*04:02:01* ($p=0.026$; OR=5.88; 95% CI=[1.20–28.72]) and *DRB1*08:01:01* ($p=0.07$; OR=3.94; 95% CI=[1.01–15.39]) allele frequencies were significantly higher in patients with oligoarthritis but not systemic JIA when compared with controls.

Conclusion: High-throughput HLA typing revealed distinct HLA-alleles associated with different JIA subtypes in the Belarusian population as well as some alleles protective both for oligoarthritic and systemic JIA.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1346

AB0004 THE ASSOCIATION OF THE RARE RS35667974 IFIH1 GENE POLYMORPHISM WITH SIX AUTOIMMUNE DISEASES: STRUCTURAL BIOLOGICAL INSIGHTS

M. Zervou¹, A. Andreou², G. Goulielmos¹, E. Eliopoulos². ¹University of Crete, Heraklion, Greece, Section of Molecular Pathology and Human Genetics, Department of Internal Medicine, School of Medicine, Heraklion, Greece; ²Agricultural University of Athens, Department of Biotechnology, Athens, Greece

Background: Genome wide association studies (GWAS) have successfully identified novel autoimmune disease-associated loci, with many of them shared by multiple disease-associated pathways but much of the genetics and pathophysiological mechanisms remain still obscure. Considering that most of the potential causal variants are still unknown, many studies showed that the missense variant rs35667974 at interferon induced with helicase C domain 1 (*IFIH1*) gene is protective for type 1 diabetes (T1D), psoriasis (PS) and psoriatic arthritis (PsA), while it was found to be also associated with ankylosing spondylitis (AS), Crohn's disease (CD) and ulcerative colitis (UC). *IFIH1* gene encodes a cytoplasmic RNA helicase that recognizes viral RNA and is involved in innate immunity through recognition of viral RNA [1].

Objectives: To investigate the role of the rare rs35667974 variant of *IFIH1* gene, which resides in exon 14 and changes a conserved isoleucine at position #923 to valine in T1D, PS, PsA, AS, CD and UC [2–4] as well as the mechanism that may affect the function in the protein structure.

Methods: Evolutionary analysis revealed heavily conserved sequence elements among species, indicating structural/functional importance of the mutation at position #923. *In silico* mutagenesis and three-dimensional (3D) homology modeling was used to localize the polymorphism under study on the *IFIH1* protein. The mutant was constructed using molecular modeling with the program Maestro (Schrodinger, LLC) [5]. Molecular mechanics/dynamics studies were applied to validate structural/functional changes caused by the Ile923V substitution. All figures depicting 3D models were generated using the PyMOL molecular-graphics system V.2.2 [6].

Results: Evolutionary and structural analysis revealed that the position of residue Ile923 is located on a protein loop (921–927) directly interacting with mRNA both to the phosphoribose chain and the base pairs. Mutation of Ile to Val at position #923 will directly affect the said interaction with mRNA [7] (Figure 1). Moreover, it has been reported that pre-mRNA or mRNA levels did not correlate with Ile923Val, suggesting that, Ile923Val did not alter the expression of *IFIH1* in a major way [7].

Figure 1. Structural representation of the interferon-induced helicase C domain-containing protein 1 [Homo sapiens] structure (4GL2 from Protein Data Bank) (in green) bound to the double stranded RNA (in orange-yellow). The location of the mutation (Ile923V) is highlighted in pink. The proximity of the aminoacid 923 to the nucleotide is apparent.

Conclusion: This study represents a comprehensive evaluation of the role of the shared rs35667974 variant of autoimmune locus *IFIH1*, reported to lead to a loss of function phenotype and, as a consequence, reduced levels of *IFIH1* protein and activity that protect against autoimmunity. Structural analysis of rare shared genetic susceptibility or protection loci may provide insight to our understanding of the pathophysiology of autoimmune diseases and the research findings may affect the better management of the diseases under study.

REFERENCES:

- [1] Nejentsev S *et al.* (2009). *Science* 324:387–389.
- [2] Smyth DJ *et al.* (2006). *Nat Genet.* 38: 617–619.
- [3] Li Y *et al.* (2010). *J Invest Dermatol* 130:2768–2772.
- [4] Ellinghaus D *et al.* (2016). *Nat Genet* 48:510–518.
- [5] Schrödinger Release 2017-1: Maestro, Schrödinger, LLC, New York, NY, 2017.
- [6] Schrödinger LLC: The PyMOL Molecular Graphics System 2016 version 2.2.
- [7] Wu B *et al.* (2013). *Cell* 152:276–289 [8] Downes K *et al.* (2010). *PLoS One* 5:e12646.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1388

AB0005 IDENTIFICATION OF KEY GENES AND PATHWAYS FOR PSORIASIS BASED ON GEO DATABASES BY BIOINFORMATICS ANALYSIS

X. Sun¹, S. X. Zhang^{2,3,4}, S. Song^{2,3,4}, T. Kong¹, C. Zheng¹, L. Cheng¹, S. Feng⁵, G. Shi⁶, X. Li^{2,3,4}, P. F. He⁷, Q. Yu^{1,7}. ¹Shanxi Medical University, School of Management, Taiyuan, China; ²The Second Hospital of Shanxi Medical University, Department of Rheumatology, Taiyuan, China; ³Shanxi Li Xiaofeng Medical Groups, Department of Rheumatology, Taiyuan, China; ⁴Ministry of Education, Key Laboratory of Cellular Physiology at Shanxi Medical University, Taiyuan, China; ⁵Shanxi Medical University, School of Humanities and Social Science, Taiyuan, China; ⁶Shanxi Medical University, Basic Medical College, Taiyuan, China; ⁷Shanxi Medical University, Institute of Medical Data Sciences, Taiyuan, China

Background: Psoriasis is an immune-mediated, genetic disease manifesting in the skin or joints or both, and also has a strong genetic predisposition and autoimmune pathogenic traits¹. The hallmark of psoriasis is sustained inflammation that leads to uncontrolled keratinocyte proliferation and dysfunctional differentiation. And it's also a chronic relapsing disease, which often necessitates a long-term therapy².

Objectives: To investigate the molecular mechanisms of psoriasis and find the potential gene targets for diagnosis and treating psoriasis.

Methods: Total 334 gene expression data of patients with psoriasis research (GSE13355 GSE14905 and GSE30999) were obtained from the Gene Expression Omnibus database. After data preprocessing and screening of differentially expressed genes (DEGs) by R software. Online tool Metascape³ was used to analyze Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis of DEGs. Interactions of proteins encoded by DEGs were discovered by Protein-protein interaction network (PPI) using STRING online software. Cytoscape software was utilized to visualize PPI and the degree of each DEGs was obtained by analyzing the topological structure of the PPI network.

Results: A total of 611 DEGs were found to be differentially expressed in psoriasis. GO analysis revealed that up-regulated DEGs were mostly associated with defense and response to external stimulus while down-regulated DEGs were mostly associated with metabolism and synthesis of lipids. KEGG enrichment analysis suggested they were mainly enriched in IL-17 signaling, Toll-like receptor signaling and PPAR signaling pathways, Cytokine-cytokine receptor interaction and lipid metabolism. In addition, top 9 key genes (CXCL10, OASL, IFIT1, IFIT3, RSAD2, MX1, OAS1, IFI44 and OAS2) were identified through Cytoscape.

Conclusion: DEGs of psoriasis may play an essential role in disease development and may be potential pathogenesis of psoriasis.

REFERENCES:

- [1] Boehncke WH, Schon MP. Psoriasis. *Lancet* 2015;386(9997):983-94. doi: 10.1016/S0140-6736(14)61909-7 [published Online First: 2015/05/31].
- [2] Zhang YJ, Sun YZ, Gao XH, *et al.* Integrated bioinformatic analysis of differentially expressed genes and signaling pathways in plaque psoriasis. *Mol Med Rep* 2019;20(1):225-35. doi: 10.3892/mmr.2019.10241 [published Online First: 2019/05/23].
- [3] Zhou Y, Zhou B, Pache L, *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun* 2019;10(1):1523. doi: 10.1038/s41467-019-09234-6 [published Online First: 2019/04/05].

Figure 1: (A) Heatmap of DEGs: The red and blue colors represent up-regulation and down-regulation, respectively. (B) Volcano map of gene expression: The horizontal axis is $\log_2$ FC, the vertical axis is $\log_{10}$ adj. P.Val. Each dot represents a gene, blue indicates low expression genes, red indicates high. (C, D, E, F) Visualization GO enrichment analysis and KEGG pathway enrichment analysis. (G, H) PPI network and hub gene identification: Top 9 hub genes in the PPI network were screened by Cytoscape plugin Cytobubba based on their degree value (high degree value) to yellow (low degree value).

Acknowledgements: This project was supported by National Science Foundation of China (82001740), Open Fund from the Key Laboratory of Cellular Physiology (Shanxi Medical University) (KLCP2019) and Innovation Plan for Postgraduate Education in Shanxi Province (2020BY078).

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1773

AB0006 GENETIC RISK PROFILE FOR PSORIATIC ARTHRITIS PREDISPOSITION IN ITALIAN PATIENTS

M. S. Chimenti¹, C. Ciccacci², G. De Benedittis³, A. Latini³, P. Conigliaro¹, G. L. Fonti¹, P. Triggianese¹, P. Borgiani³, R. Perricone¹. ¹University of Rome Tor Vergata, Rheumatology, Allergology and Clinical Immunology, Rome, Italy; ²UniCamillus - Saint Camillus International University of Health Sciences, Genetics Unit, Rome, Italy; ³University of Rome Tor Vergata, Genetics Unit, Rome, Italy

Background: Psoriatic arthritis (PsA) is a chronic inflammatory joint disease typically associated with psoriasis and classified in the group of spondyloarthritis (1). The pathogenesis is based on an interplay of different genes interacting with several environmental factors including stress, trauma, infections, triggering an inflammatory response related to the activation of innate and acquired immunity in different tissues and organs (2). However, the risk for the development of PsA is not clearly understood.

Objectives: The aim of this study was to evaluate, in a cohort of Italian PsA out-patients of the Rheumatology Unit of the University of Rome Tor Vergata, the association of genetic variants in candidate genes for PSA susceptibility and their possible contribute in the modulation of clinical and laboratory features.

Methods: The genes were selected according to previous studies describing these genes as involved in susceptibility to rheumatoid arthritis (RA) (3), since

a common genetic background can be shared between these diseases. Nine SNPs (single nucleotide polymorphism) in eight candidate genes were analysed: STAT4 (rs7574865), TRAF3IP2 (rs33980500), TNFAIP3 (rs6920220 and rs2230926), MIR146A (rs2910164), PSORS1C1 (rs2233945), IL-10 (rs1800872), HCP5 (rs3099844) and ERAP1 (rs27524). Polymorphisms were analysed in 163 consecutive PsA out-patients and 198 healthy controls (HC). Genotyping was performed by allelic discrimination by TaqMan assay. Alleles frequencies differences between cases and controls or between phenotypic groups were compared using Pearson's χ^2 test.

Results: We have observed an association between PSA susceptibility and the variant alleles of STAT4 [OR= 1.60 (1.15-2.21), P= 0.005], TRAF3IP2 [OR= 1.65 (1.01-2.65), P= 0.04], ERAP1 [OR= 1.40 (1.05-1.85), P= 0.02] and TNFAIP3 (rs6920220) [OR= 1.75 (1.19-2.57), P= 0.004]. On the contrary, the variant allele of IL-10 polymorphism seems to play a protective role [OR= 0.74 (1.05-1.85), P= 0.05]. Moreover, in order to define a genetic risk profile, we have counted the total number of risk alleles in each subject, considering as risk alleles the allelic variant of rs7574865 (STAT4), rs33980500 (TRAF3IP2), rs6920220 (TNFAIP3) and rs27524 (ERAP1) SNPs. Then, we have compared the risk allele number distribution between patients and HC (Fig.1). Classes with 3 or more risk alleles are significantly more represented in patients than in HC (OR= 2.03, P=0.004). The risk to develop the disease increases significantly in subjects with at least four risk alleles (OR= 2.96, P=0.002).

Figure 1. Number of risk alleles in patients and controls: rs7574865 (STAT4), rs33980500 (TRAF3IP2), rs6920220 (TNFAIP3) and rs27524 (ERAP1) SNPs.

Conclusion: We confirm the associations between five SNPs, already studied in RA, and PSA susceptibility, suggesting a common inflammatory pathway in chronic inflammatory rheumatological diseases. Moreover, we show how the genotyping of only few associated SNPs could help to define a genetic risk profile for PSA development.

REFERENCES:

- [1] Calabresi E, et al. One year in review 2019: psoriatic arthritis. Clin Exp Rheumatol. 2020;38:1046-55.
- [2] Chimenti MS, Triggianese P, De Martino E, Conigliaro P, Fonti GL, Sunzini F, Caso F, Perricone C, Costa L, Perricone R. An update on pathogenesis of psoriatic arthritis and potential therapeutic targets. Expert Rev Clin Immunol. 2019 Aug;15(8):823-836.
- [3] Ciccacci C, et al. Polymorphisms in STAT-4, IL-10, PSORS1C1, PTPN2 and MIR146A genes are associated differently with prognostic factors in Italian patients affected by rheumatoid arthritis. Clin Exp Immunol. 2016;186:157-63.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1976

AB0007 STAT4 RS7574865 G/T POLYMORPHISM AS MARKER OF PREDISPOSITION TO SYSTEMIC LUPUS ERYTHEMATOSUS WITH JUVENILE ONSET

M. Kaleda¹, M. Krylov², I. Nikishina¹. ¹V.A. Nasonova Research Institute of Rheumatology, Pediatric Department, Moscow, Russian Federation; ²V.A. Nasonova Research Institute of Rheumatology, Genetic Department, Moscow, Russian Federation

Background: Systemic lupus erythematosus (SLE) is a chronic autoimmune disease with a significant genetic predisposition. Recent studies have identified STAT4 (signal transducers and transcription activators 4) as a susceptibility gene for SLE.

Objectives: To investigate the hypothesis of the association of STAT4 rs7574865 G/T polymorphism with the predisposition to SLE in children and its relationship with some of SLE manifestations.