($CI_{95\%}$ 0.83-0.98)). Lupus arthritis was more common in risk TT-genotype SLE carriers than in other SLE patients ($\chi^2=5.902$ $p=0.015$; $p_{2-1}=0.027$). We revealed significant increase of CT genotype (RUNX1 rs9979383) in healthy donors vs SLE patients ($\chi^2=4.14$; $p=0.042$; $dOR=0.53$ ($CI_{95\%}$ 0.29-0.98); $LR^+=0.69$ ($CI_{95\%}$ 0.45-0.99); $LR^-=1.3$ ($CI_{95\%}$ 1.01-1.56)). Lupus arthritis was more common in SLE CT-genotype carriers than in other SLE patients ($\chi^2=4.66$ $p=0.031$; $p_{2-1}=0.058$).

Significant differences in IL6 rs1800795, IL6R rs2228145 and IL6R rs4845618 genotypes distribution between studied groups were not found (χ^2 , $p=0.427$, $p=0.559$ and $p=0.407$, correspondingly) but GG-genotype (IL6 rs1800795) carriage in SLE patients was associated with increased APS frequency ($\chi^2=4.45$, $p=0.035$; $dOR=0.19$ ($CI_{95\%}$ 0.04-0.9); $LR^+=0.28$ ($CI_{95\%}$ 0.07-0.93); $LR^-=1.41$ ($CI_{95\%}$ 1.03-1.64)).

Conclusion: Our data suggest the susceptibility to SLE in TT genotype of STAT4 rs7574865 polymorphism, protective role of CT genotype of RUNX1 rs9979383 for SLE and association between GG-genotype of IL6 rs1800795 and APS in SLE patients in Belarusian population. Lupus arthritis was associated with TT genotype of STAT4 rs7574865 and CT genotype of RUNX1 rs9979383.

REFERENCES:

- [1] Chen L, Morris DL, Vyse TJ. Genetic advances in systemic lupus erythematosus: an update. *Curr Opin Rheumatol* 2017;29:423–33.
- [2] Hochberg MC. Updating the American College of Rheumatology Revised Criteria for the classification of Systemic Lupus Erythematosus. *Arthritis Rheum* 1997;40:1725.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2523

AB0010 ASSOCIATION OF SAA1 GENE POLYMORPHISM -13T/C (RS12218) WITH ANKYLOSING SPONDYLITIS IN RUSSIAN POPULATION

I. Guseva¹, K. Sakharova², M. Krylov¹, E. Samarkina¹, S. Erdes². ¹V.A. Nasonova Research Institute of Rheumatology, Laboratory of Immunology and Molecular Biology of Rheumatic Diseases, Moscow, Russian Federation; ²V.A. Nasonova Research Institute of Rheumatology, Laboratory of Spondyloarthritis, Moscow, Russian Federation

Background: Ankylosing spondylitis is a chronic systemic inflammatory disease. Inflammation and high levels of serum amyloid A (SAA) protein are predisposing factors for secondary AA amyloidosis. The role of SAA1 gene polymorphisms in AS is not well understood.

Objectives: To investigate the association of SAA1 gene polymorphism -13T/C (rs12218) with ankylosing spondylitis and to evaluate the influence of this polymorphism on SAA protein concentration.

Methods: 123 AS patients (72 males, 51 females; age - M (SD) 37.51 (12.77) years; disease duration - 14.28 (11.22) years; BASDAI - 5.59 (1.13); B27-positive - 111 (90.2%) pts) and 95 gender, age matched healthy individuals (control group) were included in this study. SAA1 gene polymorphism -13T/C was genotyped using allele-specific RT-PCR assay. SAA protein concentration was measured using nephelometry in AS patients.

Results: The distribution of genotypes TT, TC and CC differed statistically between AS and control groups (24.4%, 56.1%, 19.5% and 41.1%, 44.2%, 14.7% respectively, $\chi^2=6.9$, $p=0.03$).

The presence of the C allele was associated with the development of AS ($OR=1.55$ [CI 1.04-2.33], $p=0.03$). The SAA1 -13T/C polymorphism tended to be associated with SAA protein value in AS patients: TT+TC genotypes -13.8mg/l [4.2; 91.0], CC genotype -7.8mg/l [1.6; 29.6], $p=0.07$. ESR, CRP and BASDAI values did not correlated with SAA1 -13T/C polymorphism ($p=0.6$, $p=0.4$, $p=0.4$ respectively).

Conclusion: The results of our study demonstrated for the first time that SAA1 gene polymorphism -13T/C (rs12218) is associated with susceptibility to AS. It is also shown that this polymorphism can affect the SAA protein level. Our findings need to be verified in AS patients with high levels of SAA protein in various ethnic and population groups.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2998

AB0011 EXPRESSION PROFILE AND POTENTIAL FUNCTION OF CIRCRNAS IN PERIPHERAL BLOOD MONONUCLEAR CELLS FROM PATIENTS WITH PRIMARY GOUT

Y. F. Qing¹, F. Dai¹, Q. B. Zhang², Y. P. Tang¹, Z. R. Dong¹, Y. X. He¹, Y. Jiang¹, Y. Q. Huang², J. Zheng¹. ¹Institute of Rheumatology and Immunology, Affiliated Hospital of North Sichuan Medical College, Department of Rheumatology and Immunology, Affiliated Hospital of North Sichuan Medical College, Nanchong, China; ²Institute of Rheumatology and Immunology, Affiliated Hospital of North Sichuan Medical College, Department of Geriatrics, Affiliated Hospital of North Sichuan Medical College, Nanchong, China

Background: Autophagy is a phenomenon of "self-phagocytosis" in eukaryotic cells, which maintains cell homeostasis by transporting intracellular materials to lysosomes for degradation and recycling. In recent years, studies have shown that autophagy may be involved in the pathogenesis of rheumatoid arthritis(RA) [1], but its specific mechanism is still unclear.

Objectives: The expression levels of autophagy-related genes(ATG) unc-51-like kinase 1(ULK1), ATG13, ATG17, microtubule associated protein 1 light chain 3 (LC3), and P62 in peripheral blood mononuclear cells (PBMC) of patients with RA were detected, and their role and clinical significance in the pathogenesis of RA were explored.

Methods: Real-time fluorescent quantitative PCR was performed to detect the expression levels of ULK1, ATG13, ATG17, LC3, and P62 in PBMCs of 50 RA patients, 50 healthy controls (HC), and 25 moderate to severe RA patients before and after treatment. Then, t test, χ^2 test, Mann-Whitney U test, Pearson test were used for statistical analysis.

Results: 1.The levels of hsCRP, white blood cell(WBC), neutrophils(GR), platelet(PLT) and plateletcrit(PCT) in RA group were higher than those in HC group ($P<0.05$). Lymphocytes (LY), red blood cell(RBC), hemoglobin(HGB), hematocrit(HCT), mean corpuscular hemoglobin(MCH), mean red blood cell volume(MCV) and mean red blood cell hemoglobin concentration(MCHC) in RA group were lower than those in HC group ($P<0.05$). 2.The expressions of ULK1, ATG17, and LC3 in RA group were higher than those in HC group, while the expressions of P62 was lower than those in HC group($P<0.05$) (Figure 1). The correlation analysis suggested that ATG17 was positively correlated with tender joint count (TJC), swollen joint count (SJC), and health assessment questionnaire (HAQ) ($P<0.05$); ULK1 and HAQ were negatively correlated ($P<0.05$). 3. Compared with before treatment with TNFi, ATG17, HAQ, DAS-28, ESR, hsCRP, WBC, GR, PLT and PCT were significantly reduced after treatment ($P<0.05$); the expressions of RBC, HCT, MCV and MCH were significantly increased after treatment, ($P<0.05$); ULK1, ATG13, LC3, P62 and other related clinical and laboratory indicators were not significantly different before and after treatment with TNFi ($P>0.05$).

Figure 1. The expression levels of ATGs in HC and RA groups.

Conclusion: There is abnormal expression of autophagy genes in the peripheral blood of RA patients. ULK1, ATG17, LC3 and P62 may be related to the pathogenesis of RA, among them, ATG17 may regulate the pathogenesis of RA by participating in the TNF- α pathway.

REFERENCES:

- [1] Rockel Jason S, Kapoor Mohit, Autophagy: controlling cell fate in rheumatic diseases. *J. Nat Rev Rheumatol*, 2016, 12: 517-31.

Disclosure of Interests: Yu-Feng Qing Grant/research support from: Science and Technology Project of Nanchong City (no.18SXHZ0522), Fei Dai: None declared, Quan-Bo Zhang Grant/research support from: the National Natural Science Foundation of China(General Program) (no.81974250), and Science and Technology Plan Project of Sichuan Province (no.2018JY0257), Yi-Ping Tang: None declared, Zeng-Rong Dong: None declared, Yi-Xi He: None declared, Yi Jiang: None declared, Yu-Qin Huang: None declared, Jianxiong Zheng: None declared

DOI: 10.1136/annrheumdis-2021-eular.3081

AB0012 THE EFFECTS OF COMMON THERAPEUTICS ON AIRE METHYLATION STATUS AND THE LEVELS OF IL-16, IL-1B, AND IFN- γ IN LPS-INDUCED MACROPHAGE CELLS CAN SULPHASALAZINE'S INFLUENCE ON METHYLATION STATUS OF AIRE EXPLAIN ITS AUTOIMMUNE SIDE EFFECTS?

Z. B. Çelik¹, C. Gunaydin², A. K. Cengiz³. ¹19 Mayıs University Medical Faculty, Medical Biology, Samsun, Turkey; ²19 Mayıs University Medical Faculty, Pharmacology, Samsun, Turkey; ³19 Mayıs University Medical Faculty, Physical Medicine and Rehabilitation-Rheumatology, Samsun, Turkey

Background: The autoimmune regulator gene (*AIRE*) has very important role in self-tolerance. Estrogen induced hypermethylation of *AIRE* promoter was put forward as one of the reasons for susceptibility to autoimmunity in females. [1] Sulphasalazine is a commonly used drug in rheumatology and is known to be responsible for various autoimmune side effects.

Objectives: We aimed to investigate the effects of commonly used treatments on the methylation status of *AIRE* and levels of interleukin-16 (IL16), interleukin-1 β (IL1 β), and interferon- γ (IFN γ), in lipopolysaccharides (LPS)-induced RAW264.7 macrophage cells which mimic inflammatory state in rheumatoid arthritis in vitro.[2]

Methods: RAW264.7 cells were stimulated by LPS (1 μ /mL). Cell viability test was performed to determine drug concentrations. Following drug treatments, cell media was isolated for the determination of IL16, IL1 β , and IFN γ levels by ELISA. Also, the cells detached by trypsin-EDTA were used for DNA isolation and bisulfite modification. Then, the promoter methylation status of *AIRE* was analyzed with methylation-specific PCR and agarose gel electrophoresis.

Results: Our results demonstrated that the *AIRE* promoter is highly methylated in absence of any inflammatory stimulus. LPS treatment changed methylation status to unmethylated form. Leflunomide (LEF), sulfasalazine (SLZ) and methotrexate (MTX) significantly altered methylation status as the methylated ($p=0.041, p<0.001, p<0.001$, respectively). Surprisingly, SLZ treatment caused significantly more methylation in *AIRE* promoter than other drugs ($p<0.001$, Fig 1A). Analysis of IL16, IL1 β and IFN γ levels demonstrated that LPS caused potent increase in all three cytokines ($p<0.001$, Fig 1B-D). LEF and SLZ significantly prevented LPS-induced increase in these cytokines ($p=0.025, p<0.001$ for IL-16, $p<0.001$ for IL-1 β and IFN γ , Fig 1B-D). Although MTX exacerbated LPS-induced increase in IL16 levels, it inhibited LPS induced increase in IL1 β and IFN γ levels ($p<0.001$, Fig 1B-D).

Figure 1. PCR gel image. All data expressed as mean $\pm$ SD. *** $p<0.001$ versus control, ## $p<0.001$, # $p<0.05$ versus LPS, +++ $p<0.001$ versus sulfasalazine.

Conclusion: Our results suggest that although LEF, SLZ and MTX potently suppresses LPS-induced inflammatory cytokines, they affect *AIRE* methylation differently. Changes in the methylation status of *AIRE* are important for autoimmunity. Relative increase in methylation of *AIRE* promoter by SLZ usage can be responsible for its autoimmune side effects, like drug induced lupus or hypersensitivity reactions, which are not common in MTX and LEF usage.

IL16 is a key regulator of biological properties of CD4 $^{+}$ T cells, it also regulates migration of CD4 $^{+}$ CD25 $^{+}$ Treg cells.[3] MTX treatment was shown to increase Treg cells in early rheumatoid arthritis patients. LEF and SLZ's effect on Treg cells was shown to be different from that of MTX.[4] Our results demonstrated that MTX exacerbated LPS-induced increase in IL16 levels in contrary to LEF and SLZ. This difference may also be responsible for the different effect of these medications on Treg functions.

The different effect of commonly used disease modifying drugs on IL16 levels and methylation of *AIRE* promoter is interesting and deserves attention.

REFERENCES:

- Berrih-Aknin, Le Panse R, Dragin N. *AIRE*: a missing link to explain female susceptibility to autoimmune diseases. *Ann.N.Y.Acad.Sci.* 2018;1412(1):21-32.
- Lin Y-Y, Jean Y-H, Lee H-P et al. Excavatolide B attenuates rheumatoid arthritis through the inhibition of osteoclastogenesis. *Marine Drugs*, 2017; 15(1):9.
- Skundric DS, Cruikshank WW, Montgomery PC et al. Emerging role of IL-16 in cytokine-mediated regulation of multiple sclerosis. *Cytokine*, 2015;75(2):234-248.
- Oh JS, Kim Y-G, Lee SG et al. The effect of various disease modifying anti-rheumatic drugs on suppressive function of CD4 $^{+}$ CD25 $^{+}$ regulatory T cells. *Rheumatol Int.* 2013;33:381-388.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.3335

AB0013

ASSOCIATION BETWEEN STAT4 POLYMORPHISM AND MANIFESTATIONS OF SLE

V. Agarwal¹, S. Kakati¹, P. Debaruah². ¹Assam Medical College and Hospital, Medicine, Dibrugarh, India; ²Assam Medical College and Hospital, Anatomy, Dibrugarh, India

Background: SNP rs7574865, located within the third intron of STAT4 gene at chromosome 2, has been associated with susceptibility to SLE among different ethnic groups.^{1,2} Interestingly, we recently have documented an association between this gene and susceptibility to systemic lupus erythematosus (SLE) in Indian population.³

Objectives: To determine whether the STAT4 (rs7574865) SNP is associated with clinical and immunological manifestations in SLE.

Methods: The study was carried out on 100 unrelated SLE (SLICC criteria 2012) patients from North-East India. Genotyping of STAT4 rs7574865 SNP was done using Taqman probe and Real-Time Polymerase chain reaction. An association study was performed between the alleles and genotypes of STAT4 rs7574865 with the clinical and immunological manifestations included in the SLE SLICC classification criteria. For all analysis, the statistical significance was fixed at 5% level of significance ($p<0.05$).

Results: The mean duration of illness was 2.69 $\pm$ 2.55 years. Cases and Controls remained in Hardy-Weinberg equilibrium. The occurrence of Photosensitivity and hyperpigmentation was significantly higher in TT genotype group (97.22% and 77.77%, respectively) with $p<0.001$ in each case. SLE patients with nephritis (Albuminuria >500mg/24 hours) and elevated serum creatinine were both significantly higher in TT genotype group as compared to GT and GG ($p<0.001$ and $p=0.001$ respectively). The Anti-dsDNA antibody was significantly associated with TT genotype ($p<0.001$).

Conclusion: Our study provides evidence regarding the association between STAT4 rs7574865 gene polymorphism is risk factor for cutaneous manifestations, Lupus nephritis and Anti ds-DNA positivity in SLE. So, our findings reinforce the need for further association studies including prospective studies with larger subjects in order to replicate such findings.

REFERENCES:

- Graham RR, Ph D, Hom G, Ph D, Behrens TW, Bakker PIW De, et al. and the Risk of Rheumatoid Arthritis and Systemic Lupus Erythematosus. *N Engl J Med.* 2007;357(10):977-86.
- Yuan H, Feng JB, Pan HF, Qiu LX, Li LH, Zhang N, et al. A meta-analysis of the association of STAT4 polymorphism with systemic lupus erythematosus. *Mod Rheumatol.* 2010;20(3):257-62.
- Gupta V, Kumar S, Pratap A, Singh R, Kumari R, Kumar S, et al. Association of ITGAM, TNFSF4, TNFAIP3 and STAT4 gene polymorphisms with risk of systemic lupus erythematosus in a North Indian population. *Lupus.* 2018;27(12):1973-9.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.3470

AB0014

METHOTREXATE - IMPLICATIONS OF PHARMACOGENETICS IN THE TREATMENT OF PATIENTS WITH RHEUMATOID ARTHRITIS

R. Pinheiro Torres^{1,2}, F. Pimentel Dos Santos^{1,2,3}, J. Branco^{1,2,3}. ¹Hospital Egas Moniz, Rheumatology, Lisboa, Portugal; ²CEDOC, Rheumatology, Lisboa, Portugal; ³Nova Medical School, Rheumatology, Lisboa, Portugal

Background: Methotrexate (MTX) is an anti-folate drug with anti-proliferative and anti-inflammatory effects. MTX proved to be the most highly effective, fast-acting disease modifying anti-rheumatic drug (DMARD), being widely used for the treatment of rheumatoid arthritis (RA) (1).

Objectives: This review aims to describe the main genetic variants identified concerning proteins that play a role in methotrexate's kinetics and efficiency profile.

Methods: A literature review was conducted since January of 2000 until December 2020, by searching the PubMed and Embase bibliographic databases, employing the following MeSH terms: methotrexate, pharmacogenetics, pharmacokinetics and rheumatoid arthritis. The search was limited to articles in English language. Two independent reviewers screened the titles and abstracts followed by a full-text review to assess papers regarding their eligibility. A total of 48 articles matched the research criteria and were analysed.

Results: Genetic variants of four main proteins, with different functions, have been consistently described.

Reduced folate carrier 1 (RFC1), a constitutively expressed folate transport protein that has high affinity for MTX is responsible, almost exclusively, for the transport of folate and MTX into the cell. The most commonly studied variant of the gene is the 80G > A variant (rs1051266), mapped within exon 2, on chromosome 21. It seems to improve RA responses to MTX, clinical efficacy with long disease remission (2).