P. Alonso Lecue^{1,4}, J. Rodríguez Carrio⁵, D. Prieto-Peña^{1,3}, V. Portilla^{1,3}, R. Blanco^{1,3}, A. Corrales^{1,3}, J. M. Cifrián-Martínez^{1,4,6}, R. López-Mejías¹, M. A. González-Gay^{1,3,6,7}. ¹IDIVAL, Research Group on Genetic Epidemiology and Atherosclerosis in Systemic Diseases and in Metabolic Bone Diseases of the Musculoskeletal System, Santander, Spain; ²Hospital Universitario Marqués de Valdecilla, 'López Albo' Post-Residency Programme, Santander, Spain; ³Hospital Universitario Marqués de Valdecilla, Department of Rheumatology, Santander, Spain; ⁴Hospital Universitario Marqués de Valdecilla, Department of Pneumology, Santander, Spain; ⁵Faculty of Medicine, Universidad de Oviedo, Department of Functional Biology, Immunology Area, Oviedo, Spain; ⁶Universidad de Cantabria, School of Medicine, Santander, Spain; ⁷Faculty of Health Sciences, University of the Witwatersrand, Cardiovascular Pathophysiology and Genomics Research Unit, School of Physiology, Johannesburg, South Africa

Background: Interstitial lung disease (ILD) is one of the most significant complications of connective tissue diseases (CTD), leading to an increase of the morbidity and mortality in patients with CTD [1]. A specific T cell subset termed angiogenic T cells (TAng), that promote endothelial repair and revascularization, have been involved in the pathogenesis of CTD [2-4]. However, to the best of our knowledge, no information regarding the role of TAng in CTD-ILD* is available.

Objectives: To study, for the first time, the potential role of TAng related to vascular damage in CTD-ILD*.

Methods: Peripheral venous blood was collected from 40 patients with CTD-ILD* and three comparative groups: 44 CTD-ILD* patients, 21 idiopathic pulmonary fibrosis (IPF) patients and 20 healthy controls (HC). All subjects were recruited from the Rheumatology and Pneumology departments of Hospital Universitario Marqués de Valdecilla, Santander, Spain. Quantification of TAng was performed by flow cytometry. TAng were considered as triple-positive for CD3, CD31 and CXCR4.

Results: Patients with CTD-ILD* exhibited a significantly lower TAng frequency than CTD-ILD* patients ($p < 0.001$). Similar results were obtained when patients with CTD-ILD* were compared with HC ($p = 0.004$) although no difference was observed between CTD-ILD* and IPF. In addition, a significant increase of TAng frequency was shown in patients with CTD-ILD* in relation to IPF patients ($p < 0.001$), while no difference was observed between CTD-ILD* and HC.

Conclusion: Our results reveal a decrease of TAng frequency related to vascular damage in CTD-ILD*. Furthermore, we disclose that the presence of ILD is associated with lower TAng frequency.

REFERENCES:

- [1] Expert Rev Clin Immunol 2018;14(1):69-82.
- [2] Circulation 2007;116(15):1671-82.
- [3] Ann Rheum Dis 2015 74(5):921-7.
- [4] PLoS One 2017;12(8):e0183102.

Acknowledgements: Personal funds, VP-C: PREVAL18/01 (IDIVAL); SR-M: RD16/0012/0009 (ISCIII-ERDF); LL-G: INNVAL20/06 (IDIVAL); RP-F: START PROJECT (FOREUM); RL-M: Miguel Servet type I CP16/00033 (ISCIII-ESF).

Disclosure of Interests: Verónica Pulito-Cueto: None declared, Sara Remuzgo Martínez: None declared, Fernanda Genre: None declared, Belén Atienza-Mateo: None declared, Victor Manuel Mora-Cuesta: None declared, David Iturbe-Fernández: None declared, Leticia Lera-Gómez: None declared, Raquel Pérez-Fernández: None declared, Pilar Alonso Lecue: None declared, Javier Rodríguez Carrio: None declared, Diana Prieto-Peña: None declared, Virginia Portilla: None declared, Ricardo Blanco Speakers bureau: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen and MSD, Consultant of: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen and MSD, Grant/research support from: Abbvie, MSD and Roche, Alfonso Corrales: None declared, Jose Manuel Cifrián-Martínez: None declared, Raquel López-Mejías: None declared, Miguel A González-Gay Speakers bureau: Pfizer, Abbvie, MSD, Grant/research support from: Pfizer, Abbvie, MSD

DOI: 10.1136/annrheumdis-2021-eular.2676

AB0027

INCREASED CIRCULATING CD39+FOXP3+CD4+ TREG CELLS IN EARLY RHEUMATOID ARTHRITIS FACILITATE THE ANTIINFLAMMATORY ACTION OF METHOTREXATE

L. Nuño¹, A. Villalba¹, M. Novella-Navarro¹, I. Monjo¹, D. Peiteado¹, S. García-Carazo¹, A. Puig-Kröger², A. Balsa¹, M. E. Miranda-Carus¹. ¹Hospital Universitario La Paz, Rheumatology, Madrid, Spain; ²Gregorio Marañón Hospital, Immunology, Madrid, Spain

Background: Methotrexate (MTX) remains the first line of treatment in Rheumatoid Arthritis (RA)^{1,2}. Inhibition of AICAR transformylase by MTX results in augmented release of adenine nucleotides to the extracellular space¹; these are rapidly hydrolysed by the combined action of ectonucleotidases CD39 and CD79 rendering the antiinflammatory agent adenosine¹. CD39, the rate-limiting enzyme in this cascade, is highly expressed by a subset of human FoxP3+CD4+ regulatory T cells (Treg39+)²⁻⁴ and MTX may act synergistically with Tregs in the control of inflammation.

Objectives: To study the expression of CD39 on circulating Treg cells of untreated early Rheumatoid Arthritis (ERA) patients and its relation with the ex vivo effect of MTX.

Methods: Peripheral blood was drawn from 22 DMARD- and steroid- naïve ERA patients with a disease duration < 24 weeks, 15 longstanding RA patients (LRA,

disease duration > 2 years) and 37 age and gender-matched healthy controls (HC). LRA patients were receiving low-dose weekly MTX and were naïve to biologicals. 10 ERA patients who had achieved remission 12 months after initiating MTX donated blood for a second time (ERA-R). The frequency of Treg and Treg cell subsets was assessed by flow cytometry. CD4+CD25+CD127- (total T reg), CD4+CD27+CD127-CD39+ Treg (Treg39+) and CD4+CD25-CD39- responder T (Tresp39-) cells were isolated by Ficol-Hypaque, followed by sorting. The suppressor potency of Tregs was assessed in cocultures of isolated Tregs with Tresp, established at different Treg/Tresp ratios. Proliferation was determined by CFSE dilution; cytokine secretion was measured by ELISA of culture supernatants.

Results: As previously described⁵, ERA but not LRA patients demonstrated a superior frequency of circulating Treg (CD4+CD25+CD127-FoxP3+) cells. In addition, the proportion of Tregs that expressed CD39 (Treg39+) was significantly increased in ERA but not LRA. Total ERA Tregs were significantly more potent suppressors of proliferation, TNF α and IFN γ secretion when compared with HC or LRA Tregs, and this difference was partially and significantly abrogated in the presence of adenosine deaminase, or the adenosine A2A receptor (A2AR) antagonists DMPX (3,7-dimethyl-1-propargylxanthine) or ZM 241385, but not in the presence of the adenosine A1 receptor antagonist DPCPX (8-cyclopentyl-dipropylxanthine). When MTX was added to the culture medium, the suppressor potency of total Tregs was further enhanced in all 3 groups of patients, and this enhancement was significantly higher in ERA total Treg/Tresp39- cocultures as compared with HC or LRA total Treg/Tresp39- cocultures. The effect of MTX was also partially and significantly abrogated by adenosine deaminase, DMPX or ZM 241385 but not by DPCPX. We then tested the suppressor potency of isolated Treg39+ together with the enhancer effect of MTX on this potency, and observed that there were no longer differences among ERA, LRA and HC; this further suggests that the differences observed in assays using total Tregs can be attributable to the increased Treg39+ proportions present in ERA. The frequency and function of ERA-R Treg cells were not different from HC or LRA Tregs.

Conclusion: The suppressor action of CD39+Tregs is mediated at least in part by adenosine through A2AR ligation, and the superior suppressive potency of total ERA Tregs is associated with their higher proportion of TregCD39+ cells as compared with HC or LRA. In addition, the augmented suppressor effect observed in the presence of MTX is partly mediated by an increased adenosine production acting on A2AR and is more marked in ERA patients reflecting again their higher proportion of Treg39+ cells. This indicates that MTX cooperates with Treg39+ cells in the control of inflammation.

REFERENCES:

- [1] Montesinos MC, et al. Arthritis Rheum. 2007.
- [2] Peres RS, et al. Proc Natl Acad Sci U S A. 2015.
- [3] Deaglio S, et al. J Exp Med. 2007.
- [4] Borsellino G, et al. Blood. 2007.
- [5] Benito-Miguel M, et al. J Immunol. 2009.

Acknowledgements: FIS PI 20/00141; FIS RD16/0012/0012; Fondo Europeo de Desarrollo Regional (FEDER), unrestricted research grant from Gebro Pharma

Disclosure of Interests: Laura Nuño: None declared, Alejandro Villalba: None declared, Marta Novella-Navarro: None declared, Irene Monjo: None declared, Diana Peiteado: None declared, Sara García-Carazo: None declared, Amaya Puig-Kröger: None declared, Alejandro Balsa Grant/research support from: BMS, Gebro Pharma, Maria-Eugenia Miranda-Carus Grant/research support from: BMS, Gebro Pharma

DOI: 10.1136/annrheumdis-2021-eular.2680

AB0028

EXPRESSION OF FATTY ACID TRANSPORTERS AND FREE FATTY ACID UPTAKE IN CD8+ T CELLS IN AUTOIMMUNE ARTHRITIS PATIENTS

S. Keck¹, F. V. Kraus¹, H. M. Lorenz¹, M. Souto-Carneiro¹. ¹University Hospital Heidelberg, Department of Rheumatology, Heidelberg, Germany

Background: RA CD8+ T cells (CD8s) have been shown to vastly contribute to tissue damage and RA disease activity by presenting and maintaining an active effector phenotype promoting autoinflammation. In order to meet their high metabolic requirements RA CD8s alter not only their glycolytic and glutaminolytic profiles, but also rely on increased lipid metabolism characterized by increased free fatty acid (FFA) uptake and lipid biosynthesis.

Objectives: In order to classify the importance of lipid metabolism in RA CD8s in terms of initiation and perpetuation of effector functions and therefore exploring potential novel therapy targeting points or biomarkers additional investigations have to be made. By analysing FFA uptake and the expression of free fatty acid transporters (FAT) upon TCR-stimulation in RA, PsA, SpA patients and healthy controls (HC), we aimed to further characterize the importance of RA CD8s' lipid metabolism.

Methods: Blood samples from 23 RA patients, 22 SpA patients, 20 PsA patients and 7 HC recruited at Heidelberg University Hospital were processed via negative magnetic separation selection to obtain purified CD8s. CD8s were cultured for 72 hours in [1,6-¹³C] glucose containing medium in the presence of anti-CD3 and anti-CD28. FFA uptake was measured by Palmitate-Bodipy 488 staining (ThermoFisher), the expression of FAT CD36, FABP4 and GPR84 besides CD69, CD36, CD45RA, CD3 and CD8 were assessed by FACS analysis.