AB0091

INHIBITION OF AUTOPHAGY PREVENTS PROGRESSION OF FIBROSIS IN MURINE MODELS OF SYSTEMIC SCLEROSIS

A. Zehender¹, Y. N. Li¹, N. Y. Lin¹, A. H. Györfi¹, A. Soare¹, C. Bergmann¹, A. Ramming¹, G. Schett¹, J. H. W. Distler¹. ¹Medicine 3 of the University Clinic Erlangen, Medizin 3, Erlangen, Germany

Background: Autophagy is catabolic process allowing cells to degrade unnecessary or dysfunctional cellular organelles. Failure of appropriate regulation of autophagy, however, can severely perturb tissue homeostasis. Recent studies demonstrate that autophagy is activated in several fibrotic diseases such as liver fibrosis, renal interstitial fibrosis, cardiac fibrosis.

Objectives: The objective of this work was to characterize the activation of autophagy in systemic sclerosis (SSc) and to decipher its role in the pathogenesis of SSc.

Methods: Activation of autophagy in skin samples of patients and murine models of SSc was assessed by co-staining of LC3B and P62 with the lysosomal marker LAMP2. The role of the autophagy was investigated in the model of bleomycin-induced dermal fibrosis. Beclin1 was overexpressed using adenovirus encoding for *Beclin1*. To knockdown *Atg7* *in vivo* was achieved by subcutaneous injections of *Atg7* siRNA or non-targeting siRNA. *In vivo*, 3-methyladenine (3-MA) was administered i.p. in a concentration of 15 mg/kg ones daily. Protein expression was measured by Western blot. Target genes were analyzed by qPCR. To monitor the autophagic flux, we generated adenoviral vectors encoding for tandem fluorescent-tagged LC3 (mRFP-EGFP-LC3).

Results: In the present study, we demonstrate that autophagy is activated in fibroblasts in SSc skin and also in experimental fibrosis models as compared to respective non-fibrotic control tissue with enhanced activity in *in vivo* and *in vitro* autophagy reporter studies. The aberrant activation of autophagy had profound stimulatory effects on fibroblasts. Activation of autophagy by forced expression of BECLIN1 promoted fibroblast-to-myofibroblast transition and stimulated the collagen release by cultured human fibroblasts and induced fibrosis in murine model. Nevertheless, inhibition of autophagy can deactivate myofibroblasts and induce regression of tissue fibrosis. Knockdown of *ATG7* or *BECLIN1* in human fibroblasts reduced the expression of α SMA and the number of stress fibers in myofibroblasts, indicating re-differentiation of myofibroblasts into resting fibroblasts upon inhibition of autophagy. Similar results were obtained with the autophagy inhibitors CQ and 3-MA. *In vivo*, siRNA mediated knockdown of *Atg7* effectively prevented progression of fibrosis in a model of established bleomycin-induced skin fibrosis. Inactivation of autophagy decreased dermal thickness, myofibroblast counts and hydroxyproline content to below pretreatment levels, indicating regression of bleomycin-induced skin fibrosis. In addition, treatment of mice with the autophagy inhibitor 3-MA ameliorated bleomycin-induced skin fibrosis.

Conclusion: We demonstrate that autophagy activity is enhanced in fibroblasts of SSc patients and in murine models of SSc. The increased activation of autophagy induces fibroblast-to-myofibroblast transition and promotes fibrotic tissue remodeling. However, inhibition of autophagy can deactivate myofibroblasts and induce regression of tissue fibrosis.

REFERENCES:

- Wynn, T. Cellular and molecular mechanisms of fibrosis. *J Pathol* 214, 199-210 (2008).
- Klionsky DJ, Abeliovich H, Agostinis P, *et al.* Guidelines for the use and interpretation of assays for monitoring autophagy in higher eukaryotes. *Autophagy* 4, 151-175 (2008).
- Wang, CW & Klionsky, DJ. The molecular mechanism of autophagy. *Mol Med* 9, 65-76 (2003).
- Hernández-Gea V, Ghiassi-Nejad Z, Rozenfeld R, *et al.* Autophagy releases lipid that promotes fibrogenesis by activated hepatic stellate cells in mice and in human tissues. *Gastroenterology* 142, 938-946 (2012).

Disclosure of Interests: Ariella Zehender: None declared, Yi-Nan Li: None declared, Neng-Yu Lin: None declared, Andrea-Hermina Györfi: None declared, Alina Soare: None declared, Christina Bergmann: None declared, Andreas Ramming: None declared, Georg Schett: None declared, Jörg H.W. Distler Consultant of: Actelion, Active Biotech, Anamar, ARXX, Bayer Pharma, Boehringer Ingelheim, Celgene, Galapagos, GSK, Inventiva, JB Therapeutics, Medac, Pfizer, Ruiyi and UCB., Grant/research support from: Anamar, Active Biotech, Array Biopharma, aTyr, BMS, Bayer Pharma, Boehringer Ingelheim, Celgene, Galapagos, GSK, Inventiva, Novartis, Sanofi-Aventis, RedX, UCB., Employee of: stock owner of 4D Science and Scientific head of FibroCure

DOI: 10.1136/annrheumdis-2021-eular.819

AB0092

ANTISYNTHEASE SYNDROME: CLINICAL PROFILE, SEROLOGIC AND TREATMENTS USED IN A COHORT OF PATIENTS FOLLOWED AT THE VIRGEN MACARENA HOSPITAL

P. Muñoz Reinoso¹, I. García Hernández¹, M. Ferrer Galván², F. J. Toyos Sáenz de Miera¹, L. Fernández de la Fuente Bursón¹, J. J. Pérez Venegas¹. ¹Hospital

Universitario Virgen Macarena, Reumatología, Sevilla, Spain; ²Hospital Universitario Virgen Macarena, Neumología, Sevilla, Spain

Background: The antisynthetase syndrome (SAS) is characterized by the presence of antisynthetase antibodies, anti-JO1, PL7 y PL12 are the most common; and the classic triad of myositis, arthritis, and diffuse interstitial lung disease (ILD)¹. Most patients present incomplete forms and the severity of the ILD determines the prognosis of the disease².

Objectives: to analyze epidemiological, clinical and serological characteristics and treatments used in a cohort of patients with SAS.

Methods: descriptive study of review of medical records. Data were collected from 15 patients with SAS followed in the Rheumatology and Pneumology consultations of the Virgen Macarena Hospital (Seville) in the last 10 years. The analysis was carried out using the R software.

Results: 15 patients were included, 8 men and 7 women. The median age was 56 years (33-77). Seven patients (47%) used to smoke. Four patients (27%) met the classical triad. All of them presented ILD and 8 patients (53%) had arthritis and / or myositis. Five (33%) had mechanic's hands and six of them (40%) presented Raynaud. Seven (47%) suffered from dyspnea before the SAS diagnosis. The median diagnostic delay was 1 month (0-43). Seven (47%) patients had anti-JO1, 1 (7%) anti-PL7, 2 (13%) anti-PL12 and 2 (13%) patients anti-Ro52. Radiological patterns detected by HRCT were: 5 (33%) NINE, 4 (37%) NIU and 6 (40%) others. The initial treatment included mostly (66%) glucocorticoids (GC) and one or more cFAME. In maintenance, mycophenolate was used in 7 patients (47%), cyclosporine 5 (33%), cyclophosphamide in 3 cases (20%), azathioprine in 3 patients (20%) and methotrexate in 3 of them (20%). Four (37%) patients required a combination of DMARDs and 2 cases needed (13%) biological therapy, Rituximab and Tocilizumab. Changes in the mean value of the initial respiratory function tests (FVC1 and DLCO1) and during follow-up (FVC2 and DLCO2) were not relevant (FVC1 81.5% [42-110], FVC2 81% [59-115]; DLCO1 83% [10-112], DLCO2 80.5% [47-108]). Nine patients (60%) remained clinically stable and 3 patients (20%) progressed radiologically. Four patients died from ILD progression.

Conclusion: In this study, the incomplete diagnosis of SAS predominated. The most detected antibody was anti-JO1. ILD is present in all cases, with NINE being the most frequent pattern so multidisciplinary management is necessary. Most used treatments were GC and FAMES combined, some cases required biological therapy.

REFERENCES:

- Irazoque F, *et al.* Epidemiology, etiology and classification. *Reumatol Clin.* 2009;5:2-5.
- Johnson C, *et al.* Clinical and pathologic differences in interstitial lung disease based on antisynthetase antibody type. *Respir Med.* 2014; 108(10):1542-8.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1427

AB0093

CLINICAL AND SEROLOGICAL CHARACTERISTICS OF SYSTEMIC SCLEROSIS: EXPERIENCE FROM A TERTIARY CARE CENTER IN PAKISTAN

S. Asif^{1,2,2}, M. Haroon¹, A. Khan², M. Faig¹. ¹Fatima Memorial Hospital, Lahore, Rheumatology, Lahore, Pakistan; ²Fatima Memorial Hospital, Lahore, Rheumatology, Lahore, Pakistan

Background: Systemic sclerosis (SS) is less studied Connective tissue disease in our population. It is characterized by different manifestations which if left undiagnosed and untreated lead to serious complications. The hallmark of this disease is fibrosis of various organs including skin and also involving pulmonary, gastrointestinal and cardiovascular system. We aim to evaluate the clinical and serological characteristics of SS in our population. In addition, we evaluated the prevalence of Fibromyalgia and Depression in patients with SS.

Objectives: We aimed to determine Clinical and Serological Characteristics of Systemic Sclerosis visiting our department of Rheumatology, Fatima Memorial hospital, Lahore.

Methods: It is a cross-sectional study of 38 patients visiting outdoor and indoor in Rheumatology department of Fatima Memorial Hospital from September 2019 to January 2020. All systemic sclerosis patients with age 16 years or above were included in this study. After taking informed consent, we filled the performas to record all the required information. We evaluated the clinical, serological and radiological features of Systemic Sclerosis.

Results: A total of 38 patients were included in the study with mean age was 34.47 ± 1.53 years. Female gender comprised of 32(84.2%), with male to female ratio of 1:5.3. Thirty-four (89.5%) had developed Raynaud phenomenon and sclerodactyly was found in 34(89.5%), digital ulcers in 25 (65.8%), and tendon friction rub in 12 (31.6%) patients. Skin tightness proximal to elbow was present in 9 (23.7%). Microstomia (decreased mouth opening) was present in 32(84.2%) patients. Twenty-eight (73.7%) patients had arthritis. Respiratory symptoms comprised of shortness of breath in 36(94.7%), palpitation in 24 (63.2%) and chest