	Univariate analysis	Multivariate analysis
	Odd Ratio (95% Confidence Interval)	
Gender	1.43 (0.76; 2.72)	
Disease duration	1.02 (0.99; 1.06)	
Age at TNF- α initiation	0.97 (0.95; 0.995)	0.97 (0.94; 0.999)
Magnetic sacroiliitis	0.46 (0.21; 0.99)	0.63 (0.26; 1.52)
Baseline BASDAI	1.02 (0.99; 1.04)	
Baseline CRP	0.99 (0.98; 1.02)	
Positive B27	0.81 (0.4; 1.68)	
Peripheral involvement	0.79 (0.4; 1.56)	
NSAIDs	2.91 (1.47; 5.77)	1.08 (0.41; 2.79)
PPIs	6.4 (3.25; 12.7)	6.9 (2.8; 17)
Antibiotics	1.87 (0.84; 4.16)	
Psychotics	1.94 (0.75; 4.97)	
Corticosteroids	1.11 (0.39; 3.11)	
Methotrexate	2.29 (0.97; 5.38)	

Conclusion: Co-medication with PPIs was considered as an independent factor associated with TNF- α failure. The hypothesis that this effect is due to their interference with the gut microbiota is only speculative but, regardless of the reason for this interaction, clinicians should be aware of the potential negative effect on TNF- α response.

REFERENCES:

- [1] Bazin, T., Hooks, K.B., Barnetche, T. *et al.* Microbiota Composition May Predict Anti-Tnf Alpha Response in Spondyloarthritis Patients: an Exploratory Study. *Sci Rep* 8, 5446 (2018). <https://doi.org/10.1038/s41598-018-23571-4>
- [2] Masson M, Kostine M, Barnetche T, Truchetet ME, Richez C, Schaefferbeke T. Ab0661 Co-Medications May Alter the Response to Tnf-Inhibitors in Spondyloarthritis Patients: A Pharmacomicrobiomic Effect? *Annals of the Rheumatic Diseases*. 1 juin 2020;79(Suppl 1):1625-6.

Disclosure of Interests: None declared.

DOI: 10.1136/annrheumdis-2021-eular.1716

AB0470

EXPERIENCE WITH THE USE OF INTERLEUKIN 6 INHIBITOR IN SEVERE ANKYLOSING SPONDYLITIS

D. Rumiantceva¹, E. Agafonova¹, M. Urumova¹, S. Erdes¹. ¹V.A. Nasonova Research Institute of Rheumatology, Laboratory of Spondyloarthritis and Psoriatic Arthritis, Moscow, Russian Federation

Background: Inhibitors of interleukin 6 (IL6) have been found to be ineffective in ankylosing spondylitis (AS) based on the results of randomized clinical trials (RCTs) on tocilizumab (TCZ) and sarilumab [1, 2]. However, there is evidence that IL6 is actively involved in the pathogenesis of the disease [3]. In addition, the efficacy of IL6 inhibitors has been shown in patients with secondary AA amyloidosis [4].

Objectives: to analyze the effectiveness of IL6 inhibitor - TCZ in patients with AS with high disease activity and secondary AA-amyloidosis.

Methods: The analysis included 10 patients with AS receiving TCZ therapy, of which 8 (80%) men, 2 (20%) women, and 100% HLA B27 positive. The average age of the patients was 40 $\pm$ 8.6 years, the average age of disease onset was 13.4 $\pm$ 7.5 years, the average duration of AS was 25.6 $\pm$ 6.5 years. 9 patients had histologically confirmed secondary AA-amyloidosis: 100% had kidney amyloidosis, 6 (66.6%) patients also had gastrointestinal amyloidosis and 3 (33.4%) patients had heart amyloidosis. All patients had arthritis and coxitis.

Results: As the first biological drug, TCZ was prescribed to 2 patients due to the presence of manifestations of secondary AA-amyloidosis, and 8 patients had previous experience of treatment with one or more inhibitors of tumor necrosis factor- α (TNF- α). The average duration of TCZ treatment was 22.0 months [1.0; 36.0]. 8 (80%) patients continue to take TCZ therapy to date, and 2 (20%) - are transferred to TNF- α inhibitors. Table 1 shows the main clinical and laboratory characteristics of patients at the initiation of TCZ therapy and during the last

Table 1. The main clinical and laboratory characteristics of patients at the initiation and after of TCZ therapy.

	Before TCZ	After TCZ	p
CRP, Me [25%; 75%], mg/l	95.2 [52.2; 189.0]	10.8 [0.8; 8.0]	p<0,05
Proteinuria, Me [25%; 75%], g/day.	1,6 [1,0; 2,1]	0,08 [0; 0,3]	p<0,05
BASDAI, Mean (SD)	6,1 (1,6)	3,3 (0,9)	p<0,05
ASDAS CRP, Mean (SD)	4,3 (1,1)	2,2 (0,7)	p<0,05
IBP, N (%)	10 (100%)	2 (20%)	p<0,05
Arthritis, N (%)	10 (100%)	2 (20%)	p<0,05
Cocitis, N (%)	10 (100%)	5 (50%)	p<0,05
Enthesitis, N (%)	7 (70%)	6 (60%)	p>0,05

hospitalization at the V.A. Nasonova Research Institute of Rheumatology. As you can see, against the background of TCZ therapy, both the level of CRP, proteinuria and the indices of AS activity (BASDAI, ASDAS CRP) significantly decreased. The number of patients with inflammatory back pain (IBP), arthritis and coxitis also decreased significantly.

Conclusion: The above data showed that the method of treating AS with IL6 inhibitors, in certain clinical situations, primarily such as ineffectiveness of TNF- α inhibitors, high CRP level and the development of secondary AA-amyloidosis, can be highly effective. Therefore, it is advisable to continue the set of clinical observations on the treatment of especially severe patients with AS treated with non-standard methods.

REFERENCES:

- [1] Sieper J, Porter-Brown B, Thompson L, *et al.* Assessment of short-term symptomatic efficacy of tocilizumab in ankylosing spondylitis: results of randomised, placebo-controlled trials *Ann Rheum Dis* 2014; 73: 95–100. doi:10.1136/annrheumdis-2013-203559.
- [2] Sieper J, Braun J, Kay J, *et al.* Sarilumab for the treatment of ankylosing spondylitis: results of a Phase II, randomised, double-blind, placebo-controlled study (ALIGN). *Ann Rheum Dis*. 2015; 74: 1051–1057. doi:10.1136/annrheumdis-2013-204963.
- [3] Londono J, Romero-Sanchez MC, Torres VG, *et al.* The association between serum levels of potential biomarkers with the presence of factors related to the clinical activity and poor prognosis in spondyloarthritis. *Rev Bras Reumatol* 2012; 52: 536–44.
- [4] Yilmaz S, Tekgöz E, Çınar M. Recurrence of proteinuria after cessation of tocilizumab in patients with AA amyloidosis secondary to FMF. *Eur J Rheumatol*. 2018; 5: 278–80. DOI: 10.5152/eurjrheum.2018.17183.

Disclosure of Interests: None declared.

DOI: 10.1136/annrheumdis-2021-eular.1740

AB0471

PATIENTS WITH RADIOGRAPHIC axSpA WHO PROGRESSED FROM ASAS20 AT WEEK 16 TO ASAS40 AT WEEK 52: RESULTS FROM COAST-W

X. Baraliakos¹, A. Deodhar², S. Liu Leage³, Y. Schymura⁴, R. Bolce⁵, D. Sandoval⁶, J. A. Walsh⁶, J. Sieper⁷. ¹Ruhr-University Bochum, Rheumatology, Bochum, Germany; ²Oregon Health & Science University, Division of Arthritis & Rheumatic Diseases, Portland, United States of America; ³Eli Lilly and Company Corporate Center, Rheumatology, Indianapolis, United States of America; ⁴Eli Lilly and Company Corporate Center, Statistics, Indianapolis, United States of America; ⁵Eli Lilly and Company Corporate Center, Medical Affairs, Indianapolis, United States of America; ⁶University of Utah, Rheumatology, Salt Lake City, United States of America; ⁷Charité - Universitätsmedizin Berlin, Campus Benjamin Franklin, Rheumatology, Berlin, Germany

Background: The timeframe for maximum treatment response varies across patients with radiographic axial spondyloarthritis (r-axSpA). Understanding which patients may benefit from additional time on treatment could influence treatment decisions.

Objectives: This post-hoc analysis aims to determine the percentage of patients, previously exposed to TNFi, who progressed from ASAS20 at week 16 to ASAS40 response at week 52 with ixekizumab (IXE) treatment and to explore factors that may associate with additional improvement after 16 weeks.

Methods: Patients who achieved Assessment of SpondyloArthritis International Society (ASAS)20 at week 16 from COAST-W (NCT02696798), a Phase 3, randomized, double-blind, placebo-controlled trial, in tumour necrosis factor inhibitor (TNFi)-experienced patients who fulfilled the ASAS criteria for r-axSpA, were analysed. Patients treated with IXE 80mg Q4W were categorized according to their ASAS response at week 52: sustaining ASAS20 but not reaching an ASAS40 response or achieving ASAS40. Patient demographics and disease characteristics at baseline were analysed by descriptive statistics, and the individual components determining ASAS response at baseline and at week 52 are provided.

Results: At week 16, 22.8% (n=26/114) of patients achieved ASAS20 but not ASAS40; of these, 2 patients discontinued the study before week 52. Amongst the patients who continued through week 52, 50% (12/24) of patients achieved ASAS40; the other 50% sustained their ASAS20 response. Patients who achieved ASAS40 at week 52 were older, had longer disease duration, were less likely to be HLA-B27 positive, and had worse BASDAI and BASFI scores at baseline (Table 1, part a). Achieving ASAS40 appeared to depend most on the Patient Global Assessment of Disease Activity and spinal pain score over time (Table 1, part b).

Table 1. a) Demographics and Characteristics at Baseline (week 0) and b) Descriptive observed data for the individual components of the ASAS response criteria at week 52, mean (SD)

a)	IXEQ4W	
	Patients sustaining an ASAS20 but not reaching an ASAS40 response at week 52 (n=12)	Patients achieving an ASAS40 response at week 52 (n=12)
Age, years	47.3 (13.3)	49.6 (12.3)
Duration of symptoms since axSpA onset, years	16.4 (9.2)	18.6 (12.6)
HLA B27 positive, n (%)	12 (100)	9 (75)
Current tobacco use, n (%)	4 (33.3)	1 (8.3)
ASDAS Score	4.0 (1.0)	4.2 (1.1)
BASDAI Score	7.4 (1.2)	7.8 (1.7)
Serum CRP concentration (mg/L)	23.0 (47.1)	28.0 (66.0)
Spinal Pain due to Ankylosing Spondylitis (NRS)	8.2 (0.8)	8.5 (1.6)
Patient Global Assessment of Disease Activity (NRS)	7.8 (1.3)	8.6 (1.2)
BASFI	7.1 (1.4)	7.4 (2.3)
BASDAI stiffness score	7.1 (1.5)	7.6 (2.2)
b)	IXEQ4W	
Patient Global Assessment of Disease Activity (NRS)	5.5 (1)	3.2 (1.3)
Spinal Pain due to Ankylosing Spondylitis (NRS)	5.6 (1)	3 (1.9)
BASFI	5.0 (1.5)	3.7 (2.1)
BASDAI stiffness score	4.1 (1.4)	3.4 (2.3)

Values are mean (SD) unless stated otherwise. ASAS= Assessment of SpondyloArthritis International Society; ASDAS= Ankylosing Spondylitis Disease Activity Score; axSpA = axial spondyloarthritis; BASDAI= Bath Ankylosing Spondylitis Disease Activity Index; BASFI= Bath Ankylosing Spondylitis Functional Index; CRP= C-reactive protein; HLA= human leucocyte antigen; NRS = numeric rating scale; SD= standard deviation

Conclusion: For patients with r-axSpA previously exposed to TNFi and showing modest response to IXE over 16 weeks, longer exposure to IXE may be required to achieve ASAS40.

Acknowledgements: The authors would like to acknowledge Philana Fernandes, an employee of Eli Lilly, for her editorial and writing support.

Disclosure of Interests: Xenofon Baraliakos Speakers bureau: Abbvie, BMS, Chugai, Celgene, Eli Lilly, Pfizer, Galapagos, UCB, MSD, Consultant of: Abbvie, BMS, Chugai, Celgene, Eli Lilly, Pfizer, Galapagos, UCB, MSD, Grant/research support from: Abbvie, Celgene, MSD, Atul Deodhar Consultant of: AbbVie, Amgen, Boehringer Ingelheim, Bristol Myers Squibb, Celgene, Eli Lilly, Giliad, GlaxoSmith & Kline, Janssen, Novartis, Pfizer, UCB, Grant/research support from: AbbVie, Boehringer Ingelheim, Eli Lilly, GlaxoSmith & Kline, Novartis, Pfizer, UCB, Soyi Liu Leage Employee of: Eli Lilly, Yves Schymura Employee of: Eli Lilly, Rebecca Bolce Shareholder of: Eli Lilly, Employee of: Eli Lilly, David Sandoval Shareholder of: Eli Lilly, Employee of: Eli Lilly, Jessica A. Walsh Consultant of: AbbVie, Novartis, Eli Lilly, UCB, Amgen & Pfizer, Grant/research support from: AbbVie, Pfizer, Merck, Joachim Sieper Speakers bureau: Abbvie, Janssen, Eli Lilly and Novartis, Consultant of: Abbvie, Janssen, Eli Lilly and Novartis.

DOI: 10.1136/annrheumdis-2021-eular.1823

AB0472

DISEASE ACTIVITY AND OUTCOME IN PREGNANCIES OF PATIENTS WITH SpA - DATA FROM THE GERMAN PREGNANCY REGISTER RHEIKSS

A. Weiß¹, C. Bungartz¹, J. Richter², S. Spaethling-Mestekemper³, X. Baraliakos⁴, P. M. Aries⁵, R. Fischer-Betz², A. Strangfeld¹. ¹German Rheumatism Research Centre, Epidemiology Unit, Berlin, Germany; ²University Hospital Düsseldorf, Department of Rheumatology & Hiller Research Unit, Düsseldorf, Germany; ³Rheumapraxis München-Pasing, Rheumatology, München, Germany; ⁴Rheumazentrum Ruhrgebiet, Rheumatology, Herne, Germany; ⁵Rheumatologie, Rheumatology, Hamburg, Germany

Background: Spondyloarthritis (SpA) is a severe chronic inflammatory disease, which affects quality of life and functional status. It frequently occurs in women of childbearing age. Active disease and TNFi discontinuation at early pregnancy were found to be risk factors for flares during pregnancy (1).

Objectives: To compare disease activity during pregnancy in patients with or without bDMARD exposure at conception and during pregnancy and to assess pregnancy outcomes.

Methods: RHEIKSS is a prospective longitudinal cohort study including patients with confirmed diagnosis of inflammatory rheumatic disease. Pregnant patients are eligible to be enrolled until the 20th week of gestation regardless of drug treatment. During observation, information on treatment, disease and pregnancy course, and outcome is collected from rheumatologists and patients. For this analysis, pregnancies of patients with SpA were selected and stratified into three groups according to their exposure to bDMARDs.

Results: Of 140 SpA pregnancies included, 74 (53%) were not exposed to bDMARDs at conception (group 1), 38 (27%) were exposed to bDMARDs at conception, but not during pregnancy (group 2) and 28 (20%) were continuously exposed to bDMARDs at conception and during pregnancy (group 3). Certolizumab (50%), Adalimumab (20%), Etanercept (8%) and Infliximab (8%) were the most frequently prescribed bDMARDs at beginning of pregnancy. Baseline characteristics according to treatment exposure are shown in Table 1. Frequency of flares was highest in group 2: 21%, 38%, and 39% of patients flared during the 1st, 2nd, and 3rd trimester. These rates were 20%, 25%, and 21% in group 1 and 8%, 20%, and zero in group 3. The difference in flare rates was also mirrored in the course of physician assessed global disease activity (Figure 1). Whereas patients in group 1 seemed to have a quite stable disease activity during pregnancy, those who were in group 2 had an increasing activity of disease during pregnancy with an even higher increase of disease activity after giving birth. Patients in group 3 had the lowest disease activity.

Of 137 singleton pregnancies, 130 (95%) ended in live birth. Of 6 spontaneous abortions 2 were in every of the three groups. One pregnancy in group 1 was terminated in gestational week 22 due to suspect malformation. One baby of the triple pregnancy was born and two aborted. All babies of the twin pregnancies were born healthy.

Conclusion: SpA patients treated with bDMARDs at conception are not at higher risk for adverse pregnancy outcomes. Our results in a larger patient population confirmed that discontinuation of bDMARDs after conception is associated with increased disease activity during pregnancy and after birth and a higher risk of flares.

REFERENCES:

- [1] van den Brandt S et al., Arthritis Res Ther. 2017; 19(1):64.

Table 1. Baseline characteristics; numbers are n (%) if not otherwise specified; * value at beginning of pregnancy; first 22 weeks after conception

Parameter	no bDMARD at conception (group 1) n=74	bDMARD at conception and discontinued during pregnancy (group 2) n=38	bDMARD at conception and continued during pregnancy (group 3) n=28	Total n=140
Singleton	72 (97)	37 (97.4)	28 (100)	137 (97.9)
Twin	1 (1.4)	1 (2.6)	0	2 (1.4)
Triple	1 (1.4)	0	0	1 (0.7)
New-York criteria fulfilled	21 (33)	17 (49)	10 (48)	48 (40)
disease duration in years, mean (SD)	6.4 (5.9)	7 (4.1)	5.8 (4)	6.4 (5.1)
age*, mean (SD)	33.4 (4.9)	32.3 (4)	31.6 (3.4)	32.7 (4.4)
severity of illness*: asymptomatic	4 (6)	0 (0)	3 (14)	7 (6)
mild	31 (48)	6 (17)	4 (19)	41 (34)
moderate	24 (38)	21 (60)	14 (67)	59 (49)
severe	5 (8)	8 (23)	0	13 (11)
HLA-B27 positive	41 (62)	24 (80)	15 (75)	80 (69)
CRP in mg/l *, mean (SD)	6.6 (8.2)	5.4 (8.2)	5.2 (4.9)	6 (7.6)
CRP >5mg/l *	25 (41)	9 (30)	8 (35)	42 (37)
physician global* (NRS 0-10), mean (SD)	2.6 (2)	2.3 (2.5)	1.7 (1.4)	2.4 (2.1)
BASDAI* (0-10), mean (SD)	3.2 (2)	2.9 (2.3)	2.8 (1.5)	3.1 (2)
patient global* (NRS 0-10), mean (SD)	3.3 (2.7)	3 (2.8)	3 (2.3)	3.1 (2.6)