AB0511

EFFECT OF SECUKINUMAB ON THE REDUCTION OF MYOCARDIAL DYSFUNCTION IN PATIENTS WITH ANKYLOSING SPONDYLITIS

S. Lapshina¹, L. Feiskhanova¹. ¹Kazan State Medical University, Department of Hospital Therapy, Kazan, Russian Federation

Background: In ankylosing spondylitis (AS), early subclinical changes in the myocardium mostly remain undiagnosed. The impact of genetic engineering biological therapy on these changes is also unclear.

Objectives: Aim: to establish the relationship between taking of secukinumab and changes in systolic and diastolic myocardial function in patients with AS.

Methods: 69 patients with AS were examined, of which the first group included 33 people (average age 38.8±4.74, 63.6% of men) who received the interleukin 17 inhibitor (IL17) - secukinumab, the second group - 36 people who did not receive biological therapy, average age 42.5±11 years, 66.7% of men. The control group included 40 healthy individuals, comparable in gender and age. Patients underwent tissue dopplerography of the heart and transthoracic echocardiography. The research materials were subjected to statistical processing using the program STATISTICA 10.0.

Results: The patients with AS had significantly higher left ventricular mass index and ejection fraction ($p < 0.01$) compared to the control group. Moreover, among patients who do not take biological therapy, the indicators were the highest (tab.1).

Table 1. Comparison of transthoracic echocardiography parameters in the study groups according to the Craskell-Wallis test.

Показатель	group 1	group 2	control group	p
LVMI	85 [81; 100]	93 [81,3; 114,2]	79,5 [72; 86,5]	$p = 0,002$
EF	63 [62; 64]	64,5 [63; 66]	60 [59,5; 64,5]	$p = 0,003$

LVMI – ventricular myocardial mass index, EF – ejection fraction.

A comparison of tissue dopplerography of the heart between the groups established statistically higher values of peak velocity of the ring of the mitral valve in early diastole (e'L) in the 1st group (14 [9,5; 14]) and control (13 [11; 15,5]) compared with the indicator of the second group (10,5 [8,3; 13]), $p < 0,01$; the peak speed of movement of the septum in early diastole, (e'S) in patients with secukinumab is higher (11 [9; 11]) compared to 2th (8,1 [7; 12]) and control groups 10 [8,5; 11]), $p < 0,05$, and in the 2nd group lower than in the control ($p = 0,03$). Significant differences in the parameters were supported by a comparison of the frequency of detection of diastolic dysfunction in the study groups. Diastolic dysfunction was diagnosed only in this group - in 38.9%. Against the background of the use of secukinumab, an increase in the systolic function of the myocardium was revealed.

Conclusion: The patients with AS are characterized by a high frequency of subclinical heart dysfunction. Inhibition of IL17 can reduce diastolic dysfunction, which is confirmed by an improvement in myocardial contractility during secukinumab therapy.

REFERENCES:

[1] ankylosing spondylitis, spondylarthritis, diastolic myocardial dysfunction, secukinumab

Disclosure of Interests: None declared.

DOI: 10.1136/annrheumdis-2021-eular.3474

AB0512

CLINICAL AND RADIOLOGICAL FEATURES OF SHOULDER INVOLVEMENT IN SPONDYLARTHROSIS

C. Ines¹, H. Ferjani¹, K. Maatallah¹, S. Miri¹, H. Hajji¹, M. Yasmine¹, D. Kaffel¹, W. Hamdi¹ on behalf of Rheumatology department of Med kassab institute of orthopaedics.. ¹Manouba, Rheumatology, Mannouba, Tunisia

Background: Peripheral joint involvement is frequent among patients with peripheral spondylarthritis (SA). However, the root joint involvement, such as hip involvement, has been described as being associated with axial spondylarthritis, while data for shoulder involvement are scarce. Cuff tendonitis and enthesitis are common features in the shoulder in patients with SA.

Objectives: This study aimed to identify the prevalence of shoulder involvement in SA patients and describe its clinical and radiological features.

Methods: We conducted a retrospective study including SA patients, all fulfilling the assessment of Spondyloarthritis International Society (ASAS) criteria.

For all patients, we collected the following data: Age, the clinical presentation of SA, the inflammatory biomarkers C-reactive protein (CRP), and the disease activity assessed by the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI).

Regarding shoulder involvement, we assessed the following items: tenderness, mobility, rotator cuff tests, as well as the results of X-ray and Ultrasound examination (US).

Results: We included one hundred and thirty-one SA patients (mean age 39,77 years). Among them, sixty-two percent were male. Ten percent of patients were smokers, 46,6% had the peripheral beginning of the disease. Fourteen patients complained of shoulder pain. The majority of them were male.

X-rays showed no abnormalities (n=2), while a destructive form was noted in (n=7).

Rotator cuff rupture or enthesopathy was also revealed by the reduction of sub-acromial space (n=5), condensation of the greater tuberosity (n=3), crooked aspect of the acromion (n=2), and Moloney's line disruption (n=1).

US showed no abnormalities (n=1), supraspinatus tendinopathy (n=5), supraspinatus transfixing tear (n=1), infraspinatus tendinopathy (n=2), supraspinatus enthesopathy (n=2), infraspinatus enthesopathy (n=1), moderate synovitis (n=1), subacromiodeltoid bursitis (n=1).

Shoulder involvement was correlated significantly with the peripheral beginning of SA ($P < 0.05$). There were no significant differences in gender, tobacco use, CRP value, and the disease activity between the two groups ($P > 0.05$).

Conclusion: The shoulder involvement was rare in our study. It is characterized by cuff tendonitis and enthesitis, especially in supraspinatus insertion. However, the glenohumeral synovitis was uncommon in our series, even in the peripheral form of the disease.

REFERENCES:

[1] Hip and Shoulder Involvement and Their Management in Axial Spondyloarthritis: a Current Review Clementina López-Medina^{1,2,3,4} & M. Carmen Castro-Villegas^{1,2,3} & Eduardo Collantes-Estévez^{1,2,3}, 23 July 2020.

Acknowledgements: I would like to thank all the authors listed above, for they helped me make writing my first abstract less harder.

Disclosure of Interests: None declared.

DOI: 10.1136/annrheumdis-2021-eular.3503

AB0513

THE RELATIONSHIP OF PERIARTICULAR TISSUES LESIONS AND JOINTS DEGENERATIVE CHANGES IN PATIENTS WITH INFLAMMATORY BOWEL DISEASES

G. Gainullina¹, E. Kirillova¹, D. Abduganieva¹, E. Belousova¹, A. Odincova².

¹Kazan State Medical University, Therapy, Kazan, Russian Federation;

²Republican Clinical Hospital, Gastroenterology, Kazan, Russian Federation

Background: Inflammatory bowel diseases (IBD) are associated with a variety of extra-intestinal manifestations. The most frequent of these is joint involvement, which affects 16–33 % of IBD patients, whereas 5–10 % are affected by enthesitis [1, 2]. Recent studies have shown the superiority of ultrasound over clinical findings in the evaluation of joints and periarticular tissues [3].

Objectives: To assess of joint and enthesal involvement in patients with Crohn's disease (CD) and ulcerative colitis (UC) using ultrasound with Power Doppler.

Methods: We prospectively included 70 IBD patients into the study. Peripheral joints and entheses were imaged by ultrasound, using Samsung Accuvix A30 5-13 MHz linear array transducer. Ultrasound examination of 14 peripheral joints (hip, knee, ankle, shoulder, acromioclavicular, elbow, wrist) and 35 entheses was performed. Vascularization was assessed with Power Doppler (PD). Statistical analysis was done by Mann-Whitney test by "Statistica" software.

Results: In 70 patients UC was in 40 (57%), CD - in 30 (43%). The mean age of patients was 32 (27; 36) years. The mean duration of the disease was 48 (12; 84) months. The majority of patients had highly active disease: moderate and severe activity was observed in 47 (67%).

Osteophytes were found in 32 (46%) IBD pts, in 10 pts (14%) - osteophytes were found in three or more joints. We divided pts into two groups. In first group we included pts without degenerative changes in the joints, in the second - with identified osteophytes.

Osteophytes were found more often in the acromioclavicular joints - 13 pts (41%) and in the hip joints - 10 pts (31%). With the same frequency (9 pts), osteophytes were detected in the wrist, knee and ankle joints. The mean age of pts and mean duration of IBD were significantly higher in the second group patients ($p = 0.00002$ and $p = 0.019$).

Enthesitis (echogenicity reduction and thickening) without vascularization were detected in 26 patients of the first group (68%) and 29 patients of the second group (91%) ($p = 0.009$). The rate of vascularized enthesitis in the two groups was the same: enthesitis with vascularization were observed in 14 pts of the first group (37%) and 14 of the second group (44%) ($p = 0.822$). Structure damage (erosions and enthesophytosis) was more frequent in the second group ($p = 0.025$ and $p = 0.001$). There were no significant differences in the incidence of tendinitis and tenosynovitis between two groups.

Conclusion: Joint and periarticular tissues damage in IBD patients increases with age and duration of the disease. The presence of degenerative changes in the joints is associated with more frequent detection of enthesitis and enthesopathies.

REFERENCES:

[1] The First European Evidence-based Consensus on Extra-Intestinal Manifestations in IBD. JCC. 2016; 10 (3): 239-254.