Methods: Patients from a longitudinal PsD cohort without a prior history of CVEs were included. In the first available serum sample, a targeted nuclear magnetic resonance (NMR) metabolomics platform was used to quantify 64 metabolite measures comprised of lipoprotein subclasses, fatty acids, glycolysis precursors, ketone bodies and amino acids. The study outcome included any of the following CVEs occurring within the first 10 years of biomarker assessment: angina, myocardial infarction, congestive heart failure, transient ischemic attack, cerebrovascular accident, revascularization procedures and CV death. The association of each metabolite with incident CVEs were analyzed separately using Cox proportional hazards regression models first adjusted for age and sex, and subsequently for traditional CV risk factors. Variable selection was performed using penalization with boosting after adjusting for age and sex. The added predictive value of the selected metabolites to improve risk prediction beyond traditional CV risk factors was assessed using the area under the receiver operator characteristic curve (AUC).

Results: A total of 977 patients with PsD, followed between 2002 and 2019, were analyzed (mean age 49.1 ± 12.6 years, 45.1% female). During a mean follow-up of 7.1 years, 70 (7.2%) patients developed incident CVEs. In Cox regression models adjusted for CV risk factors, alanine, tyrosine, total high-density lipoprotein (HDL) cholesterol, medium and large HDL particles, and the degree of unsaturation of fatty acids were significantly associated with decreased CV risk. Glycoprotein acetyls, apolipoprotein B, remnant cholesterol, very low-density lipoprotein (VLDL) cholesterol, and very small VLDL particles were associated with an increased CV risk. In proportional sub-distribution hazards regression models adjusted for age and sex, 13 metabolites were selected (Table 1). The age- and sex-adjusted expanded model (base model + 13 metabolites) significantly improved prediction of CVEs beyond the base model (only age and sex) with an AUC of 79.9 vs. 72.6, respectively ($p=0.019$) (Figure 1).

Table 1. Regression coefficients of the selected metabolites in a model adjusted for age and sex.

Category	Metabolite	Model adjusted for Age and Sex
Amino Acids	Alanine	-0.1179
	Glycine	-0.0339
	Tyrosine	-0.1010
Fatty acid ratios, relative to total fatty acids	Docosahexaenoic acid	-0.0862
	Unsaturation degree, double bonds per fatty acid	-0.1265
Fluid Balance	Albumin	+0.0685
Glycerides	Triglycerides in IDL cholesterol	+0.1546
Glycolysis precursors	Glucose	+0.1391
Inflammation	Glycoprotein acetyls	+0.1478
Ketone bodies	Acetoacetate	+0.0464
Lipoprotein subclasses	HDL ₃ Cholesterol	-0.0211
	Medium HDL	-0.0296
	Large HDL	-0.0309

Figure 1. Predictive performance of a model with age and sex alone is compared to a model with age and sex plus selected metabolites.

Conclusion: Using NMR metabolomics profiling, we identified a variety of metabolites associated with a lower and higher risk of developing CVEs in patients with PsD. Further study of their underlying association with CVEs is

needed to clarify the clinical utility of these biomarkers to guide CV risk assessment in this population.

REFERENCES:

- [1] Eder L, Wu Y, Chandran V, et al. Incidence and predictors for cardiovascular events in patients with psoriatic arthritis. *Ann Rheum Dis* 2016;75(9):1680-6.
- [2] Soininen P, Kangas AJ, Wurtz P, et al. Quantitative serum nuclear magnetic resonance metabolomics in cardiovascular epidemiology and genetics. *Circ Cardiovasc Genet* 2015;8(1):192-206.

Acknowledgements: Keith Colaco is supported by the Enid Walker Estate, Women's College Research Institute, Arthritis Society (TGP-19-0446), National Psoriasis Foundation (Early Career Grant) and the Edward Dunlop Foundation. Lihi Eder is supported by a Young Investigator Award from the Arthritis Society and an Early Researcher Award from the Ontario Ministry of Science and Innovation. The study was supported in part by a discovery grant from the National Psoriasis Foundation and an operating grant from the Arthritis Society (YIO-16-394).

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1154

OP0222 THE INCIDENCE OF INTERSTITIAL LUNG DISEASE IN PSORIATIC ARTHRITIS COMPARED TO RHEUMATOID ARTHRITIS. DATA FROM OVER 89 000 bDMARD TREATMENT COURSES DERIVED FROM FIVE NORDIC REGISTERS

S. Aarrestad Provan¹, B. Michelsen¹, L. Ljung^{2,3}, T. Jonmundsson⁴, B. Gudbjornsson⁴, D. Di Giuseppe⁵, M. L. Hetland^{6,7}, G. B. Reynisdóttir⁴, B. Blintborg^{6,7}, E. Kristianslund¹, H. Relas⁸, K. Aaltonen⁹, D. Nordström⁸, T. K. Kvien¹, J. Askling^{5,10}. ¹Diakonhjemmet Hospital, Division of Rheumatology and Research, Oslo, Norway; ²Clinical Epidemiology Section, Karolinska Institutet, Medicine Solna, Stockholm, Sweden; ³Stockholm Health Service, Center for Rheumatology, Academic Specialist Center, Stockholm, Sweden; ⁴University Hospital, Rheumatology, Reykjavik, Iceland; ⁵Clinical Epidemiology Section, Karolinska Institutet, Medicine Solna, Stockholm, Sweden; ⁶University of Copenhagen, Department of Clinical Medicine, Faculty of Health and Medical Sciences, Copenhagen, Denmark; ⁷Copenhagen Center for Arthritis Research, Rigshospitalet Glostrup, Center for Rheumatology and Spine Diseases, Copenhagen, Denmark; ⁸ROB-FIN, Helsinki University Hospital and Helsinki University, Department of Medicine and Rheumatology, Helsinki, Finland; ⁹Ministry of Social Affairs and Health, Pharmaceuticals Pricing Board, Helsinki, Finland; ¹⁰Karolinska University Hospital, Rheumatology, Theme Inflammation and Infection, Stockholm, Sweden

Background: Interstitial lung disease (ILD) is an established extra-articular manifestation of rheumatoid arthritis (RA). Few studies have investigated the prevalence of ILD in patients with psoriatic arthritis (PsA). Methotrexate (MTX) is frequently used in the treatment of both RA and PsA and has been suggested to be a risk factor for the development of ILD. It is of interest to understand the interaction between disease and treatment in the development of ILD.

Objectives: To compare the incidence of ILD between patient with PsA and RA treated with biologic disease modifying antirheumatic drugs (bDMARDs), with or without MTX as a co-medication.

Methods: Cohorts of patients with RA and PsA starting bDMARD were identified in Nordic registers (Danish nationwide clinical register for patients with RA (DAN-BIO), Register on antirheumatic and biological therapy in Finland (ROB-FIN), Icelandic nationwide database of biologic therapy (ICEBIO), Norwegian Anti-rheumatic Drug Register (NOR-DMARD), and the Swedish Rheumatology Quality Register (SRQ)). Linkages to the National Patient Registers and to the Cause of Death Registers were performed in each country to identify cases of ILD. Each individual patient could contribute several treatment courses. ILD was identified as hospital or death certificate ICD10 codes of ILD (J84.1, J84.8, J84.9, J70.2, J70.3, J70.4, J99.0, J99.1, J99.8) given during the follow-up period which was defined as the treatment course duration, plus a 30-day wash-out period added to the end of treatment course period. MTX co-medication was specified as use of MTX at the start of bDMARD. Incidence rates (IR) for any ILD were calculated per 1000 person years at risk (PYR) for each country. The five cohorts were pooled and incidence rate ratios (IRR) for PsA vs. RA were calculated. Hazard ratios (HR) for any ILD in PsA vs. RA were estimated in Cox regression models adjusted for age, gender and repeated observations, and stratified for the use of MTX co-medication.

Results: Overall 47 987 individual patients representing 89 239 bDMARD treatment courses and contributing 201 279 PYR were included in the study (Table 1). Methotrexate was reported as comedication in 29 916 (33.5 %) of the treatment courses (PsA vs. RA, 30.4 % vs 34.5 %). 970 cases of ILD were identified during the follow-up period. The risk of ILD was consistently lower in patients with PsA compared to patients with RA in all countries. In models stratified for co-medication the HR for ILD in PsA vs. RA was 0.34 (0.21-0.57) in patients treated with MTX and 0.26 (0.18-0.36) in patients not treated with MTX.

Table 1. Interstitial lung disease in PsA vs. RA in five Nordic biologic registers

	DENMARK		FINLAND		ICELAND		NORWAY		SWEDEN	
	RA	PsA	RA	PsA	RA	PsA	RA	PsA	RA	PsA
Number of individuals	7829	3386	4946	1091	675	470	1590	999	20596	6393
Number of treatment courses	17 072	6640	8634	1845	1280	859	2379	1427	38 279	10 824
Age baseline (SD)	57.3 (13.1)	49.0 (12.6)	53.8 (13.4)	48.8 (11.4)	53.9 (14.2)	50.1 (13.3)	53.8 (13.7)	48.7 (12.0)	57.1 (13.7)	50.6 (12.8)
Female n (%)	12 963 (76)	3929 (59)	6571 (76)	933 (51)	969 (76)	551 (65)	1815 (77)	818 (57)	29 635 (77)	6162 (57)
Number of PYR	40235	13986	21798	4910	4517	2799	4556	2653	120334	27412
ILD-events within PYR	218	22	132	8	7	2	32	6	680	28
IR pr 1000 PYR	5.4	1.6	6.1	1.6	1.5	0.7	7.0	2.3	5.7	1.0
IRR PsA vs RA crude	0.29 (0.18-0.45)		0.27 (0.11-0.55)		0.46 (0.05-2.42)		0.32 (0.11-0.78)		0.18 (0.12-0.26)	
HR PsA vs RA	0.31 (0.17-0.56)		0.46 (0.22-0.96)		0.62 (0.12-3.14)		0.19 (0.06-0.54)		0.25 (0.17-0.37)	

PYR: Patient years at risk, IR: Incidence rates, IRR: Incidence rate ratios, HR: Hazard Ratios

Conclusion: In these preliminary analyses, the incidence of ILD is lower in bDMARD treated PsA vs. RA patients, irrespective of co-medication with MTX. This indicates that the clinician should consider the rheumatological diagnosis when assessing the risk for future ILD in patients treated with bDMARDs and MTX.

Acknowledgements: Partly funded by NordForsk and FOREUM

Disclosure of Interests: Sella Aarrestad Provan Consultant of: Novartis, Grant/research support from: Boehringer-Ingelheim, Brigitte Michelsen: None declared, Lotta Ljung: None declared, Thorarinn Jonmundsson: None declared, Björn Gudbjörnsson Speakers bureau: Amgen and Novartis, Daniela Di Giuseppe: None declared, Merete Lund Hetland Speakers bureau: Orion Pharma, Biogen, Pfizer, CellTrion, Merck and Samsung Bioepis, Consultant of: Eli Lilly, Grant/research support from: BMS, MSD, AbbVie, Roche, Novartis, Biogen and Pfizer, Guðrún Björk Reynisdóttir: None declared, Bente Glintborg: None declared, Eirik kristianslund: None declared, Heikki Relas: None declared, Kalle Aaltonen: None declared, Dan Nordström Speakers bureau: Abbvie, BMS, Celgene, Eli Lilly, MSD, Novartis, Pfizer, Roche and UCB., Consultant of: Abbvie, BMS, Celgene, Eli Lilly, MSD, Novartis, Pfizer, Roche and UCB., Tore K. Kvien Speakers bureau: Amgen, Celltrion, Eisai, Evapharma, Ewopharma, Hikma, Oktal, Sandoz, Sanofi., Consultant of: AbbVie, Amgen, Biogen, Celltrion, Eli Lilly, Gilead, Mylan, Novartis, Pfizer, Roche, Sandoz, Sanofi., Johan Askling Grant/research support from: Abbvie, BMS, Eli Lilly, Merck, Pfizer, Roche, Samsung Bioepis, and Sanofi

DOI: 10.1136/annrheumdis-2021-eular.2955

OP0223

DEVELOPMENT AND PRELIMINARY VALIDATION OF ULTRASONOGRAPHIC DISEASE ACTIVITY AND DAMAGE SCORES IN PSORIATIC ARTHRITIS PATIENTS: RESULTS FROM THE UPSTREAM (ULTRASOUND IN PSORIATIC ARTHRITIS TREATMENT) STUDY

A. Zabotti¹, M. Piga², A. Zanetti³, M. Canzoni⁴, N. Boffini⁵, V. Picerno⁶, G. Zanframundo⁷, E. Silvagni⁸, I. Giovannini¹, B. Raffener⁹, P. Scolieri¹⁰, P. Mancini¹¹, S. Parisi¹², A. Bortoluzzi⁸, G. Sakellariou¹³, O. De Lucia¹⁴, I. Tinazzi¹⁵, F. Figus¹⁶, L. Idolazzi¹⁷, M. Lorenzin¹⁸, S. Z. Callegger¹, A. Cauli¹⁹, G. Carrara³, C. A. Scirè³, A. Iagnocco¹⁶, on behalf of The Musculoskeletal Ultrasound Study Group of the Italian Society of Rheumatology. ¹University of Udine, Rheumatology Clinic, Santa Maria della Misericordia Hospital, Medical and Biological Sciences Department, Udine, Italy; ²Rheumatology Unit, AOU and University of Cagliari, Department of Medical Sciences and Public Health, Monserrato, Italy; ³Italian Society for Rheumatology, Epidemiology Research Unit, Milano, Italy; ⁴Local Health Unit (ASL) Rome-1, Rome-4 and Viterbo, Rheumatologist Outpatient Specialist, Rome, Italy; ⁵IRCCS Ospedale S. Raffaele, UNIRAR Department, Milano, Italy; ⁶Azienda Ospedaliera San Carlo, Dipartimento internistico, Potenza, Italy; ⁷Pavia, Department of Internal Medicine and Therapeutics, Pavia, Italy; ⁸University of Ferrara, Department of Medical Sciences, Cona, Italy; ⁹Central Hospital of Bolzano, Rheumatology Unit, Bolzano, Italy; ¹⁰Ospedale Nuovo Regina Margherita, Dipartimento delle Specialità Mediche, Rome, Italy; ¹¹Ospedale S. Spirito e Nuovo Regina Margherita, U.O.C. Radiologia, Rome, Italy; ¹²AOU Città della Salute e della Sanità di Torino, Rheumatology Unit, Turin, Italy; ¹³Scientific Clinical Institutes Maugeri IRCCS, Department of General Medicine, Pavia, Italy; ¹⁴ASST Centro Ortopedico Traumatologico G. Pini-CTO, Dipartimento di reumatologia e scienze mediche, Milano, Italy; ¹⁵Ospedale Sacro Cuore di Negrar, UOS di Reumatologia, Negrar di Valpolicella, Italy; ¹⁶Academic Rheumatology Centre,

Dipartimento Scienze Cliniche e Biologiche, Turin, Italy; ¹⁷University of Verona, Department of Medicine, Verona, Italy; ¹⁸Padua, Department of Medicine DIMED, Padua, Italy; ¹⁹Rheumatology Unit, Medical Sciences and Public Health, University of Cagliari, Cagliari, Italy

Background: The UPSTREAM (NCT03330769) is a 24-month multi-center prospective cohort study that primarily aims to evaluate the additional value of musculoskeletal ultrasound (msk-US) over clinical examination in predicting 6-month minimal disease activity in Psoriatic Arthritis (PsA). (1)

Objectives: To develop and preliminarily validate an activity msk-US score and a damage msk-US score for PsA using the UPSTREAM database.

Methods: Patients classified with PsA according to CASPAR criteria and starting a new course of therapy for clinically active peripheral joint disease were eligible. The information regarding objectives, study design, clinical and US assessment has already been published (1). The msk-US examination was performed in 42 joints, 36 tendons, 12 entheses and 2 bursae defined through a web-based exercise (2). The sonographic elementary lesions were allocated to disease activity [i.e. synovitis (sy), tenosynovitis (ts), peritendinitis (pt), bursitis (bs) all evaluated both in Grey Scale (GS) and Power Doppler (PD) and active enthesitis (en)] and to damage (i.e. joint erosion, bone proliferation, tendon tear, enthesophyte, calcification and irregular entheses bone profile). Hands and feet X-ray were assessed using the modified Sharp-Van der Heijde (mSVH) score. A principal component (PC) analysis (PCA) was performed for each score and the number of PCs was defined by means of parallel analysis using baseline data. Each PC was normalized (n) taking into account the proportion between the observed value (e.g. sy-GS count) and the maximum expected value (e.g. 42 for sy-GS). Spearman' correlation was used to investigate the construct and discrimination validity of the new scores.

Results: Between February 2017 and May 2020, 312 PsA patients (155 men), with a mean (SD) age of 52.8 13.4, were enrolled from 19 centers; 22 expert sonographers were involved with substantial agreement for US lesions evaluated ($k \geq 0.7$). The median [IQR] disease duration was 1.3 [0.1-6.1] years and the median [IQR] tender joint and swollen joint counts were 6 [3-13] and 2 [1-5], respectively. The weight derived from PCA for each sonographic lesions and the final equation for calculating the scores are reported in Figure 1 (1A activity and 1B damage). The final msk-US activity score [$n(ts-GS + ts-PD)*2.87 + n(bs-GS + bs-PD)*1.76 + n(pt-GS + pt-PD)*1.43 + n(active en)*1.00 + n(sy-GS)*0.83 + n(sy-PD)*0.45$] has the best construct and discrimination validities according to a significant correlation with all clinical variables usually related to clinical activity (Table 1). The msk-US damage score correlated with mSVH score, HAQ and other clinical variables (Table 1).

Table 1.

Variables	Msk-US activity score		Msk-US damage score	
	Spearman correlation	P-value	Spearman correlation	P-value
ESR	0.196	0.002	0.075	0.235
CRP	0.209	<0.001	0.068	0.254
TJC	0.338	<0.001	0.286	<0.001
SJC	0.338	<0.001	0.072	0.221
Dactylitis count	0.284	<0.001	-0.061	0.306
LEI	0.194	0.001	0.214	<0.001
Physician GA	0.15	0.012	0.016	0.793
Patient GA activity	0.138	0.018	-0.073	0.221
Patient GA pain	0.199	0.001	-0.027	0.648
HAQ	0.238	<0.001	0.146	0.014
BASDAI	0.237	<0.001	0.175	0.003
PSAID-9	0.7	0.004	0.148	0.013
DAPSA	0.392	<0.001	0.228	<0.001
Sharp van Der Heijde score	0.115	0.2	0.266	0.003

Figure 1.