healthcare professionals performing surgical procedures on the joints of COVID-19 patients are exposed to a risk of contagion due to exposure to respiratory droplets, blood and body fluids, but not to direct exposure to joint- or bone-related tissues. Furthermore, given that some case reports of arthritis in COVID-19 patients⁵⁻⁸ show that it is possible that COVID-19 patients display viral-mediated arthralgias and arthritis, the absence of the virus in the knee highlighted by our study suggests that it is unlikely that SARS-CoV-2 has a direct inflammatory action on the joint, but it could induce an inflammation-related reaction, manifesting as a reactive arthritis⁹.

REFERENCES:

- [1] Cheng ZJ, Shan J. 2019 Novel coronavirus: where we are and what we know. *Infection*. 2020;(0123456789):1-9. doi:10.1007/s15010-020-01401-y
- [2] Chen Y, Chen L, Deng Q, et al. The Presence of SARS-CoV-2 RNA in Feces of COVID-19 Patients. *J Med Virol*. April 2020. doi:10.1002/jmv.25825
- [3] Maiese A, Manetti AC, La Russa R, et al. Autopsy findings in COVID-19-related deaths: a literature review. *Forensic Sci Med Pathol*. 2020. doi:10.1007/s12024-020-00310-8
- [4] Baldanti F, Novazzi F, Cassaniti I, Piralla A, Di A, Bruno R. Detection of the SARS-CoV-2 in different biologic specimens from positive patients with COVID-19, in Northern Italy. *Authorea*. 2020. doi:10.22541/au.159724509.93056098
- [5] López-González M-C, Peral-Garrido ML, Calabuig I, et al. Case series of acute arthritis during COVID-19 admission. *Ann Rheum Dis*. 2020;0(0):annrheumdis-2020-217914. doi:10.1136/annrheumdis-2020-217914
- [6] Parisi S, Borrelli R, Bianchi S, Fusaro E. Viral arthritis and COVID-19. *Lancet Rheumatol*. 2020;2(11):e655-e657. doi:10.1016/S2665-9913(20)30348-9
- [7] Alivernini S, Cingolani A, Gessi M, et al. Comparative analysis of synovial inflammation after SARS-CoV-2 infection. *Ann Rheum Dis*. 2020;0(0):2-3. doi:10.1136/annrheumdis-2020-218315
- [8] Talarico R, Stagnaro C, Ferro F, Carli L, Mosca M. Symmetric peripheral polyarthritis developed during SARS-CoV-2 infection. *Lancet Rheumatol*. 2020;2(9):e518-e519. doi:10.1016/S2665-9913(20)30216-2
- [9] Di Carlo M, Tardella M, Salaffi F. Can SARS-CoV-2 induce reactive arthritis? *Clin Exp Rheumatol*. 2020;In press.

Acknowledgements: We would like to thank the clinical staff of the Pathology Unit for their collaboration in performing the autopsies.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2033

AB0672

PSYCHOLOGIC STATE OF PATIENTS WITH RHEUMATIC DISEASES AFTER THE OUTBREAK OF SARS-COV2

M. Yasmine¹, S. Mariem¹, S. Miladi¹, A. Fazaa¹, L. Souebni¹, K. Ouenniche¹, S. Kassab¹, S. Chekili¹, K. Ben Abdelghani¹, A. Laatar¹. ¹Mongi Slim Hospital, Rheumatology, Tunis, Tunisia

Background: The ongoing Coronavirus-19 disease epidemic is a public health emergency of international concern which could pose a challenge to psychological resilience. This is mainly the case of patients with rheumatic diseases (RD).

Objectives: The aim of this study was to establish the prevalence of psychiatric symptoms in patients with RD and to identify factors associated with psychological stress.

Methods: A cross-sectional study was conducted among patients with RD. They were invited to respond to a survey designed to assess their psychological response after the outbreak of the SARS-COV2 pandemic. Mental health status after the Covid-19 was evaluated using the Depression and Anxiety Scale (DASS-21). The total depression subscale score was divided into normal (0–9), mild depression (10–12), moderate depression (13–20), severe depression (21–27), and extremely severe depression (28–42). The total anxiety subscale score was divided into normal (0–6), mild anxiety (7–9), moderate anxiety (10–14), severe anxiety (15–19), and extremely severe anxiety (20–42). The Impact of Event Scale-Revised (IES-R) is a 22-item self-administered questionnaire that assesses the post-traumatic state (PTS). A cut-off point of 33 provided the best diagnostic accuracy for PTS. The level of significance was fixed at 0.05.

Results: Sixty patients were included (42 with rheumatoid arthritis and 18 with spondylarthritis) with a mean age of 51.3 ± 13.1 years [18–75]. The majority of patients were women (63.3%), married (81.7%), with a household size of 3–5 people (67.9%). The mean disease duration was 11.9 ± 5.6 [1–23] years.

According to DASS depression score almost half of the patients (47%) suffered from depression, which was classified as severe in 10% of cases. The mean DASS anxiety subscale was 12.6 ± 8.2 [1–35]. The level of anxiety was mild in 23% of cases, moderate in 25% of cases, and severe in 23.3% of cases. The mean ILES-R score was 20.4 ± 10.2 [20–38]. Women experienced more PTS than men (26.3% vs 22.7%) without reaching a statistically significant correlation ($p=0.9$). Patients with higher education had lower IES-R, depression, and anxiety subscale scores as compared to those who were illiterate without a statistically

significant correlation ($p=0.5$, $p=0.2$, $p=0.4$ respectively). A positive correlation was found between anxiety score and IES-R ($p=0.00$). Other sociodemographic variables including disease duration, parental status, marital status, and household size were not associated with IES-R and DASS subscale scores ($p>0.05$).

Conclusion: Our study showed that many of our patients with RD experienced anxiety, depression, and PTS after the outbreak of Covid-19. Professional status as well as other sociodemographic data was not associated with psychological stress.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2116

AB0673

THE ROUTINE HEALTH CARE OF PATIENTS WITH RHEUMATIC DISEASES DURING THE PANDEMIC OF COVID-19

S. Mariem¹, M. Yasmine¹, S. Miladi¹, A. Fazaa¹, L. Souebni¹, K. Ouenniche¹, S. Kassab¹, S. Chekili¹, K. Ben Abdelghani¹, A. Laatar¹. ¹Mongi Slim Hospital, Rheumatology, Tunis, Tunisia

Background: The coronavirus pandemic poses a great threat to patients with rheumatic diseases (RD). Because of the state of immunodepression, they are perceived as vulnerable and at high risk of contracting the virus. This may lead to a delay in the health care routine.

Objectives: The aim of our study was to assess the routine of patients with rheumatic diseases in the era of covid-19.

Methods: We conducted a prospective study after the outbreak of Covid-19 including patients with RD: rheumatoid arthritis (RA) and Spondylarthritis (SpA). All of them were invited to answer a questionnaire about adherence to protective measures, health services, hospital visits, treatment modalities, as well as the level of satisfaction with the medical support or information received for COVID-19. We also evaluated the post-traumatic state (PTS) with the Impact of Event Scale-Revised (IES-R). A cut-off point fixed at 33 provided the best diagnostic accuracy for PTS. The level of significance was fixed at 0.05.

Results: The study included 60 patients with a female predominance (sex ratio=1.7). The mean age was 51.3 ± 13.1 years [18–75] and the mean disease duration was 11.9 ± 5.6 [1–23] years. Forty-two patients had RA and 18 patients had SpA. The majority of respondents were women (63.3%), married (81.7%), with a household size of 3–5 people (67.9%). The mean DASS depression and anxiety subscale was 9.5 ± 7.6 [2–32] and 12.6 ± 8.2 [1–35] respectively. PTS was found in 25% of cases. Twenty-two patients stopped working in the second outbreak. The adherence to protective measures was as follows: compliance with lock-down (92.5%), adherence to stay home warnings (82%), wearing masks (96.3%), social distancing (68%), and washing hands immediately after sneezing (59.3%). Information was mainly sourced from both social media and the Internet in 78.2% of cases. Most of the patients were satisfied with the information provided and only 5 demanded more information about vaccination protocol. Of the patients having direct contact with a person affected with Covid-19 (20%), seven patients experienced symptoms. Three patients were diagnosed with SARS-COV2 and three other patients were hospitalized because of flares. The majority of the respondents (98.2%) had confidence in their own doctors' ability to recognize Covid-19. The majority of patients (91%) were afraid of getting sick and 83.1% feared stigma and discrimination after the infection. Fifty patients did attend their consultation appointment and 13.3% had a phone consultation. About a third of patients (26.7%) of patients experienced a delay in treatment. A positive correlation was found between marital status and the fear of other family members getting Covid-19 ($p=0.029$). Patients who did not consult experienced more PTS and severe anxiety than those who did not delay hospital visits ($p=0.07$, $p=0.04$ respectively).

Conclusion: Our study showed that during the outbreak of Covid-19, patients with RD exhibited a high level of protection towards the virus. Despite these precautionary measures, a delay in hospital routine was observed.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2125

AB0674

RAPID ADOPTION OF TELEMEDICINE IN RHEUMATOLOGY TRAINING: RESULTS FROM THE COVID-19 GLOBAL RHEUMATOLOGY ALLIANCE TRAINEE SURVEY

S.A. Yeoh¹, K. Young², M. Putman³, E. Graef⁴, F. Berenbaum^{5,6,7}, R. Conway⁸, R. Grainger⁹, A. Kilian¹⁰, M. König¹¹, J. Liew¹², P. M. Machado¹, S. E. Sattui¹³, J. Sparks^{14,15}, P. Sufka¹⁶, M. Ugarte-Gil¹⁷, L. Upton¹⁸, Z. Wallace^{15,19}, J. Yazdany²⁰, A. Jayatilake²¹ on behalf of the COVID-19 Global Rheumatology Alliance. ¹University College London, Rheumatology, London, United Kingdom; ²University of Texas Southwest Medical Center, Rheumatology, Dallas, United States of America; ³Medical College of Wisconsin, Rheumatology, Milwaukee, United States of America; ⁴Emerson Hospital, Department of Medicine,