disease (n = 1), polyosteoarthritis (n = 6), benign osteopathy (n = 1). 17 patients were in low disease activity when they presented the COVID-19. 8 patients presented moderate to high activity, of which 2 (25%) developed a form requiring hospitalization. 7 patients were under NSAIDs, 13 under corticosteroids, 19 under csDMARD, 7 under bDMARD. The COVID-19 disease was symptomatic in 83% of cases, with fatigue (68%), cough (52%), back pain (48%), anosmia (48%), ageusia (44%), arthritis (44%), fever (36%). The diagnosis was made with polymerase chain reaction (83%), serology (10%), CT scan (23%). The CT lung lesions reached 25-50% in 2 lupus patients with a moderate disease activity, under corticosteroids and hydroxychloroquine. The involvement was > 75% in 2 patients, who admitted to intensive care: one was followed for polyarthrosis and was hypertensive and poorly balanced diabetic, and the other was followed for Horton's disease. The treatment was administered on an outpatient basis in 77.4% of cases, in an intensive care unit in 16% of cases. 30 patients fully recovered from infection, one patient died secondary to pulmonary embolism. The subgroup of patients on immunosuppressants consisted of 23 patients (76%), of whom 5 (22%) developed a moderate to severe form of COVID-19 requiring hospitalization. Among these hospitalized patients, 2 (40%) had moderate to high disease activity of their rheumatism.

Conclusion: Most of the patients in our series presented a mild form of COVID-19, including those under immunosuppressants. Patients who were under hydroxychloroquine were not spared from COVID-19, but developed a mild form. The disease activity of the chronic rheumatism did not appear to influence the severity of the COVID-19.

REFERENCES:

- [1] Grange L, Guilpain P, Truchetet M-E, Cracowski J-L. Challenges of autoimmune rheumatic disease treatment during the COVID-19 pandemic: A review. *Therapies*. 2020;75(4):335-42.
- [2] Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *Jama*. 2020;323(13):1239-42.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2744

AB0686 IMPACT OF COVID19 IN SPONDYLOARTHRITIS (SPA) PATIENTS IN A TERTIARY HOSPITAL

C. L. Garcia Gomez¹, D. Montero¹, L. Vega¹, O. Ibaranguoitia², I. Calvo², C. E. Perez¹, J. M. Blanco¹, E. Galindez¹, A. R. Inchaurre¹, M. E. Ruiz¹, I. Torre¹, E. Cuande¹, I. Gorostiza³, M. R. Exposito-Moliner¹, M. L. Garcia Vilar¹.

¹University Hospital of Basurto, Rheumatology, Bilbao, Spain; ²Hospital of Galdakao, Rheumatology, Galdakao, Spain; ³University Hospital of Basurto, Investigation, Bilbao, Spain

Background: This tertiary hospital is the referral centre of 360.000 inhabitants, population with a Covid seroprevalence of 8,4% at final 2020.

Since march, we have had a special concern for rheumatologic patients with systemic diseases and under immunosuppressive agents, including disease modifying antirheumatic drugs (DMARDs) and biological therapy (BT). This is why a special protocol for this population was set. It included performance of serology (CLIA test) for patients under BT and PCR and CLIA testing prior to new treatments. PCR testing was also generally performed: if symptomatology consistent with Covid; before hospitalisation; to tight contacts of infected people; and before procedures.

Objectives: To evaluate the impact of COVID-19 in our SpA patients in terms of severity of viral infection and its effect on SpA.

Methods: Data of 665 SpA patients and confirmed Covid infection seen in our center from March 15th to December 15th was crossed. 3 miscoded patients with rheumatoid arthritis and 2 with non definite CLIA positivity were excluded. Finally 49 patients' clinical records were reviewed. Data regarding epidemiologic features, SpA characteristics, comorbidities, therapy received, clinical activity before and after Covid, and severity of the infection was collected. IBM SPSS v23 was used for statistical analysis.

Results: Among 49 SpA patients, 59% were male, mean aged 56,63 years (range 23-79). 62,2% presented at least 1 comorbidity. 65% were psoriatic arthritis. They mostly had longstanding disease (median 10,5 years -range- 1-35). Previously 63% had received DMARDs, mainly methotrexate, and 32 % BT.

When Covid was diagnosed 37,2% were under DMARDs and 53% under BT (69,2% TNF inhibitors, 26,9% anti-IL 17, 3,9% ustekinumab). At this point, disease activity was controlled in 82% of patients (39% in remission, and 43% in low disease activity state). Only 18% showed moderate activity.

Within the 49 patients, 34 were diagnosed by PCR and 15 by CLIA tests. 9 required hospitalisation, of whom 4 developed more severe disease (3 received glucocorticoid pulses and 2 tocilizumab). A woman with PsA under secukinumab presented pneumonia and PE. None required mechanical ventilation. There were no exits.

Due to Covid infection 9 patients (50%) stopped DMARDs treatment, (5 of them hospitalised). 9 patients withdrew BT after Covid diagnosis; 60% of the BT-hospitalised, and 28,5% of the BT- non-hospitalised. 1 suffered a flow with severe disease activity after withdrawal of IL-17 inhibitor.

Conclusion: Prevalence of SARS cov 2 infection in SpA patients was not greater than in general population. Most were asymptomatic or suffered mild disease. Only 9 were hospitalised. Factors related to hospitalisation seem similar to those of general population, even if statistical significance was not found due to the small sample. BT does not seem to relate to hospitalisation in SpA and we had no deaths to date in them.

More studies should be made to throw out conclusions.

Baseline characteristics of SpA patients with confirmed Covid19	Hospitalised (N 9)	Non-hospitalised (N 40)	Total (N 49)
Mean age (range) – yr	62,56 (43-74)	55,3 (23-79)	56,63 (23-79)
Male sex - no. (%)	6 (66,6)	23 (57,5)	29 (59,1)
Smoking habit (active, ex-smokers) - no. (%)	2 (22,2)	23 (57,5)	25 (51)
Non comorbidities - no. (%)	2 (22,2)	17 (42,5)	19 (38,8)
Hypertension - no. (%)	3 (33)	14 (35)	17 (35)
Diabetes - no. (%)	1 (11)	6 (15)	7 (14,2)
Dyslipidemia - no. (%)	4 (44)	9 (22,5)	13 (26,5)
Obesity (BMI >30) - no. (%)	4 (44)	12 (30)	16 (32,7)
Chronic obstructive pulmonary disease - no. (%)	2 (22,2)	4 (10)	6 (12,2)
Asthma - no. (%)	0 (0)	4 (10)	4 (8,2)
Cardiopathy - no. (%)	1 (11,1)	8 (20)	9 (18,4)
Median disease time since diagnosis (range) – yr	14, 5 (7-20)	10 (1-35)	10,5 (1-35)
SpA type			
Psoriatic arthritis - no. (%)	8 (88,8)	24 (60)	32 (65,3)
Ankylosing spondylitis - no. (%)	1 (11,1)	15 (37,5)	16 (32,7)
Non-Rx ankylosing spondylitis - no. (%)	0 (0)	1 (2,5)	1 (2)
Current treatment when diagnosed (N 48)			
DMARDs - no. (%)	6 (66,6)	12 (30)	18 (37,5)
BT - no. (%)	5 (55,5)*	21 (52,5)**	26 (54,1)
* 5 anti-TNF (1 etanercept), 2 anti IL-17, 1 anti IL-12-23.			
** 12 anti-TNF (4 etanercept), 5 anti IL-17.			

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2960

AB0687 EFFECT OF COVID-19 ON PATIENTS ATTENDING AMBULATORY DEEP VENOUS THROMBOSIS OUT-PATIENT CLINIC: A RETROSPECTIVE, COHORT ANALYSIS[K11] [K11]101 CHARACTERS

T. Hill¹, K. P. Iyengar¹, A. Nune². ¹Southport & Ormskirk Hospital NHS Trust, Trauma and Orthopaedics, Southport, United Kingdom; ²Southport & Ormskirk Hospital NHS Trust, Rheumatology, Southport, United Kingdom

Background: COVID-19 has been acknowledged as an procoagulant disorder with increased risk of venous thromboembolism (VTE) [1]. High rates of VTE in patients admitted to intensive care unit have been reported [2]. However effect of COVID-19 on ambulatory patients attending out-patient clinics has not been described.

Objectives: This retrospective, observational cohort study analyses the impact of COVID-19 on incidence of Deep Venous Thrombosis (DVT) in patients referred to Ambulatory Deep Venous Thrombosis Out-patient Clinic (ADOC). A comparative analysis of patients attending ADOC during a similar period in 2019 is undertaken.

Methods: Patients who attended the ADOC with suspected DVT during the 'first wave' of the COVID-19 pandemic between 01 February and 30 June 2020 at Southport and Ormskirk NHS Trust were studied and compared to a similar period in 2019. Patients characteristics, comorbidities, risk factors, incidence of DVT and relationship with COVID-19 with the evaluation of haematological parameters including D-dimer and two- tier Wells score. Additionally we examined patients outcomes including morbidity, mortality and hospital admissions.

Results: Overall, there was a decrease in the number of patients attending the ADOC from 290 in 2019 to 233 in 2020. However, a total of 38 patients tested positive for DVT, a rate of 16.3%, which is compared to an incidence of 7.9% in 2019. Due to evolving protocols, the COVID-19 status of all patients attending the ADOC could not be ascertained. However, 5/233 of the 2020 cohort patients either had a positive test result (n=4) or were symptomatic of COVID-19 (n=1).

Conclusion: There has been a 129% increase in the incidence of DVT in patients presenting to ADOC at our trust during the 'first wave' of the COVID-19