that telephone consultation should be on demand and by telephone, 25 (78.1%) considered it necessary to include video calls and the possibility of receiving reports in non-face to face care and photos. Finally, 23(71.9%) centers recorded nursing and medical telephone consultations in their medical records.

Conclusion: The pandemic had a huge impact on nursing care for rheumatic patients, with difficulties associated with extra work load and changes in the dynamics of care. A readjustment of assistance has been necessary and audio-visual aids were necessary to improve telephone (non face to face) service.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1923

AB0905-HPR TELEMEDICINE IN RHEUMATOLOGY AT TIME OF COVID PANDEMIC

G. Cuomo¹, F. Masini², K. Gjeshi², D. Lubrano², L. E. Adinolfi², C. Romano².

¹University of Campania Studies "Luigi Vanvitelli", Precision of Medicine, Napoli, Italy; ²University of Campania Studies "Luigi Vanvitelli", Department of Medical and Surgical Sciences, Napoli, Italy

Background: The outbreak of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic has imposed considerable restrictions on people's mobility in order to limit infection transmission. Before the COVID -19 pandemic tele-rheumatology was proposed to patients living in remote areas. Currently, the use of telemedicine has increased significantly. We report on the implementation of a telemedicine program for the evaluation and treatment of patients with rheumatic diseases (1).

Objectives: The aim of the study is to evaluate telemedicine as a viable approach for routine follow-up and management of rheumatic disease.

Methods: Sixty-six patients were evaluated online by the remote rheumatologist in four weeks. The population of rheumatology patients was evaluated using a IARPLUS platform Information collected included demographic information consisting of age, gender, and primary rheumatologic diagnosis. Results: The average age of patients who were seen was 54 years and 85% of patients were women.

The most common disorders included rheumatoid arthritis (22, 33%), axial spondyloarthropathies (7, 11%), and psoriatic arthritis (15,23%), systemic sclerosis (13, 20%), lupus (6, 9%), UCTD (2,3%), Sjogren and fibromyalgia (1, 1.5%).

Results: All patients were given recommendations on COVID-19 vaccine administration. Therapy was remodeled in 13 patients (in particular in 5 patients with anemia intravenous iron infusion was scheduled; 1 started immunosuppressant therapy for proteinuria, 4 increased methotrexate for disease activity, 2 discontinued Methotrexate for adverse events (hypertransaminasemia), 1 suspended OH-chloroquine due to retinal accumulation; 2 biologic-naïve patients, after a three-month of waiting due to inability to come to the hospital, started the biologic drug; 22 patients received a renewal of the therapeutic plan; 25 patients had a regular six-month follow up; 2 ticket exemption for illness; 13 consultations and/or laboratory tests (1 pneumological consultation, 1 ophthalmological consultation, 1 request for sacro-iliac MRI, 3 nailfold videocapillaroscopies, 3 FKT, 1 musculotendinous ultrasound, 1 antibodies for celiac disease, 2 antibodies anti Sars-CoV-2).

Conclusion: Telemedicine is becoming more prevalent. We report the successful use of this service in evaluation and management of rheumatic diseases in a period with limited access to rheumatologic care. We have shown that patients can be seen, evaluated, and successfully treated with a variety of medications, including biologic agents, and evaluated for both chronic inflammatory arthropathies and connective tissue diseases.

REFERENCES:

- [1] Rheumatology Care Using Telemedicine Michael M. Rezaian; Telemedicine and e-Health, 2020.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2719

AB0906-HPR BENEFIT OF A NURSE-LED PROGRAM OF CARE FOR MANAGEMENT OF PATIENTS WITH ESTABLISHED RHEUMATOID ARTHRITIS: SYSTEMATIC LITERATURE REVIEW

P. Lois¹, L. Lopez Pedraza¹, G. Candelas², C. Lajas², M. Rodero², T. Mulero López², L. León¹, B. Fernandez¹, L. Rodriguez Rodriguez¹, L. Abasolo¹ on behalf of Departamento de Investigación de la Unidad de Gestión Clínica de Reumatología del Hospital Clínico San Carlos. ¹Hospital Clínico San Carlos, Instituto de Investigación Sanitaria del Hospital Clínico San Carlos (IdISSC), Reumatología, Madrid, Spain; ²Hospital Clínico San Carlos, Reumatología, Madrid, Spain

Background: Nowadays, many countries experience an urgent demand for rheumatologists and greater cost-effectiveness. Thus, different alternatives to conventional outpatient physician-led follow-up have been investigated. The

alternative of community-based service delivery systems is intended to meet emerging needs, reducing the costs of hospital care and morbidity. In patients with long-term conditions, such as Rheumatoid Arthritis (RA), nurses are seen as essential in the delivery of promotion in self-care and decision-making, favoring the empowerment of patients.

Objectives: To systematically investigate if the nurse led-care management (NLCM) education programs are beneficial in patients with RA.

Methods: This is a systematic literature research publication on the following bibliographic databases: Medline, Embase and the Cochrane Central register of Controlled Trials from the beginning until 2019.

Results: From a yield of 116 studies, 44 were included in the review. After quality assessment and exclusion of the lowest-ranked studies, 19 studies were analyzed. Outcomes were organized into six themes (1) disease activity; (2) adherence; (3)satisfaction; (4)treatment compliance, (5) depression and (6)quality of life. All studies had significant results in at least one of the outcomes.

Conclusion: According to our results, we can conclude that, clinicians could benefit from support conducted by nurse in the management of these patients in tasks such as metrology clinic, participation in enforcement, self-administration, the correct dosage and side effects of treatments, and link the patient and other professionals or entities. It would therefore be advisable to conduct further studies to draw more accurate and reliable conclusions in this context.

REFERENCES:

- [1] Bech B, Primdahl J, Van Tubergen A, et al. 2018 update of the EULAR recommendations for the role of the nurse in the management of chronic inflammatory arthritis. *Ann Rheum Dis* 2020; 79: 61–68.
- [2] Uthman I, Almoallim H, Buckley CD, et al. Nurse-led care for the management of rheumatoid arthritis: a review of the global literature and proposed strategies for implementation in Africa and the Middle East. *Rheumatology International* 2020; 1: 3.
- [3] Garner S, Lopatina E, Rankin JA, et al. Nurse-led care for patients with rheumatoid arthritis: A systematic review of the effect on quality of care. *J Rheumatol* 2017; 44: 757–765.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.3580

AB0907-HPR UNEXPECTED DIFFERENCE IN ACCEPTANCE OF TELECONSULTATION BETWEEN PATIENTS WITH LUPUS AND RHEUMATOID ARTHRITIS WHO UNDERWENT TO A DEVELOPED AND IMPLEMENTED TELEMEDICINE INNOVATIVE PROGRAM AFTER THE DECLARATION OF QUARANTINE DUE TO THE COVID-19 PANDEMIC IN COLOMBIA

S. M. Hernández-Zambrano¹, R. A. Castiblanco-Montañez¹, J. Chavez-Chavez², D. P. Rivera-Triana², A. Aza³, L. Villarreal³, M. Martinez⁴, A. Rojas-Villarraga², P. Santos-Moreno⁵. ¹Fundación Universitaria de Ciencias de la Salud - FUCS, Nurse Department, Bogota, Colombia; ²Fundación Universitaria de Ciencias de la Salud - FUCS, Research Division, Bogota, Colombia; ³Biomab - Center for Rheumatoid Arthritis, Bogota, Colombia, Administration, Bogota, Colombia; ⁴Biomab - Center for Rheumatoid Arthritis, Bogota, Colombia, General Medicine, Bogota, Colombia; ⁵Biomab - Center for Rheumatoid Arthritis, Bogota, Colombia, Rheumatology, Bogota, Colombia

Background: The Covid-19 pandemic has generated restrictions in the mobility of people, affecting the face-to-face care of patients with chronic diseases, including autoimmune. The health emergency has created the need to establish follow-up alternatives, giving rise to telemedicine.

Objectives: To evaluate the level of attendance to teleconsultation and the face-to-face usual care in patients with rheumatoid arthritis (RA) and systemic lupus erythematosus (SLE) who underwent to a developed innovative telemedicine program after the declaration of quarantine due to the Covid-19 pandemic.

Methods: The present cross-sectional descriptive study reports retrospective data collected from patients with RA and SLE from March to June, 2020;. Since the beginning of March 2020 when Covid-19 was declared in Colombia as a health emergency establishing specific standards for outpatient care, our center has made the necessary legal and technical adjustments to develop an innovative telemedicine service to prevent the epidemiological risk. To all contacted patients were offered the option of teleconsultation or face-to-face consultation; a standardized protocol was set with clinimetry measures evaluated in both groups. Continuous variables were described using mean and standard deviation, and categorical variables were described using numbers and percentages. We performed chi-square tests of independence to determine differences between teleconsulting and conventional face-to-face consultation.

Results: A total of 5745 RA patients were followed-up, 5292 (92.1%) by teleconsulting and 453 (7.9%) by conventional face-to-face consultation; among the group of SLE patients, a total of 646 were assessed, of which there were 386 (60%) by teleconsultation and 260 (40%) by face-to-face consultation; this highlights an important difference in the level of acceptance of the teleconsultation