valve involvement (5.10%) and epididymitis (9.95%). Among female patients, the frequency of genital ulcers is significantly higher than that of men (86.24% vs. 68.69%, $P<0.001$).

Conclusion: We described the clinical characteristics of a large cohort of pediatric BD patients in China. While the clinical manifestations are largely similar to adult BD, the time from symptom onset to diagnosis is significantly delayed in pediatric patients. Our study highlights the need to improve awareness of BD among pediatric providers to facilitate rapid diagnosis and treatment in children.

REFERENCES:

- [1] Yıldız et al. Pediatric Behçet's disease- clinical aspects and current concepts. *Eur J Rheumatol* 2020; 7(Suppl 1): S38-S47.
- [2] Koné-Paut. Behçet's disease in children, an overview. *Pediatric Rheumatology* (2016) 14:10.

Table 1. Comparison of clinical features of BD in children and adults

	Children (n=848)	Adult (n=1531)	P
Age of first symptom (y)	11.59±4.11	27.08±7.91	
Sex(M/F)	412/436		
Years of diagnosis (y)	13.34±9.13	6.73±33.21	<0.001
Recurrent oral aphthosis	848 (100%)	1518 (99.15%)	0.007
Genital Ulcers	659 (77.71%)189	1152 (75.24%)379	0.176
Ocular signs	256 (30.19%)592	508 (33.18%)1023	0.134
Cutaneous signs	539 (63.56%)309	1020 (66.62%)511	0.132
Neurological signs	39 (4.60%)809	82 (5.36%)1449	0.421
Gastrointestinal signs	170 (20.05%)678	244 (15.94%)1387	0.001
Vascular signs	83 (9.79%)765	162 (10.58%)1369	0.542
Heart valve signs	31 (3.66%)817	110 (7.18%)1421	<0.001
Joint signs	275 (32.43%)573	612 (39.97%)919	<0.001
Fever	3 (0.35%)845	22 (1.44%)1509	0.013
Positive pathergy test	179 (21.11%)669	347 (22.66%)1184	0.381

Acknowledgements: China Behçet' Disease Patients Association.

Disclosure of Interests: None declared.

DOI: 10.1136/annrheumdis-2021-eular.3621

POS0082

PERFORMANCES OF THE RISK FACTORS AND PEDIATRIC VASCULITIS ACTIVITY SCORES FOR DISEASE SEVERITY IN CHILDREN WITH MULTISYSTEM INFLAMMATORY SYNDROME

B. Sözeri¹, Ş. Çağlayan¹, A. Akyol², K. Ulu¹, T. Coşkun¹, C. Hasbal Akkuş², F. Demir¹. ¹Health Sciences University, Umraniye Training and Research Hospital, Pediatric rheumatology, Istanbul, Turkey; ²Health Sciences University, Umraniye Training and Research Hospital, Pediatrics, Istanbul, Turkey

Background: Multisystem Inflammatory Syndrome in Children (MIS-C) is observed by hyperinflammation and cytokine storm. The spectrum of severity ranged from standard hospitalization to pediatric intensive care unit management. There is no specific activity score that predicts whether this hyperinflammatory state will be severe or result in mortality in pediatric patients. There are activity scores used in KD and other vasculitis such as Kobayashi score (KS) and Pediatric Vasculitis Activity Score (PVAS) that determine the severity of the disease in children.

Objectives: Based on the clinical similarity of MIS-C to these disease groups, we wanted to evaluate the performance of these disease activity scores. Also, we aimed to identify the factors associated with the disease severity of patients with MISC.

Methods: We retrospectively enrolled a single-center cohort of 45 consecutive pediatric patients with MISC admitted to Umraniye Training and Research Hospital, Pediatric Rheumatology Clinic, Istanbul, Turkey, from April 20 to December 31, 2020. Medical information of each patient including demographic data, clinical characteristics, laboratory results, and outcomes was extracted retrospectively through review of electronic medical records. We analyzed all score systems including KS, PVAS, NLR, cHIS, and C-reactive protein/albumin ratio (CAR) as assessment factors for diagnosis for severe disease and evaluation of disease activity in MISC. All scores were compared between two groups and receiver operating characteristic (ROC) curve analysis was performed to evaluate diagnostic utility.

Results: We reported 45 patients (10 female, 35male) with MISC. Their mean age was 9.65±4.93 years (7 months-18 years). All patients had fever (median 4 days), 71 % patients had acute gastrointestinal symptoms, 37.8 % of patient's conjunctivitis and only 5 patients had respiratory findings at admission. Twenty-four (46.7%) patients met criteria for classic KD. Macrophage activation syndrome and myocardial dysfunction with or without cardiogenic shock were seen 14 and 10 patients respectively. All the patients had positive serology for SARS-CoV-2, abnormal complete blood counts and coagulation tests, and elevated inflammatory markers.

We divided the disease severity into a moderate or severe group based on admission on intensive care unit (ICU). There were 15 patients with severe illness

(33%). The median age of these patients was significantly older (11.3 years vs 9.16 years, $p=0.05$). The median hospital stay period was 10 days. The median need for intensive care was on the first day (1-14th days), and the median lasted 5 (1-9) days. The majority of MISC patients were on Intravenous immunoglobulin (IVIg) (89%), and corticosteroid (73.3%). A total of 12 patients received anakinra. In the severe group, all patients had higher values of KD, PVAS, NLR, cHIS, and CAR than the patients in moderate group. For severe MISC, the area under receiver operating characteristic curve (AUC) was 0.864 (95% confidence interval [CI], 0.729–1) for the PVAS, 0.911 (95% CI, 0.827–0.995) for the NLR, and 0.853 (95% CI, 0.744–0.963) for the CAR, with optimal cut-off values of 3.5, 9.05, and 4.86, respectively. Thirty-eight (84.4%) of the 45 patients met two or more cHIS criteria at the time of their hospitalization; 39% of these patients were identified as severe group (OR 1.62, 95% CI 1.27-2.13, $p=0.04$). At the time of diagnosis, 29 patients with a Kobayashi score greater than 4 were detected, of which 15 required intensive care (OR 2.07, 95% CI 1.42-3.0, $p=0.00$).

Conclusion: This study demonstrated that both inflammatory scores (CAR and NLR) and disease activity scores (KS, PVAS and cHIS) can be used to aid the assessment for severity of MISC.

REFERENCES:

- [1] Webb BJ, et al *Lancet Rheumatol*. 2020 Dec;2(12):e754-e763.
- [2] Feldstein LR et al *N Engl J Med*. 2020;383(4):334-346

Disclosure of Interests: None declared.

DOI: 10.1136/annrheumdis-2021-eular.2916

POS0083

USEFUL EXPERIENCE OF RHEUMATOLOGISTS FOR UNDERSTANDING OF THE DISEASE ORIGIN AND APPROACHES TO THERAPY IN FIBRODYSPLASIA OSSIFICANS PROGRESSIVA

L. Nikishina¹, S. Arsenyeva¹, V. Matkava¹, A. Arefieva¹, M. Kaleda¹, T. Pachkoria¹, Z. Kolkhidova¹. ¹V.A. Nasonova Research Institute of Rheumatology, Paediatric, Moscow, Russian Federation

Background: Many monogenic genetic conditions, such as auto-inflammatory diseases (AIDs), have similar clinical manifestations and immunopathogenesis to "classic" rheumatic diseases (RD). Such cases may include Fibrodysplasia ossificans progressiva (FOP), an extremely rare genetic disease, which, according to our previous study and data from other authors¹, may represent an example of AID with catastrophic heterotopic ossification due to a mutation in the ACVR1 gene. It seems that the experience of rheumatologists, especially children's ones, will be useful in the treatment of FOP.

Objectives: To analyze the dynamics of clinical manifestations and to therapy approaches including target anti-inflammatory drug Tofacitinib (TOFA) in the one of the world's largest groups of patients (pts) with FOP.

Methods: The study was based on the analysis retrospective and prospective observation of the 35 pts (17 males and 18 females) with a verified diagnosis of FOP for the period from 1998 to 2020. In 9 pts with severe course of FOP TOFA administration were evaluated.

Results: In all 35 pts the diagnosis was verified by "classic" FOP phenotype: malformed great toes in 33 pts (94.3%); short malformed thumbs-8 (22.8%); peripheral osteochondromas-20 (57.1%); abnormalities of the cervical spine-32 (91.4%), multiple heterotopic ossifications-32 (91.4%). Genetic tests were done in 26, it confirmed mutation in the ACVR1 gene in 100%. Long term follow-up detected a lot of spondyloarthritis-like signs similar to the manifestation of RD: ankylosis of the facet joints and vertebral bodies (by the type of syndesmophytes) in most pts, sacroiliitis, confirmed by radiological methods (X-ray, CT, MRI), gradual ankylosis in the peripheral joints in 18 (56.4%), synovitis in large joints in 8 (25%) pts (knee and hip mostly). In 9 pts with the most difficult course with rapid progression of ossification due to continuous flares despite the NSAIDs and steroids intake, we tried to use TOFA after the approval of the local Ethic Committee. We use the similar dose to randomized trial for JIA (up to 5mg twice a day). The first patient was 16 y.o. at the time of TOFA administration in December 2019, the age of the other pts was from 2 to 12 y.o. By present time duration of TOFA therapy is from 6 to 15 mo. For the previous 6 months before TOFA initiation the number of flares was in average 8 per patient. After 6 months of TOFA treatment the number of new flares decreased to 0-1, except youngest patient of 2 y.o. in whom the number of flares decreased from 10 to 4 per the same period. In all 9 pts we minimize the dose or completely stop the steroids. New nodes formation stopped immediately in most pts and also the significant motion improvement of large (shoulder) joints were established. Drug tolerance was good in all pts, no AE were registered. But despite the good clinical effect without new heterotopic ossification in our first patient, we found continuous intraskeletal ossification between vertebral bodies, facet and sacroiliac joints in MRI.

Conclusion: We are confident that the processes of heterotopic ossification in FOP are very similar to new bone formation phenomenon in spondyloarthritis and reliable suppression of inflammation can interrupt the progression of the disease. We used similar justifications to our colleagues for the use of anti-cytokine drugs, but used a JAK-kinase inhibitor, it was extremely important the oral rout of drug administration and possibility to escape any injections in FOP. TOFA demonstrated positive effect