Most TE HZ infections were mild to moderate and non-serious; 6 were serious; 2 were recurrences. No visceral TE HZ occurred across the FIL RA program; there was 1 case each of genital, disseminated, and ophthalmic HZ. The disseminated HZ occurred in a pt with prior HZ history. Lymphopenia was not associated with HZ during the PBO-controlled W12 period.

Conclusion: HZ was more common in both FIL groups vs ADA or MTX up to 52 weeks but comparable vs PBO during the 12-week placebo-controlled period. In multivariate analyses, prior history of HZ, Asian region, and age ≥50 years were associated with increased HZ risk.

REFERENCES:

[1] Genovese et al. *JAMA*. 2019;322:315–25.
[2] Westhovens et al. *Ann Rheum Dis*. 2021; online first.
[3] Combe et al. *Ann Rheum Dis*. 2021; online first.
[4] Higarashi and Honda. *Drugs*. 2020;80:1183–201.

Disclosure of Interests: Kevin Winthrop Consultant of: AbbVie, Bristol-Myers Squibb, Eli Lilly and Co., Galapagos NV, Gilead Sciences, GlaxoSmithKline, Pfizer, Roche, and UCB, Grant/research support from: AbbVie, Bristol-Myers Squibb, and Pfizer, Maya H Buch Speakers bureau: AbbVie; Eli Lilly and Company; Gilead Sciences, Inc.; Merck-Serono; Pfizer; Roche; Sandoz; Sanofi; and UCB, Consultant of: AbbVie; Eli Lilly and Company; Gilead Sciences, Inc.; Merck-Serono; Pfizer; Roche; Sandoz; Sanofi; and UCB, Grant/research support from: AbbVie; Eli Lilly and Company; Gilead Sciences, Inc.; Merck-Serono; Pfizer; Roche; Sandoz; Sanofi; and UCB, Jeffrey Curtis Grant/research support from: AbbVie, Amgen, BMS, Corrona, Eli Lilly, Janssen, Myriad, Pfizer, Regeneron, Roche, and UCB, Gerd Rüdiger Burmester Speakers bureau: AbbVie; Eli Lilly; Pfizer; and Gilead Sciences, Inc., Consultant of: AbbVie; Eli Lilly; Pfizer; and Gilead Sciences, Inc., Daniel Aletaha Speakers bureau: AbbVie, Amgen, Bristol-Myers Squibb, Celgene, Eli Lilly, Medac, Merck, Merck Sharp & Dohme, Novartis, Pfizer, Roche, Sandoz, Sanofi/Genzyme, and UCB, Consultant of: AbbVie, Amgen, Celgene, Eli Lilly, Janssen, Medac, Merck, Novartis, Pfizer, Roche, Sandoz, and Sanofi/Genzyme, Grant/research support from: AbbVie, Merck Sharp & Dohme, Novartis, and Roche, Koichi Amano Speakers bureau: AbbVie GK, Astellas, Chugai Pharmaceutical Co. Ltd., Eli Lilly, GlaxoSmithKline KK, Pfizer Japan, Mitsubishi-Tanabe Pharma, Grant/research support from: Asahi Kasei Pharma, Alena Pechonkina Shareholder of: Gilead Sciences, Inc., Employee of: Gilead Sciences, Inc., Iyabode Tiarniyu Shareholder of: Gilead Sciences, Inc., Employee of: Gilead Sciences, Inc., Cianna Leatherwood Shareholder of: Gilead Sciences, Inc., Employee of: Gilead Sciences, Inc., Lei Ye Shareholder of: Gilead Sciences, Inc., Employee of: Gilead Sciences, Inc., Qi Gong Shareholder of: Gilead Sciences, Inc., Employee of: Gilead Sciences, Inc., Robin Besuyen Shareholder of: Galapagos, BV, Employee of: Galapagos, BV, James Galway Speakers bureau: Pfizer, Bristol-Myers Squibb, UCB and Celgene
DOI: 10.1136/annrheumdis-2021-eular.1408

Rheumatoid arthritis - prognosis, predictors and outcome

POS0093

HETEROGENEITY IN ADVERSE EVENT ASSESSMENT BETWEEN COUNTRIES PARTICIPATING IN AN INTERNATIONAL COLLABORATION OF REGISTRIES OF RHEUMATOID ARTHRITIS PATIENTS USING JANUS KINASE INHIBITORS (THE JAK-POT STUDY)

K. Lauper¹, D. Mongin¹, S. A. Bergstra², D. Choquette³, C. Codreanu⁴, D. De Cock⁵, L. Dreyer⁶, O. Elkayam⁷, K. Hyrich⁸, F. Iannone⁹, N. Inanc¹⁰, E. Kristianslund¹¹, T. K. Kvien¹¹, B. Leeb¹², G. Lukina¹³, D. Nordström¹⁴, K. Pavelka¹⁵, M. Pombo-Suarez¹⁶, Z. Rotar¹⁷, M. J. Santos^{18,19}, A. Strangfeld²⁰, D. Courvoisier¹, A. Finckh¹ on behalf of Epidemiology and registry working group. ¹Geneva University Hospital, Rheumatology, Geneva, Switzerland; ²LUMC, Rheumatology, Leiden, Netherlands; ³CHUM, Institut de Recherche en Rhumatologie, Montréal, Canada; ⁴University of Medicine, Center of Rheumatic Diseases, Bucharest, Romania; ⁵KU Leuven, Skeletal Biology and Engineering Research Center, Leuven, Belgium; ⁶Aalborg University Hospital, Rheumatology, DANBIO Aalborg, Aalborg, Denmark; ⁷Tel Aviv University, Rheumatology, Tel Aviv, Israel; ⁸University of Manchester, Centre for Epidemiology Versus Arthritis, Manchester, United Kingdom; ⁹University Hospital of Bari, GISEA, Rheumatology, Bari, Italy; ¹⁰Marmara University School of Medicine, Rheumatology, Istanbul, Turkey; ¹¹Diakonhjemmet Hospital, Rheumatology, Oslo, Norway; ¹²University Hospital St. Poelten, Rheumatology, St. Poelten, Austria; ¹³V.A. Nasonova Research Institute, Rheumatology, Moscow, Russian Federation; ¹⁴Helsinki University, ROB-FIN, Helsinki, Finland; ¹⁵Institute of Rheumatology, Rheumatology, Prague, Czech Republic; ¹⁶Hospital Clinico Universitario, Rheumatology, Santiago de Compostela, Spain; ¹⁷University Medical Centre Ljubljana & University of Ljubljana, Rheumatology, Ljubljana, Slovenia; ¹⁸Rheuma.pt, RA, Lisbon, Portugal; ¹⁹Faculdade de Medicina de Lisboa, Rheumatology Research Unit, Instituto de Medicina Molecular, Lisbon, Portugal; ²⁰DRFZ, Programme Area Epidemiology, Berlin, Germany

Background: Industry, regulators, and the rheumatology community have recognized the need for observational studies to monitor the safety of new anti-rheumatic agents. Registries provide a unique opportunity to understand the safety of newer therapies, but pharmacovigilance studies require large number of patients to evaluate rare drug-related adverse-events (AEs). Because JAK-inhibitors (JAKi) have only recently been approved for the treatment of rheumatoid arthritis, it makes sense to combine data from several registries in order to obtain a sufficiently large sample size to promote earlier detection of adverse events.

Objectives: The purpose of this analysis was to evaluate how AEs are assessed in the various registries in preparation for a collaborative pharmacovigilance analysis, and present preliminary results.

Methods: The “JAK-pot” collaboration includes 19 RA registries. The principal investigators of the participating registries were sent a structured questionnaire on AE assessment and 18 (94%) provided complete responses on the AE assessment procedures of their registries. We present simple descriptive statistics of the AE assessment procedures employed by the participating registries.

Results: The 19 registries represent 7186 patients initiating a JAKi (Table 1), who are on average 57 years old, with a mean disease duration 11 years, seropositive (83%), female (82%) and with moderate disease activity at treatment initiation.

Table 1.

Country, registry	N° of patients on JAKi included
Austria, BIOREG	87
Belgium, TARDIS	2113
Canada, RHUMADATA	363
Czech Republic, ATTRA	197
Denmark, DANBIO	506
Finland, ROB-FIN	229
Germany, RABBIT	620
Italy, GISEA	244
Israel, I-RECORD	96
Netherlands, METEOR	4
Norway, NOR-DMARD	97
Portugal, REUMA.PT	44
Romania, RRBR	252
Russia, ARBITER	428
Slovenia, biorx.si	141
Spain, BIOBADASER	139
Switzerland, SCQM	738
Turkey, TURKBIO	404
UK, BSRBR	484

After ineffectiveness, AEs was the second most common reason for JAKi discontinuation (25.5%), with large differences between registries (Figure 1).

Of the participating registries, 2 registries do not collect AEs, while 16 (89%) assess incident AEs, by means of a pre-specified extraction form (3 registries), by free text (5 registries), by a combination of both (6 registries) and/or the use of linkage to external electronic records (3registries). AEs are coded using a pre-defined coding system by 11 registries (MeDRA (8), other (3)), but nearly all are recording the severity of the AE (15, 94%), AE related-death (15, 94%), or AE-related hospitalisation (15, 94%). AEs of special interest, such as serious infections (15, 94%), thromboembolic events (15, 94%), or shingles (9, 56%), are recorded by most registries. Incident AEs are linked by the treating physician to specific therapies in 11 registries (69%), while the other 5 registries extrapolate potential causal associations based on therapy start and stop dates. A pre-specified adjudication process for AEs is made only by 5 registries (31%).

Conclusion: Substantial heterogeneity exists among registries regarding AE assessment within the JAK-pot collaboration. These differences must be taken into account when analysing the safety of JAKi across different countries in collaborative studies. For comparative analyses, stratified analyses by country are required to account for differential AE assessment and varying degrees of potential under-reporting.

Disclosure of Interests: Kim Lauper: None declared, Denis Mongin: None declared, Syske Anne Bergstra: None declared, Denis Choquette: None declared, Catalin Codreanu: None declared, Diederik De Cock: None declared, Lene Dreyer: None declared, Ori Elkayam: None declared, Kimme Hyrich: None declared, Florenzo Iannone: None declared, Nevsun Inanc: None declared, Eirik Kristianslund: None declared, Tore K. Kvien: None declared, Burkhard Leeb: None declared, Galina Lukina: None declared, Dan Nordström: None declared, Karel Pavelka: None declared, Manuel Pombo-Suarez: None declared, Ziga Rotar: None declared, Maria Jose Santos: None declared, Anja Strangfeld: None declared, Delphine Courvoisier: None declared, Axel Finckh Speakers bureau: Eli-Lilly, Pfizer, Consultant of: Eli-Lilly, Pfizer, Grant/research support from: BMS, Pfizer.

DOI: 10.1136/annrheumdis-2021-eular.2216

POS0094

EFFECTS OF RANKL INHIBITION ON PROMOTING HEALING OF BONE EROSION IN RHEUMATOID ARTHRITIS USING HR-pQCT: A 2-YEAR, RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED TRIAL

H. So¹, I. T. Cheng¹, S. L. Lau¹, E. Chow¹, T. Lam¹, V. W. Hung², E. Li¹, J. F. Griffith³, V. W. Lee⁴, L. Shi³, J. Huang³, Y. K. Kwok⁵, I. C. Yim⁶, T. K. Li¹, V. Lo¹, J. M. Lee⁷, J. J. W. Lee⁸, L. Qin², L. S. Tam¹. ¹The Chinese University of Hong Kong, Medicine and Therapeutics, Hong Kong, Hong Kong (SAR); ²The Chinese University of Hong Kong, Orthopedics & Traumatology, Hong Kong, Hong Kong (SAR); ³The Chinese University of Hong Kong, Imaging & Interventional Radiology, Hong Kong, Hong Kong (SAR); ⁴The Chinese University of Hong Kong, School of Pharmacy, Hong Kong, Hong Kong (SAR); ⁵Queen Elizabeth Hospital, Medicine, Hong Kong, Hong Kong (SAR); ⁶Tseung Kwan O Hospital, Medicine, Hong Kong, Hong Kong (SAR); ⁷Tai Po Hospital, Medicine and Geriatrics, Hong Kong, Hong Kong (SAR); ⁸The Chinese University of Hong Kong, The Jockey Club School of Public Health & Primary Care, Hong Kong, Hong Kong (SAR)

Background: Partial repair of bone erosions in rheumatoid arthritis (RA) is known from high-resolution peripheral quantitative computer tomography (HR-pQCT) studies in patients with moderate to high disease activity using biologics [1]. Whether RANKL inhibition by denosumab is efficacious in healing existing erosions in RA patients with low disease activity or in remission on conventional synthetic DMARDs is uncertain.

Objectives: To evaluate the effects of denosumab on erosion healing at 2-4 metacarpophalangeal head as determined by HR-pQCT in patients with RA with stable disease.

Methods: This was a randomized, placebo-controlled, double-blind study. RA patients with disease activity score 28 joints (DAS28) ≤ 5.1 were randomized (1:1) to subcutaneous denosumab 60mg or placebo once every six months for 24 months. The primary outcome was erosion healing at MCP 2-4 on HR-pQCT at 12 months. The effects of denosumab on erosion and joint space parameters on HR-pQCT and radiographs, disease activity and health assessment questionnaire-disability index (HAQ-DI) were also examined.

Results: At 24 months, HR-pQCT images were analyzed in 98 patients. Baseline demographic, clinical characteristics and imaging parameters were comparable between the two treatment groups (table 1). Seventeen patients in each group (placebo group: 17/52, 32.6%; denosumab group: 17/50, 34.0%) achieved sustained low disease activity (DAS28 ≤ 3.2) throughout the 24 months. At 12 months, changes in erosion parameters on HR-pQCT were similar between the two groups. At 24 months, new erosions (19% vs 9%, $p=0.009$) and erosion progression (34% vs 16%, $p<0.001$) were more common in the placebo group than the denosumab group. Erosion healing was seen in a significantly higher proportion of patients in the denosumab group (20% vs 6%, $p=0.045$) at 24 months. The details of the changes in HR-pQCT erosion parameters are shown in figure 1. No significant differences in the changes in joint space parameters on HR-pQCT, van der Heijde-Sharp erosion score, DAS28 and HAQ-DI were observed between the two groups at 12 and 24 months.

Conclusion: Although no differences in erosion parameters were observed at 12 months, denosumab was more efficacious than placebo in erosion repair on HR-pQCT after 24 months.

REFERENCES:

- [1] Finzel S, Rech J, Schmidt S, et al. Interleukin-6 receptor blockade induces limited repair of bone erosions in rheumatoid arthritis: a micro CT study. *Ann Rheum Dis* 2013;72:396-400.

Table 1. Baseline clinical, demographic, disease activity parameters and medications

	Placebo (n=55)	Denosumab (n=55)	Total (n=110)
Age	56.5 $\pm$ 7.1	57.2 $\pm$ 8.5	56.8 $\pm$ 7.8
Gender (Female)	47 (86)	41 (75)	88 (80)
Disease duration (years)	8.5 $\pm$ 6.8	7.3 $\pm$ 6.9	7.9 $\pm$ 6.8
Rheumatoid factor positive	40 (72)	38 (69)	78 (71)
ACPA positive	43 (78)	44 (80)	87 (79)
DAS28-CRP	2.43 $\pm$ 0.83	2.6 $\pm$ 0.92	2.51 $\pm$ 0.88
DAS28-CRP ≥ 3.2	8 (15)	13 (24)	21 (19)
HAQ-DI (0-3)	0.31 $\pm$ 0.38	0.46 $\pm$ 0.47	0.39 $\pm$ 0.43
csDMARDs	49 (89)	52 (95)	101 (92)
Combination csDMARDs	26 (47)	33 (60)	59 (54)
Glucocorticoids	5 (10)	5 (9)	10 (9)
vdH- Sharp erosion score	10.4 $\pm$ 18.4	8.9 $\pm$ 13.8	9.6 $\pm$ 16.2
vdH- Sharp JSN score	12.4 $\pm$ 17.7	11.5 $\pm$ 17.2	11.9 $\pm$ 17.4
Lumbar spine aBMD, g/cm ²	0.914 $\pm$ 0.147	0.930 $\pm$ 0.143	0.922 $\pm$ 0.145
Total hip aBMD, g/cm ²	0.837 $\pm$ 0.102	0.847 $\pm$ 0.146	0.841 $\pm$ 0.125
Femoral neck aBMD, g/cm ²	0.681 $\pm$ 0.099	0.695 $\pm$ 0.128	0.687 $\pm$ 0.114

Data are reported as mean $\pm$ SD or number (%). ACPA: Anti-cyclic citrullinated peptide antibody; DAS28: disease activity score 28; csDMARDs: conventional synthetic disease modifying anti-rheumatic drug. HAQ-DI: health assessment questionnaire disability index; vdH- Sharp score: Van der Heijde- Sharp score; aBMD: areal bone mineral density

Figure 1. Changes in erosion parameters by HR-pQCT. (A) Percentage of patients with overall erosion healing; (B) Outcome of individual erosion with healing, progression and new erosion detected across study period; change in (C) mean erosion volume; (D) total erosion volume; (E) erosion width; (F) erosion depth and (G) marginal osteosclerosis per patient.

Disclosure of Interests: Ho SO: None declared, Isaac T. Cheng: None declared, Sze-Lok Lau: None declared, Evelyn Chow: None declared, Tommy Lam: None declared, Vivian W Hung: None declared, Edmund Li: None declared, James F Griffith: None declared, Vivian WY Lee: None declared, Lin Shi: None declared, Junbin Huang: None declared, Yan Kitty Kwok: None declared, Isaac C Yim: None declared, Tena K. Li: None declared, Vincent Lo: None declared, Jolly M Lee: None declared, Jack Jock Wai Lee: None declared, Ling Qin: None declared, Lai-Shan Tam Grant/research support from: Grants from Novartis and Pfizer

DOI: 10.1136/annrheumdis-2021-eular.2752