were genotyped in 386 Caucasian IgAV patients (the largest series of Caucasian patients with IgAV ever assessed for genetic studies) and 806 sex and ethnically matched healthy controls by TaqMan assays.

Results: No statistically significant differences in the genotype and allele frequencies between patients with IgAV and healthy controls were observed when each genetic variant of *BAFF APRIL* and *BAFFR* was analyzed independently (Table 1). Likewise, no statistically significant differences in genotype and allele frequencies of *BAFF APRIL* or *BAFFR* were found when patients with IgAV were stratified according to the age at disease onset or to the presence/absence of gastrointestinal or renal manifestations. Similar results were disclosed when haplotype frequencies of *APRIL* and *BAFFR* were compared between patients with IgAV and healthy controls as well as patients with IgAV stratified according to the age at disease onset or to the presence/absence of gastrointestinal or renal manifestations.

Conclusion: Our results suggest that the *BAFF-APRIL-BAFFR* pathway does not contribute to the genetic network underlying IgAV.

REFERENCES:

- [1] J Exp Med 1999;190:1697-710; [2] Science 1999;285:260-3; [3] Nat Genet 2005;37:829-34; [4] Nat Immunol 2002;3:822-9; [5] N Engl J Med 2017;376:1615-26; [6] N Engl J Med 2013;368:2402-14; [7] Autoimmun Rev 2018;17:301-315.

Table 1. Genotype and allele frequencies of *BAFF*, *APRIL* and *BAFFR* genes in patients with IgA vasculitis and healthy controls.

Polymorphism	Locus	1/2	Data set	1/1	1/2	2/2	1	2
rs374039502	BAFF	T/A	Patients	91.9 (353)	8.1 (31)	0	95.9 (737)	4.1 (31)
			Controls	91.5 (733)	8.1 (65)	0.4 (3)	95.6 (1531)	4.4 (71)
rs11552708	APRIL	G/A	Patients	78.1 (299)	20.6 (79)	1.3 (5)	88.4 (677)	11.6 (89)
			Controls	77.9 (625)	20.4 (164)	1.6 (13)	88.1 (1414)	11.9 (190)
rs6608	APRIL	C/T	Patients	71.9 (277)	26.0 (100)	2.1 (8)	84.9 (654)	15.1 (116)
			Controls	70.0 (561)	27.6 (221)	2.5 (20)	83.7 (1343)	16.3 (261)
rs7290134	BAFFR	A/G	Patients	58.0 (224)	36.3 (140)	5.7 (22)	76.2 (588)	23.8 (184)
			Controls	57.2 (459)	36.4 (292)	6.5 (52)	75.3 (1210)	24.6 (396)
rs77874543	BAFFR	G/C	Patients	82.7 (316)	16.0 (61)	1.3 (5)	90.7 (693)	9.3 (71)
			Controls	83.0 (666)	16.6 (133)	0.4 (3)	91.3 (1465)	8.7 (139)

Acknowledgements: This study was supported by European Union FEDER funds and "Fondo de Investigaciones Sanitarias" (grant PI18/00042) from 'Instituto de Salud Carlos III' (ISCIII, Health Ministry, Spain). DP-P is a recipient of a Río Hortega programme fellowship from the ISCIII, co-funded by the European Social Fund (ESF, 'Investing in your future') [grant number CM20/00006]; SR-M is supported by funds of the RETICS Program co-funded by the European Regional Development Fund (ERDF) [grant number RD16/0012/0009]; VP-C is supported by a pre-doctoral grant from IDIVAL [grant number PREVAL 18/01]; BA-M is a recipient of a 'López Albo' Post-Residency Programme funded by Servicio Cántabro de Salud; LL-G is supported by funds of IDIVAL [grant number INNVAL20/06]; RL-M is a recipient of a Miguel Servet type I programme fellowship from the ISCIII, co-funded by the European Social Fund (ESF, 'Investing in your future') [grant number CP16/00033].

Disclosure of Interests: Diana Prieto-Peña Speakers bureau: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen and MSD, Consultant of: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen and MSD, Grant/research support from: Abbvie, MSD and Roche, Sara Remuzgo Martínez: None declared, Fernanda Gené: None declared, Verónica Pulito-Cueto: None declared, Belén Atienza-Mateo: None declared, Belén Sevilla: None declared, Javier Llorca: None declared, Norberto Ortego: None declared, Leticia Lera-Gómez: None declared, Maite Leonardo: None declared, Ana Peñalba: None declared, J. Narváez: None declared, Luis Martín-Penagos: None declared, Jose Alberto Miranda-Fillooy: None declared, LUIS CAMINAL MONTERO: None declared, PAZ COLLADO: None declared, Javier Sanchez Perez: None declared, Diego de Argila: None declared, Esteban Rubio-Romero: None declared, MANUEL LEON LUQUE: None declared, Juan María Blanco-Madrugal: None declared, E. Galindez: None declared, Javier Martín Ibanez: None declared, Santos Castañeda: None declared, Ricardo Blanco Speakers bureau: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen and MSD, Consultant of: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen and MSD, Grant/research support from: Abbvie, MSD and Roche, Miguel A González-Gay Speakers bureau: Pfizer, Abbvie, MSD, Grant/research support from: Pfizer, Abbvie, MSD, Raquel López-Mejías: None declared

DOI: 10.1136/annrheumdis-2021-eular.707

POS0114 ABERRANT MONOCYTE SUBSETS IN PATIENTS WITH BECHET'S DISEASE

C. Li^{1,2}, J. Liu¹, H. Chen¹, W. Zheng¹. ¹Peking Union Medical College Hospital, Rheumatology and Clinical Immunology, Beijing, China; ²Peking University Shougang Hospital, Rheumatology, Beijing, China

Background: Behçet's Disease (BD) is a systemic vasculitis of unknown etiology[1]. Monocytes closely related to inflammation potentially contribute to BD's pathogenesis. They are classified into three subsets: classical monocytes (CM), intermediate monocytes (IM) and non-classical monocytes (NCM). Abnormalities of monocyte subsets have been reported in many infectious, inflammatory and autoimmune diseases[2-6], but their implication in BD remains elusive.

Objectives: To investigate the distribution, phenotypes and functions of monocyte subsets in BD and explore their roles in BD pathogenesis.

Methods: The frequencies and phenotypes of monocyte subsets in BD and healthy controls (HC) were determined by flow cytometry, and their correlation with clinical parameters was analyzed. Intracellular cytokines and phosphorylated signal proteins [phosphorylated p65(p-p65) and phosphorylated p38(p-p38)] were determined in LPS-activated monocyte subsets by flow cytometry. Monocyte subsets of BD and HC were sorted and co-cultured with naïve CD4+ T cells, and Th1 cell frequencies were measured on day 5.

Results: A higher IM (9.0±3.6% vs. 4.5±2.0%, p<0.01) and lower NCM (2.6±1.2% vs. 4.2±2.0%, p<0.01) population in BD patients were noted. BD IM were positively correlated with CRP (r=0.5456, p<0.05), ESR (r=0.4683, p=0.05), and the serum level of TNF-α (r=0.7372, p<0.001) and IL-6 (r=0.5013, p<0.05). BD NCM were negatively correlated with CRP (r=0.4822, p<0.05) and the serum level of IgM (r=-0.7830, p<0.001). Moreover, BD IM decreased (12.3±3.8% vs. 5.7±3.6%, p<0.05), while BD NCM increased (2.6±1.3% vs. 3.5±1.5%, p<0.01) after BD patients achieved remission. CD11b and CD64 expression on CM, IM and NCM in BD were enhanced. BD CM promoted TNF-α (61.0±11.4% vs 48.3±9.9%, p<0.05) and IL-6 (7.2±5.4% vs 1.9±1.7%, p<0.05) production and facilitated Th1 differentiation. BD IM promoted IL-6 production (6.2±3.8% vs 2.6±1.6%, p<0.05). Furthermore, we demonstrated a higher level of p-p65 (12.8±2.9% vs 3.3±1.1%, p<0.01) in BD CM and increased p-p65 (3.2±0.6% vs 0.01±0.01%, p<0.01) and p-p38 (1.1±0.6% vs 0.03±0.01%, p<0.01) in BD IM.

Conclusion: To our knowledge, our study is the first study on monocyte subsets in BD. Our data highlighted the aberrant populations of IM and CM in BD, potentially implicated in BD pathogenesis.

REFERENCES:

- [1] Ahmet, Gül, Pathogenesis of Behçet's disease: autoinflammatory features and beyond, Semin. Immunopathol. 37 (2015) 413-418.
- [2] G.Fingerle, A.Pforte, B.Passlick, et al.The novel subset of CD14+/CD16+ blood monocytes is expanded in sepsis patients, Blood. 82 (1993) 3170-3176.
- [3] M.Moniuszko, A. Bodzenta-Lukaszyk, K. Kowal, et al. Enhanced frequencies of CD14+/CD16+, but not CD14+/CD16+, peripheral blood monocytes in severe asthmatic patients, Clin. Immunol. 130 (2009) 338-346.
- [4] M.D.Sanchez, Y.Garcia, C.Montes, et al.Functional and phenotypic changes in monocytes from patients with tuberculosis are reversed with treatment, Microbes. Infect. 8 (2006) 2492-2500.
- [5] S.Koch, T.Kucharzik, J.Heidemann, et al. Investigating the role of proinflammatory CD16+ monocytes in the pathogenesis of inflammatory bowel disease, Clin. Exp. Immunol. 161 (2010) 332-341.
- [6] H. Zhu, F. Hu, X. Sun, et al.CD16(+) Monocyte Subset Was Enriched and Functionally Exacerbated in Driving T-Cell Activation and B-Cell Response in Systemic Lupus Erythematosus, Front. Immunol. 7 (2016) 512.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1157

POS0115 RISK FACTORS FOR CORONARY HEART DISEASE AND ATHEROSCLEROSIS IN PATIENTS WITH BECHET'S DISEASE

R. Goloeva¹, Z. Alekberova¹, T. Popkova¹, S. Glukhova¹. ¹V.A. Nasonova Research Institute of Rheumatology, systemic rheumatic diseases, Moscow, Russian Federation

Background:

Objectives: To evaluate traditional risk factors for clinically manifest and asymptomatic atherosclerosis in pts with BD.

Methods: Traditional risk factors for atherosclerosis were evaluated in 95 pts with Bechet's disease (BD) and 40 healthy age and gender matched controls. The male-to-female ratio was 3.7:1, pts' mean age was 29.7 (23-35) y, the mean age at the disease onset - 19.9 (14-25) y, and mean disease duration - 9.6 (4-15) y. The common carotid artery intima-media thickness (IMT) was evaluated by high resolution B-mode ultrasonography. A plaque was defined as a local intima-media thickening ≥1.2mm, with reference IMT thickness <0.9mm. Serum concentration of high-sensitivity C-reactive protein (hs CRP) was measured by immunonephelometric assay (BN-100 Analyzer; Dade Behring). Lipid profile evaluation included total cholesterol, TGs, HDL, LDL and atherogenic index.

Results: Traditional risk factors for atherosclerosis (one or more) were found in 85.3% of BD pts, dyslipidemia was the most common risk factor and was present in 76.84% of pts, mostly due to increased total cholesterol (TC) and