easy" to complete; remaining responses: "neutral" 20%, "somewhat difficult" 10%, "extremely difficult" 0%.

Conclusion: We have created and tested a system of remote clinical management for patients with RA and AS. Amongst the 108 responders, just 31% required a face-to-face appointment, with treatment changes made accordingly. With a backlog of 3,800 awaiting allocation to follow-up appointments, remote clinical management will allow us to safely and efficiently prioritise patients requiring urgent follow-up for treatment optimisation. We will integrate this system into our standard care pathway beyond the COVID-19 pandemic to streamline our service, deliver effective care and provide evidence to support the use of costly biologic drugs.¹ We plan to investigate the barriers for non-responders.

REFERENCES:

- [1] Holroyd CR, Seth R, Bukhari M, et al. The British Society for Rheumatology biologic DMARD safety guidelines in inflammatory arthritis. *Rheumatology*. 2019; 58 (2): e3–e42.

Disclosure of Interests: Tamir Malley: None declared, John Jackman: None declared, Sarah Manderson: None declared, Larissa Saldana Pena Grant/ research support from: Pfizer's Global Medical Grants program, Ellie Evans: None declared, Joe Barrett Grant/research support from: Pfizer's Global Medical Grants program, Anushka Soni Grant/research support from: Pfizer's Global Medical Grants program, Raashid Luqmani Grant/research support from: Pfizer's Global Medical Grants program

DOI: 10.1136/annrheumdis-2021-eular.309

POS0153-HPR IDENTIFYING TARGETS FOR BEHAVIOURAL INTERVENTIONS IN FIBROMYALGIA: PHYSICAL ACTIVITY BEHAVIOUR

K. Mayana¹, Y. Prior². ¹University of Salford, School of Health and Society, Manchester, United Kingdom; ²University of Salford, School of Health and Society, Salford, United Kingdom

Background: There has been much interest in the efficacy of exercise and physical activity interventions in people with Fibromyalgia. This has resulted in guidelines recommending exercise as the first line of management (Macfarlane et al., 2017). Notwithstanding the efficacy of exercises, adherence to structured exercise and interventions aiming to increase physical activities remains a considerable challenge in this population. Exploring behavioural factors that may hinder exercise and physical activity participation in people with Fibromyalgia may help to identify targets for interventions to improve adherence and engagement with physical activities and support self-management.

Objectives: To explore people with Fibromyalgia's perspectives on the personal, social and contextual factors impacting their physical activity behaviour and pain perception.

Methods: Twelve participants with Fibromyalgia were recruited from Social Media, Fibromyalgia charities, organisations and associations across the UK. Semi-structured, one-to-one in-depth telephone interviews were conducted with each participant. Interviews were audio-recorded and transcribed verbatim. Qualitative data analysis was conducted by two researchers independently, using Thematic Analysis. Critical Realism framework was employed to inform the interpretation of the findings.

Results: Four main themes emerged from the data: (i) Lack of guidance on physical activity: most participants expressed increased pain intensity following a vigorous physical activity which, have led to spending more time on sedentary behaviour to recover from pain. (ii) Participants expressed fear of fatigue as a barrier to physical activity participation: This also led to increased sedentary and protective behaviours. (iii) Impact of treatments on physical activity: Participants who received multimodal therapies and patient education reported better coping strategies e.g., pacing physical activities and less pain. (iv): Impact of social support on physical activity participation: Participants felt that a lack of understanding from employers, their family and friends and the wider society has negatively impacted their physical activity behaviours, access to workplace support and psychological wellbeing.

Conclusion: A multimodal approach, incorporating patient education with behavioural components is essential to increase adherence to and engagement with exercise and physical activity interventions to support self-management of Fibromyalgia. There is a need to improve the public perception of this syndrome to increase social and workplace support.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.3239

POS0154-HPR FACTORS FACILITATING THE SELF- AND SHARED-MANAGEMENT OF JIA BY CHILDREN, YOUNG PEOPLE, THEIR FAMILIES, AND PROFESSIONALS INVOLVED IN THEIR CARE: A REALIST EVALUATION

S. Stones¹, V. Swallow², L. Milnes¹. ¹University of Leeds, School of Healthcare, Leeds, United Kingdom; ²Sheffield Hallam University, Department of Nursing & Midwifery, Sheffield, United Kingdom

Background: Juvenile idiopathic arthritis (JIA) is a long-term condition, often requiring some element of lifelong management. Therefore, it is logical that children and young people are empowered to become competent at self-managing their health and wellbeing, while families are supported in their shared-management role during childhood, relinquishing control at age- and developmentally appropriate periods in their child's development. However, there are limited theoretical foundations underpinning optimal self- and shared-management support for children, young people and families living with JIA.

Objectives: To explain the factors facilitating the self- and shared-management of JIA by children, young people, and their families, with professional support from healthcare professionals, third sector organisations, and education professionals.

Methods: Guided by the Individual and Family Self-management Theory and the Shared Management Model, a three-stage realist evaluation was undertaken: 1) initial JIA self- and shared-management question theories were elicited from a document review, integrative review, and stakeholder insights [1]; 2) seven initial question theories were tested using qualitative research methods with 20 participants (young people, families, healthcare professionals, education professionals, and third sector representatives); 3) analysis of findings using a theory-driven approach to thematic analysis, in order to identify demi-regularities to extend or refute the initial question theories. The analysis drew on deductive, inductive, and retroductive reasoning.

Results: There were six refined JIA self- and shared-management question theories: 1) meaningful and bespoke self-management support across the life course for children and young people with JIA; 2) recognised and valued shared-management support for the families of children and young people with JIA, with autonomy in mind; 3) individual healthcare plans as a shared management communication tool to facilitate optimal management of JIA; 4) consistent recognition, value, and encourage of self- and shared-management support from the paediatric rheumatology multi-disciplinary team and associated professionals; 5) child, young-person, and family-focused paediatric rheumatology care and support services across the lifecourse; and 6) bespoke and inclusive approaches by education providers to enable children and young people with JIA to feel safe, supported, and able to fulfil their potential.

Conclusion: There is an increasing recognition of the importance of self- and shared-management of JIA. However, there is a lack of an overall, cohesive approach to self- and shared-management between healthcare providers, education providers, and the third sector. Findings from this study illuminate the factors facilitating JIA self- and shared-management at individual, interpersonal, institutional and infrastructural levels. Further work is required to empirically test these refined question theories with interventions designed to enhance JIA care, education, and support.

REFERENCES:

- [1] Stones et al. (2020). A realist approach to eliciting the initial programme theories for the self- and shared-management of juvenile idiopathic arthritis by children, young people, families and professionals involved in their care. *Pediatric Rheumatology* 18(Suppl 2): O062.

Acknowledgements: The authors thank all of the participants who helped to shape the findings of this study. This work formed part of a PhD study funded by the University of Leeds.

Disclosure of Interests: Simon Stones Speakers bureau: Janssen, Consultant of: Parexel, Envision Pharma Group, 67 Health, On The Pulse Consultancy, Veronica Swallow: None declared, Linda Milnes: None declared

DOI: 10.1136/annrheumdis-2021-eular.1286

POS0155-HPR DIAPHRAGMATIC BREATHING RELAXATION TECHNIQUE TO DECREASE ANXIETY DURING JOINT INFILTRATION: A RANDOMIZED CONTROLLED TRIAL

R. Ben Aissa^{1,1}, S. Boussaid², H. Tbin², H. Sahli¹, S. Jemmal¹, S. Rekik¹, M. El Euch². ¹Rabta Hospital, Department of Rheumatology, Tunis, Tunisia; ²Rabta Hospital, Department of Rheumatology, Tunis, Tunisia

Background: Joint infiltration is the injection of therapeutic substances directly into a joint. It may be a stressful experience for patients as imagined different from other usual injections. Several techniques are used to manage anxiety and pain during such a procedure.

Objectives: To evaluate the effectiveness of diaphragmatic breathing relaxation on reducing anxiety and pain during joint infiltration.

Methods: Patients scheduled for a joint infiltration at the rheumatology department's daycare unit were recruited. All infiltrations were performed using steroids without anesthetic therapy except for the hip. Patients were randomized into two groups (cases=38, controls=34). Cases learned from a trained health agent diaphragmatic breathing relaxation technique to perform it immediately before and during the procedure while controls received the usual procedure. We used the Visual Analogue Scale (VAS) to assess self-estimated both anxiety (VAS-Anx) and pain (VAS-Pain) as evaluated on pre and post-joint infiltration. VAS-Pain was evaluated as expected then as experienced respectively on pre and