based on medical evidence "there is an almost religious way of thinking on how to deal with the pathology. It is not an exact science when you choose the physicians you choose the treatment". 2) *Relationship with the self and the others* as some patients did not feel understood or even shameful and hopeless about their condition. 3) *Patients' and Health Professionals' beliefs about the pathology management* where common thoughts were the perceived (ab)use of passive therapies, the movement as something dangerous and that OA is "something that you try to resist to, but (surgery) is your destiny". 4) *facilitators* and 5) *barriers of the adherence to therapeutic exercise* that revolve around the cost of the therapy, the time needed and the willingness to change life habits. 6) *Patients' attitudes towards pathology* in which the oldest patients perceive OA as "something I have to accept since I am getting old" and the youngest as "Something I have to fight". 7) *Relationship with food* in which diet is seen as something that "you force yourself to follow" which is useful only to lose weight and not to preserve a high health status and where overeating is used "to eat your feelings".

Conclusion: Patients suffering from hip and knee OA seem to experience an uncertain care process. The lack of clear explanations and the attitude towards conservative treatment, which is considered as "a pastime while waiting for surgery," fosters the importance of providing patients with adequate information about the treatment, to shift their beliefs and improve their awareness. This will enhance a patient-centred and shared decision-making treatments.

REFERENCES:

- [1] Fernandes L, Hagen KB, Bijlsma JWW, *et al.* EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis. *Ann. Rheum. Dis.* 2013;72:1125–35.
- [2] Basedow M, Esterman A. Assessing appropriateness of osteoarthritis care using quality indicators: a systematic review. *J Eval Clin Pract* 2015;21:782–9.

Figure 1. Categories and Subcategories stemmed from the analysis of the patients' interviews

Acknowledgements: This work is part of the project funded by EULAR Health Professionals Research Grant 2020.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.843

POS0161-HPR BARRIERS AND FACILITATORS RELATED TO SELF-MANAGEMENT OF SHOULDER PAIN: A SYSTEMATIC REVIEW AND QUALITATIVE SYNTHESIS

A. O'Shea¹, J. Drennan², C. Littlewood³, H. Slater⁴, J. Sim⁵, J. McVeigh¹.

¹University College Cork, Discipline of Physiotherapy, College of Medicine and Health, Cork, Ireland; ²University College Cork, School of Nursing and Midwifery, College of Medicine and Health, Cork, Ireland; ³Manchester Metropolitan University, Department of Health Professions, Faculty of Health, Psychology & Social Care, Manchester, United Kingdom; ⁴Curtin University, School of Physiotherapy and Exercise Science, Perth, Australia; ⁵Keele University, School of Medicine, Faculty of Medicine and Health Sciences, Keele, United Kingdom

Background: Shoulder pain is a significant cause of pain and disability in the general population.¹ Current research suggests that shoulder pain can be resistant to treatment and is often recurrent.² Systematic review-level evidence shows modest short-term benefits for most current treatments.³ Effective self-management strategies that equip patients with the necessary tools to self-manage their condition are urgently required. However, engagement with self-management is poor, potentially compromising treatment outcomes and contributing to ongoing disability.⁴

Objectives: The objectives of this review were to:

- Systematically identify and appraise relevant qualitative evidence on barriers and facilitators relating to self-management from the perspectives of people with shoulder pain and healthcare professionals.
- Collate and synthesise this evidence, to gain an understanding of factors that influence self-management of shoulder pain.
- Develop evidence-based recommendations to inform the implementation and delivery of self-management programmes for shoulder pain.

Methods: A meta-aggregative approach to the synthesis of qualitative evidence was used. Twelve databases were searched, from inception to 13 July 2020, to identify studies exploring barriers and facilitators related to self-management of shoulder pain from the perspectives of people with shoulder pain and clinicians involved in the care of such patients. Two independent reviewers identified eligible articles, extracted the data and conducted critical appraisal. Two reviewers independently identified and developed categories, with validation by two further researchers. Categories were discussed among the wider research team and a comprehensive set of synthesized findings was derived.

Results: Sixteen studies were included in the review, exploring several shoulder conditions: shoulder instability; rotator cuff-related pain; dysfunction post rotator cuff surgery; and degenerative rotator cuff tears. From the perspective of patients, three synthesized findings were identified that influenced self-management: (1) support for self-management, including subthemes related to patient-centred support, knowledge, time, access to equipment, and patient digital literacy; (2) personal factors, including patient beliefs, patient expectations, patient motivation, pain, and therapeutic response; and (3) external factors, including influence of the clinician and therapeutic approach. From the perspective of clinicians, two synthesized findings were identified that influenced adherence to self-management: (1) support for self-management, including education, patient-centred support, patient empowerment, time, and clinician digital literacy; and (2) preferred management approach, including clinician beliefs, expectations, motivation, therapeutic approach, and therapeutic response.

Conclusion: Patients and clinicians identified several barriers and facilitators that influenced self-management of shoulder pain. Clinicians' awareness of these factors could positively influence patient management, enhance patients' ability to self-manage, and improve treatment outcomes.

REFERENCES:

- [1] Walker-Bone K, Palmer KT, Reading I, Coggon D, Cooper C. Prevalence and impact of musculoskeletal disorders of the upper limb in the general population. *Arthritis and Rheumatism*. 2004;51(4): 642-651.
- [2] Croft P, Pope D, Silman A. The clinical course of shoulder pain: prospective cohort study in primary care. *Primary Care Rheumatology Society Shoulder Study Group. British Medical Journal*. 1996;313(7057): 601-602.
- [3] Littlewood C, May S, Walters S. A review of systematic reviews of the effectiveness of conservative interventions for rotator cuff tendinopathy. *Shoulder & Elbow*. 2013;5(3): 151-167.
- [4] Littlewood C, Malliaras P, Mawson S, May S, Walters S. Patients with rotator cuff tendinopathy can successfully self-manage, but with certain caveats: a qualitative study. *Physiotherapy*. 2014;100(1): 80-85.

Disclosure of Interests: Aidan O'Shea: None declared, Jonathan Drennan: None declared, Chris Littlewood: None declared, Helen Slater Speakers bureau: AbbVie PTY LTD 2018, Julius Sim: None declared, Joseph McVeigh: None declared

DOI: 10.1136/annrheumdis-2021-eular.1279

Metabolic bone disease / Osteoporosis

POS0162 PREDICTIVE FACTORS OF A NEW FRAGILITY FRACTURE AFTER WRIST FRAGILITY FRACTURE

S. Azevedo¹, H. Parente¹, D. Esperança Almeida², F. Guimarães¹, J. Rodrigues¹, D. Faria¹, D. Peixoto¹, J. Tavares-Costa¹, C. Afonso¹, F. Teixeira¹.

¹Unidade Local de Saúde do Alto Minho, Rheumatology Department, Ponte de Lima, Portugal; ²Hospital de Braga, Rheumatology Department, Braga, Portugal

Background: Fragility fractures (FF) are fractures that result from mechanical forces that would not ordinarily result in fracture, known as low-level (or 'low energy') trauma.¹ Studies have shown that history of wrist fracture increases the risk for subsequent FF.²

Objectives: To assess predictive factors of FF occurring after a wrist fracture.

Methods: Retrospective monocentric study that included patients with a wrist FF observed at the emergency department (ED) in a tertiary center, between 1st January 2017 and 31st December 2018. Wrist fractures were identified through the 10th International Classification of Diseases and FF were identified after revision of the clinical record. Patients with relevant missing data were excluded. Seven hundred thirty-three wrist FF were identified. After calculating a representative sample (90% confidence interval), 188 patients were included. Their clinical records until 31st December 2020 (2 to 3 years after FF) were reviewed. SPSS was used for statistical analysis and significance level was defined as 2-sided p<0.05. In multivariate analysis we included variables with a significant association in univariate analysis and those with clinical relevance (reported in others studies).

Results: Wrist fractures represented 44.3% of the FF observed at the ED.

Most patients were woman (83.5%) with a mean age of 70.7 (SD=11.2) years-old at the time of their wrist fracture. A previous FF was seen in 22.9% of patients and 13.3% had a new FF during the follow-up period.

We found an association between the occurrence of a new FF and the number of comorbidities ($p=0.012$), number of visits to the ED due to falls ($p<0.001$), previous diagnosis of chronic pulmonary disease ($p=0.029$) and hematologic pathologies ($p=0.047$), and the need for hospitalization at time of the wrist FF ($p=0.018$). No associations were found between the age at the wrist fracture time, number of drugs taken daily nor its type (anxiolytics, antiepileptics, corticoids), previous fractures (and localization), overweight/obesity and other cardiovascular risk factors, endocrinopathies, psychiatric or neurologic disease or other comorbidities. After adjustment for age, gender, anti-osteoporotic treatment and comorbidities, the main predictors of a new FF were visits to the ED for falls ($p=0.005$), chronic pulmonary disease ($p=0.040$), hematologic pathologies ($p=0.004$) and need for hospitalization ($p=0.040$) (table 1).

Table 1. Multivariate analyses: linear multiple regression for predictive factors of new fragility fracture.

Determinants	Unstandardized Coefficients B	Standardized Coefficients Beta	95.0% CI	p-value
Age	-0.025	0.975	0.924 – 1.030	NS
Gender	2.065	7.889	0.757 – 82.165	NS
Number of comorbidities	0.186	1.204	0.846 – 1.713	NS
Visits to the emergency service for falls	-2.136	0.118	0.026 – 0.529	0.005
Chronic pulmonary disease	-1.326	0.266	0.075 – 0.940	0.040
Hematologic pathologies	-4.296	0.014	0.001 – 0.255	0.004
Need for hospitalization	-2.764	0.063	0.004 – 0.887	0.040
Anti-osteoporotic treatment	0.157	1.170	0.227 – 6.017	NS

CI: Confidence Interval; NS: non-significant;

Conclusion: Certain comorbidities seem to be associated with new FF. Patients with visits to the emergency service after falls and those who needed hospitalization due to the wrist fracture were more prone to have a new FF. There might be a substantial missed opportunity for intervention in these patients.

REFERENCES:

- [1] Osteoporosis: assessing the risk of fragility fracture. London: National Institute for Health and Care Excellence (UK); 2017 Feb. PMID: 32186835.
- [2] Crandall CJ, Hovey KM, Cauley JA, Andrews CA, Curtis JR, Wactawski-Wende J, Wright NC, Li W, LeBoff MS. Wrist Fracture and Risk of Subsequent Fracture: Findings from the Women's Health Initiative Study. *Journal of bone and mineral research: the official journal of the American Society for Bone and Mineral Research*. 2015;30(11):2086–2095. doi: 10.1002/jbmr.2559.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.3298

POS0163

INCIDENT FRACTURE RISK PREDICTION USING THE FRAGILITY SCORE CALCULATED BY LUMBAR SPINE RADIOFREQUENCY ECHOGRAPHIC MULTI SPECTROMETRY (REMS) SCANS

D. Ciardo¹, P. Pisani², F. A. Lombardi², R. Franchini², F. Conversano², S. Casciaro². ¹National Research Council, Institute of Clinical Physiology, Lecce, Italy; ²National Research Council, Institute of Clinical Physiology, Lecce, Italy

Background: The main consequence of osteoporosis is the occurrence of fractures due to bone fragility, with important sequelae in terms of disability and mortality. It has been already demonstrated that the information about bone mass density (BMD) alone is not sufficient to predict the risk of fragility fractures, since several fractures occur in patients with normal BMD [1].

The Fragility Score is a parameter that allows to estimate skeletal fragility thanks to a trans-abdominal ultrasound scan performed with Radiofrequency Echographic Multi Spectrometry (REMS) technology. It is calculated by comparing the results of the spectral analysis of the patient's raw ultrasound signals with reference models representative of fragile and non-fragile bones [2]. It is a dimensionless parameter, which can vary from 0 to 100, in proportion to the degree of fragility, independently from BMD.

Objectives: This study aims to evaluate the effectiveness of Fragility Score, measured during a bone densitometry exam performed with REMS technology at lumbar spine, in identifying patients at risk of incident osteoporotic fractures at a follow-up period of 5 years.

Methods: Caucasian women with age between 30 and 90 were scanned with spinal REMS and DXA. The incidence of osteoporotic fractures was assessed during a follow-up period of 5 years. The ability of the Fragility Score to discriminate between patients with and without incident fragility fractures was

subsequently evaluated and compared with the discriminatory ability of the T-score calculated with DXA and with REMS.

Results: Overall, 533 women (median age: 60 years; interquartile range [IQR]: 54-66 years) completed the follow-up (median 42 months; IQR: 35-56 months), during which 73 patients had sustained an incident fracture.

Both median REMS and DXA measured T-score values were significantly lower in fractured patients than for non-fractured ones, conversely, REMS Fragility Score was significantly higher (Table 1).

Table 1. Analysis of T-score values calculated with REMS and DXA and Fragility Score calculated with REMS. Median values and interquartile ranges (IQR) are reported. The p-value is derived from the Mann-Whitney test.

	Patients without incident fragility fracture	Patients with incident fragility fracture	p-value
T-score DXA [median (IQR)]	-1.9 (-2.7 to -1.0)	-2.6 (-3.3 to -1.7)	0.0001
T-score REMS [median (IQR)]	-2.0 (-2.8 to -1.1)	-2.7 (-3.5 to -1.9)	<0.0001
Fragility Score [median (IQR)]	29.9 (25.7 to 36.2)	53.0 (34.2 to 62.5)	<0.0001

By evaluating the capability to discriminate patients with/without fragility fractures, the Fragility Score obtained a value of the ROC area under the curve (AUC) of 0.80, higher than the AUC of the REMS T-score (0.66) and of the T-score DXA (0.64), and the difference was statistically significant (Figure 1).

Figure 1. ROC curve comparison of Fragility Score, REMS and DXA T-score values in the classification of patients with incident fragility fractures.

Furthermore, the correlation between the Fragility Score and the T-score values was low, with Pearson correlation coefficient $r=-0.19$ between Fragility Score and DXA T-score and -0.18 between the Fragility Score and the REMS T-score.

Conclusion: The Fragility Score was found to be an effective tool for the prediction of fracture risk in a population of Caucasian women, with performances superior to those of the T-score values. Therefore, this tool presents a high potential as an effective diagnostic tool for the early identification and subsequent early treatment of bone fragility.

REFERENCES:

- [1] Diez Perez A et al. *Aging Clin Exp Res* 2019; 31(10):1375-1389.
- [2] Pisani P et al. *Measurement* 2017; 101:243-249.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2311

POS0164

THE NEED FOR ANTI-OSTEOPOROTIC INTERVENTION IN POSTMENOPAUSAL WOMEN WITH RHEUMATOID ARTHRITIS BASED ON THE FRACTURE RISK ASSESSMENT

O. Dobrovolskaya¹, A. Feklistov¹, O. Nikitinskaya¹, A. Efremova¹, N. Toroptsova¹. ¹V.A. Nasonova Research Institute of Rheumatology, Department of Osteoporosis, Moscow, Russian Federation

Background: Rheumatoid arthritis (RA) is a chronic disabling disease that is associated with bone loss. Previous studies estimated that approximately