and SMARCA/BRG1 genes may suggest an epigenetic imbalance of chromatin remodelling factors involved in inflammation pathways with a potential role in PsA/psoriasis immunopathogenesis.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.3948

POS0362

INVESTIGATING THE ANKYLOSING SPONDYLITIS-ASSOCIATED REGULATORY SNPS AT THE RUNX3 LOCUS WITH A FUNCTIONAL GENOMICS APPROACH

M. Vecellio¹, P. B. Wordsworth¹, C. J. Cohen¹. ¹University of Oxford, NDORMS, Oxford, United Kingdom

Background: Of the >100 genetic associations with ankylosing spondylitis (AS), RUNX3 transcription factor (TF) involved in diverse immunological processes, is robustly (10^{-15}) associated.¹ The biggest challenge is to understand the mechanism behind this association. We demonstrated the association between AS and the single nucleotide polymorphism (SNP) *rs4648889* located in a 2kb regulatory locus upstream of the *RUNX3* promoter can be explained by allele-specific effects on TF recruitment that alter gene expression, specifically in CD8+ T-cells.² We recently dissected the full plethora of TFs affected by *rs4648889* AS-associated allele showing the NuRD complex and IRF5 differentially bound and having a crucial role in CD8+ T-cells function.³ Nevertheless, other post-GWAS approaches are needed to elucidate the function of *RUNX3* AS-associated SNPs (i.e. the lead SNP *rs6600247*). **Objectives:** The purpose of this work is to define the functional effect of *rs6600247* in CD8+ T-cells and monocytes. In specific, we expect to define the impact of AS-associated allele to TF binding and to evaluate chromosome looping between *rs6600247* and the *RUNX3* promoter.

Methods: The epigenetic landscape of SNP *rs6600247* was defined using Roadmap database. *In vitro* functional studies were performed to characterize the effects of this SNP on TFs binding. Chromosome conformation capture (3C) provided critical functional evidence for looping among AS-associated SNPs and the *RUNX3* promoter.

Results: (1) *In silico* data revealed a c-MYC ChIP-seq peak in GM12878 lymphoblastoid cells overlapping *rs6600247*; (2) Mobility shift assays (EMSA) and WB-EMSA showed reduced DNA/protein binding in the presence of the AS-risk allele in CD14+ monocytes. C-MYC binding-site is disrupted and binding abolished in the presence of the AS-risk allele; (3) 3C experiments indicate low interaction frequency between SNP *rs6600247* and *RUNX3* promoter.

Conclusion: The enhancer upstream the *RUNX3* gene has a plausible functional role in AS, probably by regulating gene transcription and DNA looping. These observations are critically important in defining dysregulated pathways and potential therapeutic drug targets.

REFERENCES:

- [1] IGAS et al. *Nat Genet.* 2013 Jul;45(7):730-8.
- [2] Vecellio M. et al, *Ann Rheum Dis.* 2016 Aug;75(8):1534-40.
- [3] Vecellio et al *Arthritis Rheumatol* 2020. doi: 10.1002/art.41628.

Acknowledgements: We thank Professor Julian Knight for his constructive advices and comments to the present work.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.4098

Adaptive immunity (T cells and B cells) in rheumatic diseases

POS0363

IDENTIFICATION OF MOLECULAR PHENOTYPES AND IMMUNE CELL INFILTRATION IN PSORIATIC ARTHRITIS PATIENTS' SKIN TISSUES BY INTEGRATED BIOINFORMATICS ANALYSIS

T. Cheng^{1,2,3}, S. X. Zhang^{1,2,3}, J. Qiao^{1,2,3}, R. Zhao^{1,2,3}, S. Song^{1,2,3}, Y. Zhang^{1,2,3}, P. Zhao⁴, G. Y. Liu¹, P. F. He⁵, X. Li^{1,2,3}. ¹The Second Hospital of Shanxi Medical University, Department of Rheumatology, Taiyuan, China; ²Shanxi Li Xiaofeng Medical Groups, Department of Rheumatology, Taiyuan, China; ³Ministry of Education, Key Laboratory of Cellular Physiology at Shanxi Medical University, Taiyuan, China; ⁴Shanxi Provincial People's Hospital, Department of Rheumatology, Taiyuan, China; ⁵Shanxi Medical University, Medical Data Sciences, Taiyuan, China

Background: Psoriatic arthritis (PsA) is an inflammatory musculoskeletal disease associated with cutaneous psoriasis¹. Heterogeneity of clinical manifestation often makes differential diagnosis difficult². Thus, the underlying molecular pathogenesis of PsA need to be further studied to diagnose early and ensure optimal management of arthritis and key comorbidities.

Objectives: This research was conducted to identify molecular phenotypes and immune infiltration in the skin tissues of psoriatic arthritis patients according to bioinformatics analysis.

Methods: The mRNA expression profiles of GSE13355 (116 samples), GSE14905 (56 samples) and GSE30999 (162 samples) were obtained from the publicly GEO databases. Non-negative matrix factorization (NMF), functional

enrichment and cibersort algorithm were applied to illustrate the conditions of PsA patients' skin tissues for classification after screening the differentially expressed genes (DEGs) between lesion biopsy and non-lesion biopsy.

Results: Two subsets (Sub1 and Sub2) were identified and validated by NMF typing of 612 detected DEGs (Figure 1a). A total of 54 signature genes (18 in Sub1 and 36 in Sub2) were obtained (Figure 1b). GO and KEGG enrichment analysis showed the signature genes in Sub1 were mainly involved in proliferation and differentiation of immune cells, whereas genes in Sub2 were related to humoral immune response mediated by antimicrobial peptide (Figure 1c,d). Further, immune cell infiltration results revealed Sub2 had higher levels of resting NK cells ($P<0.001$), macrophages M1($P<0.001$), resting mast cells ($P<0.001$) and regulatory T cells ($P<0.001$) but lower concentrations of activated CD4⁺ memory T cells ($P<0.001$), activated NK cells ($P<0.05$), activated dendritic cells($P<0.001$), eosinophils ($P<0.05$) and neutrophil ($P<0.001$) (Figure 1e).

Conclusion: The pathogenesis of psoriatic arthritis is related to both cellular immunity and humoral immunity. It is indispensable to adjust the treatment strategies according to patient's immune status.

REFERENCES:

- [1] Ritchlin CT, Colbert RA, Gladman DD. Psoriatic Arthritis. The New England journal of medicine 2017;376(10):957-70. doi: 10.1056/NEJMra1505557 [published Online First: 2017/03/09].
- [2] Veale DJ, Fearon U. The pathogenesis of psoriatic arthritis. Lancet (London, England) 2018;391(10136):2273-84. doi: 10.1016/s0140-6736(18)30830-4 [published Online First: 2018/06/13].

Figure (a) NMF consensus clustering of 334 PsA patients based on 612 DEGs. (b) Heat map of differentially expressed genes: Each small square represents a gene; the color indicates the expression of the gene. Each column indicates the expression level of genes in a sample; the right side is the gene name. (c,d) Visualization GO enrichment analysis and KEGG pathway enrichment. (e) Violin plot show the fraction of 22 immune cells subpopulations in two subtypes based on CIBERSORT algorithm.

Acknowledgements: This project was supported by National Science Foundation of China (82001740), Open Fund from the Key Laboratory of Cellular Physiology (Shanxi Medical University) (KLCP2019) and Innovation Plan for Postgraduate Education in Shanxi Province (2020BY078).

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2171