population) patients had a titers >340 units/ml of SSB antibodies. The percentage of patients positive for RF antibodies was higher in SSA and/or SSB positive patients ($p = 0.01$).

Table 1. Clinical and demographic features from 647 patients with rheumatoid arthritis and a wish to conceive.

Variable	SSA and SSB anti-bodies [-], n = 622	Positive SSA and/or SSB antibodies [+], n = 25	p-value
Age at inclusion in the cohort (y, SD)	32.3 (4.0)	32.3 (4.4)	0.98
Median disease duration at first visit, years (IQR)	5.2 (2.2 – 9.8)	4.9 (2.1 – 11.4)	0.77
ACPA positive, n (%)	408/614 (66.5)	16 (69.6)	0.75
Rheumatoid Factor positive, n (%)	426/617 (69.0)	23 (92.0)	0.01

Conclusion: Our study shows an overall low prevalence of SSA and SSB autoantibodies in female RA patients with a wish to conceive. Moreover, about half of the patients positive for SSA or SSB had, based on literature, moderate or low titers of these antibodies (2). Especially for RF negative patients, the current advice to test for SSA and SSB antibodies in all RA patients with a wish to conceive should be reconsidered.

REFERENCES:

- [1] Sammaritano LR et al. 2020 American College of Rheumatology Guideline for the Management of Reproductive Health in Rheumatic and Musculoskeletal Diseases. *Arthritis Care Res (Hoboken)*. 2020 Apr; 72(4):461-488.
- [2] Jaeggi E et al. The Importance of the Level of Maternal Anti-Ro/SSA Antibodies as a Prognostic Marker of the Development of Cardiac Neonatal Lupus Erythematosus: A Prospective Study of 186 Antibody-Exposed Fetuses and Infants. *Journal of the American College of Cardiology*. 2010 2010/06/15; 55(24):2778-2784

Disclosure of Interests: Hieronymus TW Smeele: None declared, Luis Fernando Perez-Garcia Consultant of: Yes, Galapagos, J.M.J. Cornette: None declared, Marco WJ Schreurs: None declared, Radboud Dolhain Speakers bureau: Yes UCB, Roche, Abbvie, Genzyme, Novartis, Consultant of: Yes, Galapagos, Grant/research support from: Yes, UCB
DOI: 10.1136/annrheumdis-2021-eular.2435

POS0542

DISEASE ACTIVITY IN PREGNANT WOMEN WITH RHEUMATOID ARTHRITIS: IMPACT ON THE SOLUBLE FMS-LIKE TYROSINE KINASE-1 TO PLACENTAL GROWTH FACTOR RATIO

H.T. Smeele¹, R.I. Neuman^{2,3}, A.H.J. Danser², R. Dolhain¹, W. Visser^{2,3}.
¹Erasmus MC, Rheumatology, Rotterdam, Netherlands; ²Erasmus MC, Department of Internal Medicine; Division of Pharmacology and Vascular Medicine, Rotterdam, Netherlands; ³Erasmus MC, Department of Gynecology and Obstetrics, Rotterdam, Netherlands

Background: An elevated sFlt-1 indicates soluble Fms-like tyrosine kinase-1 (sFlt-1)/ placental growth factor (PlGF) ratio has recently been validated as significant predictor of preeclampsia (PE)(1). However, raised sFlt-1 levels due to inflammation are observed in patients with rheumatoid arthritis (RA)(2). The use of the sFlt-1/PlGF ratio for the prediction of PE has not been evaluated in pregnant women with diseases that are characterized by high levels of inflammation, such as RA

Objectives: We investigated whether sFlt-1 and/or PlGF are altered in pregnant women with RA according to disease activity, and evaluated whether a sFlt-1/PlGF ratio of ≤ 38 could be used to predict the absence of PE in pregnant RA-patients.

Methods: This study was embedded in a nationwide, observational, prospective cohort study on pregnant women with RA (PARA-study). sFlt-1 and PlGF levels were measured, using automated analyzer (Cobas-6000, e-module; Roche-Diagnostics), in the third trimester of pregnancy.

Results: A total of 221 women, aged 21–42 years, were included. Values of sFlt-1, PlGF and sFlt-1/PlGF ratio were not significantly correlated with the DAS28-CRP (Figure 1). CRP correlated weakly with PlGF ($r = -0.14$, $p = 0.03$), while no correlation was found with sFlt-1 or the sFlt-1/PlGF ratio.

Preeclampsia occurred in four out of 214 women with a ratio ≤ 38 (2%) in contrast to three out of seven women with a ratio > 38 (43%), $p < 0.001$ (Table 1). When stratified to a sFlt-1/PlGF ratio of ≤ 38 only 2% of women developed preeclampsia, compared to 43% of the women with a ratio > 38 . The observed sensitivity and specificity for a sFlt-1/PlGF ratio cut-off of ≤ 38 were 42.1% and 98.1%, respectively, with a negative predictive value (NPV) of 98%. All women that developed PE, developed PE >4 weeks after collecting of blood samples in the 3rd trimester, therefore the observed NPV could be even higher.

Table 1. Pregnancy Outcome according to sFlt-1/PlGF ratio with a cut-off value of 38.

Parameter	Ratio ≤ 38	Ratio > 38	P-value
N	214	7	
GA at birth, weeks	39 (38 - 40)	37 (36 - 40)	0.05
<34	2 (1)	0 (0)	1.00
34 - 37	18 (8)	2 (29)	0.12
Male, n (%)	113 (53)	5 (71)	0.45
Birth weight, grams	3420 (2998 - 3800)	2620 (2360 - 2850)	<0.01
Maternal Outcome			
Gestational Hypertension	15 (7)	2 (29)	0.09
Preeclampsia	4 (2)	3 (43)	<0.001
Time to delivery, days	66 (52 - 75)	52 (44 - 66)	0.08
Fetal Outcome			
Birth weight percentile <10	33 (15)	5 (71)	<0.01
Fetal/Neonatal Death	1 (1)	0 (0)	1.00

Data are reported as median (interquartile range) or number (percentage). sFlt-1 indicates soluble Fms-like tyrosine kinase-1; PlGF, placental growth factor; GA, gestational age. Time to delivery is defined as the amount of days between blood sampling and delivery.

Conclusion: Our study shows that in pregnant women with RA, the sFlt-1/PlGF ratio is not altered due to disease activity, and a cut-off of ≤ 38 of this ratio can be used to exclude preeclampsia.

REFERENCES:

- [1] H. Zeisler et al. Predictive Value of the sFlt-1:PlGF Ratio in Women with Suspected Preeclampsia. *N Engl J Med* 2016; 374: 13-22. DOI 10.1056/NEJMoa1414838.
- [2] S. Ballara et al. Paleolog. Raised serum vascular endothelial growth factor levels are associated with destructive change in inflammatory arthritis. *Arthritis Rheum* 2001; 44: 2055-2064. DOI 10.1002/1529-0131(200109)44:9<2055::AID-ART355>3.0.CO;2-2.

Figure 1. Correlations between sFlt-1, PlGF and sFlt-1/PlGF ratio with DAS28-CRP and CRP. Correlation coefficients for sFlt-1 (A-B), PlGF (C-D) and sFlt-1/PlGF ratio (E-F) with DAS28-CRP and CRP.

Acknowledgements: The kits for measurement of sFlt-1/PlGF ratio were a kind gift from Roche Diagnostics, Germany. We thank all participants of the PARA study. Additionally, we extend our gratitude to the laboratory workers, in particular Nadine Davelaar and Priyanka Bangoer, and research assistants for their contribution to the data collection. We thank ReumaNederland (LLP project number: LLP-26) for their financial support.

Disclosure of Interests: Hieronymus TW Smeele: None declared, R.I. Neuman: None declared, A.H.J. Danser: None declared, Radboud Dolhain Speakers bureau: Yes UCB, Roche, Abbvie, Genzyme, Novartis, Consultant of: Yes, Galapagos, Grant/research support from: Yes, UCB, W. Visser: None declared
DOI: 10.1136/annrheumdis-2021-eular.2438