Table 1. The analysis of forced vital capacity (FVC) in KORIL cohort

	V1 (enrollment)	V2 (1-year)	V3 (2-year)	V4 (3-year)
FVC (mL), mean±SD	2549.5 ± 743.3	2479.5 ± 764.4	2435.8 ± 742.8	2405.7 ± 731.2
FVC % of predicted, mean±SD	84.70 ± 16.74	84.7 ± 18.1	84.3 ± 17.6	83.1 ± 19.1
Rate of FVC decline, mL·year ⁻¹ (95% CI)	-	-14 (-49, 21)	-62 (-104, -21)	-
A 10-point decline from V1 in predicted FVC value, n (%)	-	10 (7.1)	16 (13.3)	8 (9.4)
Relative decline of 10% from V1 in predicted FVC value, n (%)	-	17 (12.1)	23 (18.9)	17 (19.3)
Relative decline of 10% from the previous visit in predicted FVC value, n (%)	-	17 (12.1)	17 (14.2)	13 (15.3)

Conclusion: Annually, about 10-15% of patients experience a relative ≥10% decline from the previous visit in FVC predicted value in RA-ILD.

Acknowledgements: This work was supported by the Korea Health Technology R&D Project through the Korea Health Industry Development Institute, funded by the Ministry of Health and Welfare, Republic of Korea (grant no.HI14C1277).

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2810

POS0556

USEFULNESS OF FDG-PET/CT FOR PREDICTING SPONTANEOUS REGRESSION IN MTX ASSOCIATED LYMPHOPROLIFERATIVE DISORDER WITH RHEUMATOID ARTHRITIS

T. Kameda¹, S. Nakashima¹, H. Shimada¹, R. Wakiya¹, M. Mahmoud Fahmy Mansour¹, M. Kato¹, K. Sugihara¹, Y. Ushio¹, H. Dobashi¹. ¹Kagawa University, Department of Internal Medicine, Division of Haematology, Rheumatology and Respiratory Medicine, Faculty of Medicine, Kita-gun, Japan

Background: Recently, there are many reports from Japan about methotrexate associated lymphoproliferative disorder (MTX-LPD). We are investigating the predictive factor of spontaneous regression (SR) in MTX-LPD. On the other hand, FDG-PET/CT is used for diagnosis of LPD including malignant lymphoma. In addition, it was reported that imaging biomarkers such as metabolic tumor volume (MTV) and total lesion glycolysis (TLG) could predict the prognosis of malignant tumor (1, 2). However, there is no report that these imaging biomarkers could predict the SR of MTX-LPD.

Objectives: We investigate the usefulness of FDG-PET/CT for predictive factor of SR in MTX-LPD.

Methods: We enrolled 24 RA patients who diagnosed MTX-LPD and performed FDG-PET/CT from 2005 to 2019. We divided these cases into spontaneous regression cases (SR group; 15 cases) and cases that treated with chemotherapy after MTX discontinuation (CTx group; 9 cases), and compared the difference as follow subjects between two groups; clinical data including histopathological findings, SUVmax to evaluate malignant tumor activity by FDG-PET/CT, MTV and TLG which refer to metabolically active volume of the tumor segmented FDG-PET/CT. In addition, we analyzed cut off levels, sensitivity and specificity using statistical software JMP.

Results: Diffuse large B cell lymphoma (DLBCL) and Hodgkin lymphoma (HL) were 5 and 1 cases in SR group, and 1 and 5 cases in CTx group. In addition, MTV and TLG by FDG-PET/CT was significantly lower in SR group, although SUVmax is no difference between two groups (figure 1). Cut off levels of MTV and TLG were 103.12ml (sensitivity; 88.9%, specificity; 86.7%) and 361.75ml (sensitivity; 88.9%, specificity; 86.7%), respectively.

Conclusion: We suggested that MTV and TLG were useful for predict of SR in MTX-LPD.

REFERENCES:

- [1] Chen HH, Chiu NT, Su WC. et al. Prognostic value of whole-body total lesion glycolysis at pretreatment FDG PET/CT in non-small cell lung cancer. *Radiology*. 2012 Aug;264(2):559-66.
- [2] Chu KP, Murphy JD, La TH. et al. Prognostic value of metabolic tumor volume and velocity in predicting head-and-neck cancer outcomes. *Int J Radiat Oncol Biol Phys*. 2012 Aug 1;83(5):1521-7.

Figure 1. Comparison of the level of MTV(a) and TLG (b).

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2812

POS0557

THE RELATION BETWEEN INVOLVEMENT OF INDIVIDUAL JOINTS OF THE UPPER EXTREMITY AND GRIP FORCE IN EARLY RHEUMATOID ARTHRITIS

M. Rydholm^{1,2}, A. Sharma², I. Wikström^{1,2}, L. T. H. Jacobsson^{2,3}, C. Turesson^{1,2}.

¹Skåne University Hospital, Department of Rheumatology, Malmö, Sweden;

²Lund University, Rheumatology, Department of Clinical Sciences, Malmö, Sweden;

³Gothenburg University, Department of Rheumatology and Inflammation Research, Sahlgrenska Academy, Gothenburg, Sweden

Background: Rheumatoid arthritis (RA) has a major impact on hand function in many patients.

Objectives: The objective of this study was to investigate the relation between swelling and tenderness of individual joints in the upper extremities and grip force in patients with early RA.

Methods: An inception cohort of patients with early RA (symptom duration ≤12 months), recruited in 1995-2005, was investigated and followed in a structured program. All patients were examined by the same rheumatologist according to a structured protocol, and swelling and tenderness of individual joints was recorded. Grip force (Newton, N) was measured using the electronic instrument Grippit (AB Detektor, Gothenburg, Sweden). Average grip force values of each hand were evaluated and compared to the expected, based on age- and sex-specific reference values from the literature (1). Grip force was expressed as % of expected values. Linear regression analyses were used to assess the relation between upper extremity joint involvement in individual joints and grip force, using the presence/absence of synovitis/tenderness in each joint as covariates. For the multivariate analysis, covariates with $p < 0.10$ in the univariate analyses were selected. In cases with major collinearity (Spearman's $r > 0.3$), the covariate with the stronger association in the univariate analysis was included.

Results: A total of 215 patients with early RA (71 % women; mean age 60 years) were investigated. The median symptom duration at inclusion was 7 months; interquartile range 5-10. The median 28 joint counts for swollen joints was 7 (interquartile range, IQR 5-11) and for tender joints 4 (IQR 1-9). In univariate analyses, swelling of the wrist, each metacarpophalangeal (MCP) joint and proximal interphalangeal (PIP) joints III and V, as well as tenderness of the elbow, wrist and each MCP and PIP joint were associated with significantly reduced grip force in the right hand. In the left extremity, corresponding associations were observed for swelling of the wrist, each MCP joint and PIP joints I and V, and for tenderness of the wrist and each MCP and PIP joint. In multivariate analysis, wrist synovitis was associated with reduced grip force (β -11.0 and -14.3 % of expected, respectively). There were also independent significant associations for tenderness of MCP IV and the elbow with reduced grip force in the right hand, and for swelling of MCP I and MCP V with reduced grip force in the left hand (Table 1).