Objectives: We describe and analyze *Listeria*-related demographics and clinical features to determine the predisposing conditions for severe infections in an immunodepressed population by rheumatic diseases.

Methods: Descriptive Observational Study. A retrospective analysis of 143 patients were performed affected by listeriosis, with positive isolation of *Listeria monocytogenes* from blood, treated in the H.U. Virgen del Rocío (Seville- Spain) between 2003-2019. Of them 9 were rheumatic patients. The type of clinical manifestation was analyzed, paying special attention to the characteristics associated with patients with neurological complications or unfavorable outcome (death and / or abortion in pregnant women), immunosuppression (associated with cancer or rheumatic disease) was assessed as independent variables, chronic diseases (Hypertension, Diabetes Mellitus, dyslipidemia, COPD, Renal Insufficiency and Ischemic Heart Disease) as well as other baseline characteristics of the patient. (age, sex, pregnancy) and their toxic habits (tobacco and alcohol).

Results: The sample includes a similar proportion of men (70 cases) and women (73 cases), of all ages. Of the total patients, most (85%) required hospital admission, with a duration median (non-parametric data) of 11 days. 78% of the cases admitted showed a favorable evolution. However, 15.4% resulted in death and 5.6% in abortion. This percentage of abortions represented 29% of the total pregnant women admitted. Of all the patients admitted, a third (33%) were immunocompromised, including patients with cancer (79%) and rheumatic diseases (21%). Include lupus (33%), RA (22%), APs (11%), polymyalgia rheumatica (11%), panuveitis (11%) and ANCA vasculitis MPO specificity (11%). All of them required admission although the majority showed a favorable evolution, except one of the patient, which resulted in death, in which case in addition to lupus he presented with prostate cancer. Regarding the baseline treatment of these patients, 7 underwent treatment with synthetic DMARDs and three with biological DMARDs (1 Adalimumab, 1 Infliximab and 1 Rituximab). As a result of the *Listeria* infection, most of them had fever or digestive symptoms and two of them experienced neurological manifestations (meningoencephalitis). None of these last two (with lupus and RA) had biological DMARDs.

Conclusion: Listeriosis is an uncommon but potentially serious infection usually in older people, pregnant women and immunocompromised patients. In our sample, 33% of the patients were immunocompromised. Of the 9 patients affected by *Listeria* with rheumatic disease we find a death for meningoencephalitis. Given the impact of this infection in immunosuppressed patients should pay attention in our patients with fever and neurological manifestations.

REFERENCE:

- [1] Eleftherios Mylonakis et al. A Case Series and Review of 222 Cases. *Medicine* 2002; 81: 260-269.
- [2] Alcoba Lez M et al. Meningitis por *Listeria monocytogenes* en el adulto en España. Presentación de 10 casos y revisión de la literatura. *Rev Clin Esp* 2002; 202 (12): 638-643.
- [3] Eleftherios Mylonakis et al. Central Nervous System Infection with *Listeria monocytogenes*. 33 Years' Experience at a General Hospital and Review of 776 Episodes from the Literature. *Medicine* 1998; 77: 313-336.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.4282

Rheumatoid arthritis - comorbidity and clinical aspects - I

OP0098

POLYPHARMACY IS ASSOCIATED WITH A POORER TREATMENT RESPONSE AND INCREASED RISK OF ADVERSE EVENTS IN EARLY RHEUMATOID ARTHRITIS

S. Benamar¹, C. Lukas², C. Daien², C. Gaujoux-Viala³, L. Gossec⁴, A. C. Rat⁵, B. Combe², J. Morel³. ¹CHU Montpellier, Rheumatology, Montpellier, France; ²University Hospital of Montpellier, Rheumatology, Montpellier, France; ³University Hospital of Nîmes, Rheumatology, Nîmes, France; ⁴La Pitié-Salpêtrière Hospital, Rheumatology, Paris, France; ⁵CHU Caen, Rheumatology, Caen, France

Background: Polypharmacy is steadily increasing in patients with rheumatoid arthritis (RA). They may interfere with treatment response and the occurrence of serious adverse events. Medications taken by a patient may reflect active comorbidities, whereas comorbidity indices usually used include past or current diseases.

Objectives: To evaluate whether polypharmacy is associated with treatment response and adverse events in an early RA cohort and to establish whether polypharmacy could represent a substitute of comorbidities.

Methods: We used data from the French cohort ESPOIR, including 813 patients with early onset arthritis. Patients included in the current study had to start their first disease modifying anti-rheumatic drug (DMARD) within 24 months of inclusion in the cohort. Disease activity data were collected at one, five and ten years from the initiation of the first DMARD. For each patient, treatments were collected at baseline and at five years. Medications count included all specialties other than background RA therapy,

analgesics/NSAIDs and topicals. Polypharmacy was defined as a categorical variable based on the median and tertiles of distribution in the cohort. Treatment response was assessed by achieving DAS28 ESR remission (REM) at 1 year, 5 years and 10 years from the initiation of the first DMARD. The occurrence of severe adverse events (SAE) was measured by the occurrence of severe infection, hospitalization, or death during the 10-year follow-up. The association between patient's characteristics and achievement of REM and occurrence of SAE were tested in univariate analysis. A logistic regression model was used to evaluate associations between polypharmacy and REM at 1 year, 5 years and 10 years (we used baseline polypharmacy for the 1-year analysis and five years polypharmacy for the 5- and 10-years analyses). Multivariate adjustment was made for age, sex, BMI, duration of disease, initial DAS28 ESR, initial HAQ, smoking status, rheumatic disease comorbidity index (RDCI).

Results: The proportion of patients who achieved REM one year after the initiation of the first DMARD was 32.1% in the polypharmacy according to the median group (patients taken ≥ 2 medication) versus 67.9% in the non-polypharmacy group ($p=0.07$). At 5 years after the first DMARD, the proportion of patients with REM was 45.0% in the polypharmacy group versus 56.3% in the non-polypharmacy group ($p=0.03$). At 10 years the proportion of patients with REM was 32.5% in the polypharmacy group versus 67.5% ($p=0.06$). Patients who take greater or equal to 2 medications had a 40% lower probability of achieving REM (OR = 0.60 [0.38-0.94] $p = 0.03$) at 5 years from the first DMARD (if RDCI index was not included in the model). At 10 years, patients receiving multiple medications had a 43% lower probability of achieving REM (OR = 0.57 [0.34-0.94] $p = 0.02$). SAE incidence was 61 per 1000 patient-years. For patients who developed SAE all causes 71.4% where in the polypharmacy group versus 57.8% were in the non-polypharmacy group ($p = 0.03$; univariate analysis). These results are no longer significant after adjustment for comorbidities indices.

Conclusion: In this early RA cohort, polypharmacy is associated with a poorer treatment response and increased risk of adverse events. Polypharmacy may represent a good substitute of comorbidities for epidemiological studies.

Acknowledgements: We are grateful to Nathalie Rincheval (Montpellier) who did expert monitoring and data management and all the investigators who recruited and followed the patients (F. Berenbaum, Paris-Saint Antoine; MC. Boissier, Paris-Bobigny; A. Cantagrel, Toulouse; B. Combe, Montpellier; M. Dougados, Paris-Cochin; P. Fardellone and P. Boumier, Amiens; B. Fautrel, Paris-La Pitié; RM. Flipo, Lille; Ph. Goupille, Tours; F. Liote, Paris- Lariboisière; O. Vittecoq, Rouen; X. Mariette, Paris-Bicêtre; P. Dieude, Paris Bichat; A. Saraux, Brest; T. Schaefferbeke, Bordeaux; and J. Sibilia, Strasbourg).

The work reported on in the manuscript did not benefit from any financial support. The ESPOIR cohort is sponsored by the French Society for Rheumatology. An unrestricted grant from Merck Sharp and Dohme (MSD) was allocated for the first 5 years. Two additional grants from INSERM were obtained to support part of the biological database. Pfizer, Abbvie, Lilly and more recently Fresenius and Biogen also supported the ESPOIR cohort.

Disclosure of Interests: Soraya Benamar: None declared, Cédric Lukas Speakers bureau: Abbvie, Amgen, Janssen, Lilly, MSD, Novartis, Pfizer, Roche-Chugai, UCB, Consultant of: Abbvie, Amgen, Janssen, Lilly, MSD, Novartis, Pfizer, Roche-Chugai, UCB, Grant/research support from: Pfizer, Novartis and Roche-Chugai, Claire Daien Speakers bureau: AbbVie, Abivax, BMS, MSD, Roche, Chugai, Novartis, Pfizer, Sandoz, Lilly, Consultant of: AbbVie, Abivax, BMS, MSD, Roche, Chugai, Novartis, Pfizer, Sandoz, Lilly, Cécile Gaujoux-Viala Speakers bureau: AbbVie, BMS, Celgene, Janssen, Medac, MSD, Nordic Pharma, Novartis, Pfizer, Sanofi, Roche-Chugai, UCB, Consultant of: AbbVie, BMS, Celgene, Janssen, Medac, MSD, Nordic Pharma, Novartis, Pfizer, Sanofi, Roche-Chugai, UCB, Grant/research support from: Pfizer, Laure Gossec Speakers bureau: AbbVie, Amgen, Biogen, Celgene, Janssen, Lilly, Novartis, Pfizer, Sandoz, Sanofi-Aventis et UCB, Anne-Christine Rat Speakers bureau: Pfizer, Lilly, Consultant of: Pfizer, Lilly, Bernard Combe Speakers bureau: AbbVie; Bristol-Myers Squibb; Gilead; Janssen; Lilly; Merck; Novartis; Pfizer; Roche-Chugai; and Sanofi; Consultant of: AbbVie; Bristol-Myers Squibb; Gilead; Janssen; Lilly; Merck; Novartis; Pfizer; Roche-Chugai; and Sanofi; Grant/research support from: Novartis, Pfizer, and Roche-Chugai; Jacques Morel Speakers bureau: AbbVie, BMS, Lilly, Médac, MSD, Nordic Pharma, Pfizer, UCB, Consultant of: AbbVie, BMS, Lilly, Médac, MSD, Nordic Pharma, Pfizer, UCB, Grant/research support from: BMS, Pfizer

DOI: 10.1136/annrheumdis-2021-eular.3213

OP0099

EPIDEMIOLOGY AND MORTALITY OF RA-ASSOCIATED INTERSTITIAL LUNG DISEASE: DATA FROM A FRENCH ADMINISTRATIVE HEALTHCARE DATABASE

P.A. Juge¹, L. Wemeau Stervinou², S. Ottaviani³, G. Desjeux⁴, J. Zhuo⁵, B. Bregman⁶, V. Vannier-Moreau⁷, R. M. Flipo⁸, B. Crestani⁹, P. Dieudé³. ¹Université de Paris, AP-HP, Hôpital Bichat Claude-Bernard, Department of Rheumatology, Paris, France; ²CHU Lille, Department of Pulmonology, Lille, France; ³Assistance Publique-Hôpitaux de Paris, Hôpital Bichat-Claude-Bernard,