Table 1. Median of mean CRP^a up to W48 by response subgroups

	Tofacitinib 11 mg QD + MTX		Tofacitinib 11 mg QD + PBO	
Mean CRP, ^a median (IQR) [n]	>BL–W24	>W24–48	>BL–W24	>W24–48
HAQ-DI CRFP	2.84 (1.15–7.30) [45]	2.30 (0.82–4.75) [46]	1.45 (0.77–4.42) [56]	2.28 (0.53–7.28) [56]
HAQ-DI non-CRFP	2.81 (1.09–6.19) [195]	2.91 (1.19–5.84) [195]	2.26 (0.98–4.63) [176]	2.47 (1.13–5.53) [178]
DAS28-4(ESR) remission/LDA	2.48 (1.05–4.95) [126]	2.46 (1.07–4.76) [127]	1.70 (0.89–4.14) [115]	1.95 (0.81–3.82) [117]
DAS28-4(ESR) moderate/high disease activity	3.56 (1.17–7.13) [107]	3.58 (1.36–8.33) [107]	2.60 (0.87–5.16) [115]	2.68 (1.34–8.23) [115]

^aMean CRP was calculated as the average CRP value during each time period (>BL–W24 or >W24–48)

CRP, C-reactive protein; DAS28-4(ESR), Disease Activity Score in 28 joints, erythrocyte sedimentation rate; HAQ-DI, Health Assessment Questionnaire-Disability Index; IQR, interquartile range; LDA, low disease activity; MTX, methotrexate; n, number of pts meeting assessment criteria; QD, once daily

Figure. Cumulative probability plot at W48 for a) ACR-N; b) ΔHAQ-DI score

Endpoints were assessed using the full analysis set (defined as all pts receiving ≥1 dose of study drug with ≥1 value post-baseline), with no imputation for missing values.
ACR-N, American College of Rheumatology response, where N represents the minimum amount of improvement expressed by each pt in % ΔTJC28, % ΔSJC28, and the median of % Δ in PGA, Pain, HAQ-DI and CRP; CRP, C-reactive protein; HAQ-DI, Health Assessment Questionnaire-Disability Index; MR, modified release; MTX, methotrexate; PGA, Physician Global Assessment; TJC28, Patient Global Assessment; QD, once daily; SJC28, swollen joint count in 28 joints; TJC28, tender joint count in 28 joints

Conclusion: Although clinical/functional responses were generally similar between treatment groups, numerical improvements were seen for some efficacy endpoints with tofacitinib + MTX vs tofacitinib + PBO. A numerically higher CRFP rate may be associated with higher DAS28-4(ESR) disease activity. CRP changes up to W48 may not trend with CRFP status.

REFERENCES:

[1] Cohen et al. *Lancet Rheumatol* 2019; 1: E23-34.

Acknowledgements: Study sponsored by Pfizer Inc. Medical writing support was provided by Anthony G McCluskey, CMC Connect, and funded by Pfizer Inc.

Disclosure of Interests: Stanley B. Cohen Consultant of: AbbVie, Eli Lilly, Genentech, Gilead Sciences, Pfizer Inc, Grant/research support from: AbbVie, Eli Lilly, Genentech, Gilead Sciences, Pfizer Inc, Yi-Hsing Chen Grant/research support from: Bristol-Myers Squibb, GlaxoSmithKline, Pfizer Inc, Naonobu Sugiyama Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Jose Luis Rivas Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Annette Diehl Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Tatjana Lukic Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Jerome Paulissen Consultant of: Pfizer Inc, Haiyun Fan Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Tomohiro Hirose Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Edward Keystone Speakers bureau: AbbVie, Amgen, F. Hoffman-La Roche, Janssen, Merck, Novartis, Pfizer Inc, Sanofi

Genzyme, Consultant of: AbbVie, Amgen, Bristol-Myers Squibb, Celltrion, Eli Lilly, F. Hoffman-La Roche, Gilead Sciences, Janssen, Merck, Myriad Autoimmune, Pfizer Inc, Sandoz, Sanofi Genzyme, Samsung Bioepis, Grant/research support from: Amgen, Merck, Pfizer Inc, PuraPharm

DOI: 10.1136/annrheumdis-2021-eular.358

POS0652

SEX DIFFERENCES IN THE EFFICACY AND SAFETY OF TOFACITINIB IN RHEUMATOID ARTHRITIS PATIENTS: A POST HOC ANALYSIS OF PHASE 3 AND LONG-TERM EXTENSION TRIALS

H. N. Jones¹, V. Strand², H. Schulze-Koops³, E. Mysler⁴, C. Kinch⁵, D. C. Gruben⁶, R. Germino⁷, C. A. Connell⁸, L. Eder⁹. ¹University of Alberta, Division of Rheumatology, Edmonton, AB, Canada; ²Stanford University, Division of Immunology/Rheumatology, Palo Alto, CA, United States of America; ³University of Munich, Division of Rheumatology and Clinical Immunology, Department of Internal Medicine IV, Munich, Germany; ⁴Organización Médica de Investigación, Buenos Aires, Argentina; ⁵Pfizer Canada ULC, Inflammation and Immunology, Kirkland, QC, Canada; ⁶Pfizer Inc, Inflammation and Immunology, Groton, CT, United States of America; ⁷Pfizer Inc, Inflammation and Immunology, New York, NY, United States of America; ⁸Pfizer Inc, Clinical Development and Operations, Groton, CT, United States of America; ⁹University of Toronto, Women's College Research Institute, Department of Medicine, Toronto, ON, Canada

Background: Differences in efficacy outcomes favouring males vs females with rheumatoid arthritis (RA) have been reported with conventional synthetic (cs) disease-modifying antirheumatic drugs (DMARDs) and tumour necrosis factor inhibitors; results with Janus kinase inhibitors are less clear.

Objectives: To assess the impact of sex on efficacy, safety and persistence in tofacitinib RA clinical trials.

Methods: Efficacy and safety analyses included data pooled from Phase (P)3 randomised controlled trials (RCTs) of patients (pts) with RA and an inadequate response (IR) to methotrexate (NCT00847613; NCT00853385) or ≥1 DMARD (NCT00856544) who received tofacitinib 5 or 10 mg twice daily (BID), adalimumab (ADA) 40 mg Q2W or placebo (PBO), with background csDMARDs. Persistence analyses of pts receiving tofacitinib 5 or 10 mg BID ± csDMARDs used data pooled from two long-term extension trials (NCT00661661; NCT00413699). Efficacy outcomes to Month (M)12 included: ACR20/50/70 responses, changes from baseline (Δ; BL) in DAS28-4(ESR), CDAI, HAQ-DI and FACIT-F, and DAS28-4(ESR) remission (<2.6). Safety was evaluated to M24 for tofacitinib and ADA. Kaplan-Meier persistence analysis estimated time to discontinuation.

Results: 2265 pts were included from P3 RCTs. Demographics and BL characteristics were comparable across sexes and treatments. Tofacitinib or ADA vs PBO generally led to significantly higher ACR20/50/70 responses in both sexes through M6. To M12, ACR20/50/70 responses were broadly comparable across active treatments and between sexes, with significant differences favouring males at some time points, including M3 (Figure 1). Statistically significant differences favouring males vs females were observed in DAS28-4(ESR) remission rates at most time points, including M3 (Figure 1); a similar trend was observed for ΔDAS28-4(ESR). ΔCDAI, ΔHAQ-DI and ΔFACIT-F significantly favoured males vs females receiving tofacitinib 5mg BID at most time points, while ΔHAQ-DI and ΔFACIT-F tended to favour females receiving tofacitinib 10mg BID. Rates of adverse events (AEs), serious AEs (SAEs), severe AEs and discontinuations due to AEs were slightly higher in females vs males with tofacitinib 5mg BID; this was generally reversed with tofacitinib 10mg BID and ADA (Table 1). AEs of special interest (AESI) were comparable between sexes with tofacitinib and ADA, although low event numbers limit interpretation (Table 1). Time to all-cause discontinuation and discontinuation due to AEs/lack of efficacy with tofacitinib 5mg BID was similar between sexes. Numerical differences favouring females vs males were observed for time to all-cause discontinuation and discontinuation due to AEs with tofacitinib 10mg BID.

Table 1. Safety summary to M24 in pooled DMARD-IR P3 RCTs

	Tofacitinib 5mg BID		Tofacitinib 10mg BID		ADA	
Pts with events, n (%)	Females (N=707)	Males (N=133)	Females (N=698)	Males (N=137)	Females (N=162)	Males (N=42)
AEs	562 (79.5)	85 (63.9)	529 (75.8)	107 (78.1)	119 (73.5)	30 (71.4)
SAEs	107 (15.1)	17 (12.8)	71 (10.2)	24 (17.5)	13 (8.0)	6 (14.3)
Severe AEs	86 (12.2)	12 (9.0)	55 (7.9)	22 (16.1)	14 (8.6)	5 (11.9)
Discontinuations due to AEs	87 (12.3)	10 (7.5)	88 (12.6)	10 (7.3)	17 (10.5)	5 (11.9)
Death	6 (0.8)	4 (3.0)	0	3 (2.2)	1 (0.6)	2 (4.8)
AESI						
Serious infections	28 (4.0)	6 (4.5)	27 (3.9)	6 (4.4)	2 (1.2)	1 (2.4)
All HZ (non-serious/serious)	35 (5.0)	7 (5.3)	43 (6.2)	5 (3.6)	2 (1.2)	3 (7.1)
MACE	5 (0.7)	0	2 (0.3)	3 (2.2)	0	3 (7.1)
Malignancies (excl. NMSC)	7 (1.0)	1 (0.8)	9 (1.3)	1 (0.7)	0	1 (2.4)
NMSC	2 (0.3)	5 (3.8)	4 (0.6)	2 (1.5)	1 (0.6)	1 (2.4)
Venous thromboembolism	3 (0.4)	0	3 (0.4)	1 (0.7)	0	0

HZ, herpes zoster; MACE, major adverse cardiovascular events; NMSC, non-melanoma skin cancer

Figure. Proportion of pts achieving ACR20/50/70 and DAS28-(ESR) remission at M3, stratified by sex and treatment group (P3 RCTs)

*p<0.05, **p<0.001

Pts from ORAL Standard who were found to be non-compliant with study procedures were excluded; PBO data not presented. ACR20/50/70, American College of Rheumatology ≥20/50/70% response criteria; DAS28-(ESR), Disease Activity Score in 28 joints, erythrocyte sedimentation rate; n, number of responders; N, number of patients assessed at each visit

Conclusion: Efficacy outcomes with tofacitinib and ADA were generally higher in males and comparable in females vs previously reported mixed population response rates for advanced therapies. Safety findings did not reveal a consistent pattern between sexes. Tofacitinib persistence was similar between sexes.

Acknowledgements: Study sponsored by Pfizer Inc. Medical writing support was provided by Christina Viegelmann, CMC Connect, and funded by Pfizer Inc.

Disclosure of interests: H Niall Jones Consultant of: Pfizer Inc, Vibeke Strand Consultant of: AbbVie, Amgen, Arena, AstraZeneca, Bayer, Boehringer Ingelheim, Bristol-Myers Squibb, Celtrion, Corrona, Eli Lilly, Galapagos, Genentech/Roche, Gilead, GlaxoSmithKline, Ichnos, Inmedix, Janssen, Kiniksa, Merck, Myriad Genetics, Novartis, Pfizer Inc, Regeneron, Samsung, Sandoz, Sanofi, Scipher, SetPoint Medical, UCB, Hendrik Schulze-Koops Consultant of: AbbVie, Amgen, Biogen, Bristol-Myers Squibb, Celgene, Eli Lilly, Gilead Sciences, Hexal Sandoz, Hospira, Janssen-Cilag, MSD, Novartis, Pfizer Inc, Roche, UCB, Grant/research support from: Novartis, Pfizer Inc, Eduardo Mysler Speakers bureau: AbbVie, Bristol-Myers Squibb, Eli Lilly, Janssen, Pfizer Inc, Roche, Sanofi, Grant/research support from: Eli Lilly, Pfizer Inc, Roche, Cassandra Kinch Shareholder of: Pfizer Canada ULC, Employee of: Pfizer Canada ULC, David C Gruben Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Rebecca Germino Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Carol A. Connell Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Lihi Eder Speakers bureau: AbbVie, UCB, Consultant of: AbbVie, Celgene, Eli Lilly, Novartis, Pfizer Inc, UCB, Grant/research support from: AbbVie, Eli Lilly, Pfizer Inc, UCB

DOI: 10.1136/annrheumdis-2021-eular.359

POS0653

IMPACT OF UPADACITINIB OR ADALIMUMAB AS INITIAL THERAPY ON THE ACHIEVEMENT OF 48-WEEK TREATMENT GOALS IN PATIENTS WITH RHEUMATOID ARTHRITIS AND INADEQUATE RESPONSE TO METHOTREXATE: POST HOC ANALYSIS OF A PHASE 3 STUDY

E. Mysler¹, Y. Tanaka², A. Kavanaugh³, D. Aletaha⁴, P. C. Taylor⁵, I. H. Song⁶, T. Shaw⁶, Y. Song⁶, R. Demasi⁶, M. Ali⁶, R. Fleischmann⁷. ¹Organización Médica de Investigación, Rheumatology, Buenos Aires, Argentina; ²The First Department of Internal Medicine, University of Occupational and Environmental Health, Kitakyushu, Japan; ³Division of Rheumatology, Allergy, & Immunology, University of California San Diego Medical School, San Diego, United States of America; ⁴Division of Rheumatology, Department of Medicine 3, Medical University of Vienna, Vienna, Austria; ⁵Botnar Research Centre, Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, United Kingdom; ⁶AbbVie Inc., Rheumatology, North Chicago, United States of America; ⁷Department of Medicine, University of Texas Southwestern Medical Center, Metroplex Clinical Research Center, Dallas, United States of America

Background: In the randomized, double-blinded, Phase 3 SELECT-COMPARE study, upadacitinib (UPA) + MTX demonstrated greater clinical and functional responses vs adalimumab (ADA) + MTX in patients (pts) with RA and inadequate responses to MTX.^{1,2} Pts with insufficient response to initial therapy were switched from UPA to ADA (and vice versa) according to treat-to-target (T2T) principles.

Objectives: We analyzed 1-year treatment outcomes in SELECT-COMPARE according to initial randomization group, regardless of whether pts subsequently switched therapy.

Methods: Pts initially randomized to UPA 15 mg once daily (QD) or ADA 40 mg every other week (EOW; both + MTX) for up to 48 weeks in SELECT-COMPARE were included in the analysis. As per the protocol-directed rescue strategy, pts

experiencing <20% improvement in tender or swollen joint counts at Week 14, 18, or 22, or Clinical Disease Activity Index (CDAI) >10 at Week 26, were switched from UPA to ADA or ADA to UPA in a blinded fashion. Efficacy outcomes included CDAI remission (≤2.8) and low disease activity (LDA; ≤10), DAS of 28 joints using CRP (DAS28(CRP)) <2.6 and ≤3.2, and a composite of “deep response” (CDAI remission, HAQ-Disability Index <0.5, and pain score <20). Data are presented and attributed to initial randomized group (UPA or ADA) regardless of any subsequent switch in therapy. Time-averaged response rates were calculated as area under the curve of response rate standardized by 48 weeks. The proportions of pts who maintained Week 26 responses through 6 months of follow-up are also reported.

Results: This analysis included 651 pts initially randomized to UPA (of whom 245 switched to ADA) and 327 pts initially randomized to ADA (of whom 157 switched to UPA). Baseline characteristics including age, sex, and BMI were generally well balanced between randomized groups. At Week 48, similar proportions of pts initially randomized to UPA or ADA therapy achieved CDAI remission/LDA (27.6%/61.9% vs 24.8%/59.0%) and DAS28(CRP) <2.6/≤3.2 (45.0%/60.2% vs 43.7%/59.0%) (Figure 1). However, a small but significantly greater proportion of pts achieved a deep response with initial UPA vs initial ADA therapy (17.8% vs 12.8%; p<0.05). In addition, time-averaged response rates over 48 weeks were higher for initial UPA vs initial ADA therapy across efficacy outcomes. Similar trends were observed for other outcomes. Additionally, similar proportions of pts maintained Week 26 responses with initial UPA vs initial ADA therapy based on CDAI remission/LDA and DAS28(CRP) <2.6/≤3.2 during 6-month follow-up (Table 1).

Conclusion: Using a stringent T2T approach to RA management, rates of LDA

Figure 1. Achievement of CDAI (A) and DAS28(CRP) (B) responses corresponding with time-averaged response rates over 48 weeks by initial therapy with UPA or ADA in SELECT-COMPARE

or remission at 1 year were similar, regardless of whether pts were initially randomized to UPA or ADA. However, initial UPA therapy led to more frequent deep responses and higher time-averaged response rates vs initial ADA therapy.

REFERENCES:

- [1] Fleischmann R, et al. Arthritis Rheumatol 2019;71:1788–800.
- [2] Fleischmann R, et al. Ann Rheum Dis 2019;78:1454–62.