Table 1.

	Overall	PMR	PMR+GCA
n (%)	243	178 (73.3)	65 (26.7)
Female sex (%)	146 (60.1)	98 (55.1)	48 (73.8)
Shoulder girdle pain (%)	236 (97.1)	174 (97.8)	62 (95.4)
Pelvic girdle pain (%)	174 (71.6)	127 (71.3)	47 (72.3)
Inflammatory back pain (%)			
No	107 (44.0)	83 (46.6)	24 (36.9)
Yes	106 (43.6)	70 (39.3)	36 (55.4)
Lower limb pain (%)			
No	87 (35.8)	61 (34.3)	26 (40.0)
Yes	81 (33.3)	68 (38.2)	13 (20.0)
Weight loss (%)	112 (46.1)	78 (43.8)	34 (52.3)
CRP (mg/l) (median [IQR])	46.0 [19.0, 77.7]	44.0 [16.9, 74.2]	52.0 [27.9, 85.0]
ESR (mm/h) (mean (SD))	65.2 (30.3)	62.7 (30.2)	72.3 (29.7)
Hemoglobin (g/dl) (mean (SD))	12.1 (1.5)	12.2 (1.5)	11.7 (1.6)
Thrombocytes (1e+09/ml) (mean (SD))	341.9 (106.3)	323.9 (103.2)	375.8 (104.6)

In the univariable analyses the following factors were most strongly associated with vasculitic PET findings: female sex (OR 2.31, CI 1.17-4.58), inflammatory back pain (OR 2.73, CI 1.32-5.64), temperature $>37^{\circ}$ (OR 1.83, CI 0.90-3.7), weight loss (OR 1.83, CI 0.96-3.51), thrombocytosis (i.e., patients with a thrombocyte count 1 SD above mean have an OR of 1.51, CI 1.05-2.18), anemia (i.e., 1 g/dl decrease in Hb below mean corresponds to an OR of 1.25, CI 1.00-1.56). Patients with lower limb pain were less likely to have vasculitis (OR 0.43, CI 0.19-0.95). The estimated ORs were very similar in the multivariable model although the 95% CIs became wider.

Conclusion: Although the prevalence across published studies showed substantial variation, 6 out of 13 studies reported a prevalence of silent GCA in 18-40% of all PMR patients. The exploratory analysis of the collected IPD identified female sex, inflammatory back pain, fever, weight loss, absence of lower leg pain, thrombocytosis and anemia as factors associated with LV-GCA. These findings should be validated in future prospective cohort studies. The presence or absence of these factors may further aid in diagnosing LV-GCA in PMR patients.

REFERENCES:

- [1] Buttgeriet F, Dejaco C, Matteson EL, Dasgupta B. Polymyalgia Rheumatica and Giant Cell Arteritis: A Systematic Review. *JAMA*. 2016 Jun 14;315(22):2442-58.

Acknowledgements: The study is funded by the "Schweizerische Stiftung für die Erforschung der Muskelkrankheiten (SSEM)".

Disclosure of Interests: Daniele Silvio Gozzoli: None declared, Andrea Hemmig: None declared, Lars Hemkens: None declared, Laura Werlen: None declared, Hannah Ewald: None declared, Christoph Berger: None declared, Diego Kyburz Grant/research support from: DK reports personal fees from Abbvie, Gilead, Lilly, Novartis and Pfizer, outside of the submitted work, Stephan Imfeld: None declared, Markus Aschwanden: None declared, Mihaela Stegert: None declared, Dario Camellino: None declared, Marco Amedeo Cimmino: None declared, Corrado Campochiaro Grant/research support from: personal fees from Roche, Alessandro Tomelleri: None declared, Liesbet Henckaerts: None declared, Daniel Blockmans Speakers bureau: Paid speaker for Roche, Consultant of: Paid consultant for Roche, Patricia Moya: None declared, Hector Corominas: None declared, Russell Buchanan: None declared, Claire Owen Speakers bureau: CO has received speaking honoraria from Roche, Janssen, Novartis and Pfizer, and meeting sponsorship from Roche, UCB and Janssen, Yannick van Sleen: None declared, Elisabeth Brouwer Speakers bureau: E. Brouwer as an employee of the UMCG received speaker fees and consulting fees from Roche in 2017, 2018 which were paid to the UMCG, Consultant of: E. Brouwer as an employee of the UMCG received speaker fees and consulting fees from Roche in 2017, 2018 which were paid to the UMCG, Hiroyuki Ymashita: None declared, Thomas Daikeler: None declared

DOI: 10.1136/annrheumdis-2021-eular.1803

POS0807

RETINAL VESSEL CALIBERS AS A NON-INVASIVE BIOMARKER OF INFLAMMATORY BURDEN IN PRIMARY SYSTEMIC VASCULITIS

O. Argyropoulou¹, E. Aissopou¹, A. Argyris¹, A. Goules¹, C. Mavragani², N. Tentolouris³, P. Sfrikakis³, P. Vlachoyiannopoulos¹, A. Tzioufas¹, A. Protopogerou¹. ¹School of Medicine, National and Kapodistrian University of Athens, Pathophysiology, Athens, Greece; ²School of Medicine, National and Kapodistrian University of Athens, Physiology, Athens, Greece; ³Medical School, National and Kapodistrian University of Athens, Athens, First Department of Propaedeutic Internal Medicine, Athens, Greece

Background: Primary Systemic Vasculitides (PSV) constitute a heterogeneous group of rare and potentially life-threatening autoimmune diseases, characterized by a varying degree of inflammatory response, leading to local or generalized vascular disease. Vessel involvement accounts for the micro- and macrovascular complications of the disease, along with the classic risk factors including, among others age and chronic use of steroids (1). Early identification of high-risk

patients for cardiovascular disease (CVD) development and the contribution of inflammation towards this adverse outcome are still unmet needs. Alterations of retinal microcirculation have been independently associated with increased CVD risk in the general population (2). The potential changes of retinal vasculature and their association with disease activity and the magnitude of inflammation have not been studied in PSV so far.

Objectives: To explore the effect of disease activity and inflammation on retinal microcirculation in PSV, classified according to vessel size as large, medium or small vessel vasculitides (LVV, MVV or SVV respectively), and polymyalgia rheumatica (PMR) without vasculitis.

Methods: Fifty-nine patients, 43 with active disease [30 active vasculitis (12 LVV, 4 MVV, 14 SVV) and 13 PMR] and 16 with chronic inactive disease (13 vasculitis, 3 PMR) were studied. All patients were matched at 1:1 ratio with 59 controls, without underlying autoimmune/autoinflammatory disorder, neoplasia or infection, according to age, gender, CVD history, BMI, smoking, arterial hypertension, dyslipidemia, diabetes mellitus and treatment related to comorbidities. A total of 32 rheumatoid arthritis (RA) patients with mild to moderate inflammatory component as defined by standard of care acute phase reactants (ESR and CRP) and 16 chronic RA patients with normal ESR and CRP levels, matched 1:1 according to all the above parameters with 32 active and 16 inactive vasculitis/PMR patients respectively, served as disease controls. Digital retinal images were obtained and retinal vessel calibers were measured with a validated software to determine central retinal arteriolar and venular equivalents ratio (CRAE and CRVE respectively). For 16 patients with active vasculitis/PMR, retinal examination was performed in two time points (baseline and ≤ 6 months later).

Results: In the overall population, patients with active Vasculitis/PMR had increased CRVE (213.8 ± 21.7 vs 201.3 ± 17.1 , $p < 0.001$) and CRAE (180.0 ± 19.2 vs 164.1 ± 17.5 , $p < 0.001$) compared to healthy controls. Separating patients according to disease type, we found that in LVV, MVV and PMR, CRAE ($p \leq 0.05$) and CRVE ($p < 0.05$) were increased compared to control group, while in SVV only CRAE was increased ($p < 0.001$). Interestingly, chronic patients with disease in remission displayed higher CRAE compared to matched controls (179.8 ± 17.2 vs 169.1 ± 11.1 , $p = 0.006$). After immunosuppressive treatment for ≤ 6 months CRVE and to a lesser extent CRAE were reduced ($p = 0.048$ and 0.149 respectively) with a stronger statistical significance found in the Vasculitis group ($p = 0.026$ and 0.069 respectively). Pearson's linear correlation coefficient in active disease state (at baseline examination) revealed positive bivariate correlation only between CRVE with ESR and CRP. Patients with Vasculitis/PMR had also increased CRAE when compared to RA patients in both active and inactive disease status (182.8 ± 19.4 vs 170.1 ± 18.4 , $p < 0.001$ and 179.8 ± 17.2 vs 166.3 ± 17.9 , $p = 0.005$ respectively).

Conclusion: Systemic inflammation alters retinal microcirculation in both a reversible (venules) and irreversible (arterioles) way, independently of PSV form. Thus, common disease specific pathogenetic mechanisms related to inflammation may be implicated in vascular remodeling. Sequential follow-up of PSV patients will address whether retinal vessel calibers may serve as a biomarker of disease activity and CVD development.

REFERENCES:

- [1] Argyropoulou OD et al. *Curr Opin Rheumatol* 2017.
- [2] Shaohua G et al. *Current Atherosclerosis reports* 2020.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.1950

POS0808

CHARACTERISTICS OF GIANT CELL ARTERITIS FLARES AFTER SUCCESSFUL TREATMENT WITH TOCILIZUMAB: RESULTS FROM THE LONG-TERM EXTENSION OF A RANDOMIZED CONTROLLED PHASE 3 TRIAL

S. Unizony¹, S. Mohan², J. Han², J. H. Stone¹. ¹Harvard Medical School, Massachusetts General Hospital Rheumatology Unit, Boston, United States of America; ²Genentech, US Medical Affairs, South San Francisco, United States of America

Background: GiACTA investigated tocilizumab (TCZ) for the treatment of giant cell arteritis (GCA).

Objectives: To investigate disease flare characteristics after successful treatment with TCZ in GiACTA.

Methods: We report a post hoc analysis from part 2 of GiACTA. Part 1 was a 52-week, double-blind, randomized controlled period and part 2 was a 2-year open-label follow-up. In part 1, patients received TCZ 162mg subcutaneously every week or every other week with a 26-week prednisone taper or placebo plus a 26- or 52-week prednisone taper. Patients who were in remission at week 52 were to enter part 2 on no TCZ treatment. Part 2 treatment was at the investigator's discretion. We report characteristics of first disease flare in patients assigned to TCZ in part 1 who were in sustained remission at week 52 and experienced flare in part 2. Flare was defined as reappearance of cranial symptoms (headaches, jaw claudication, visual manifestations, scalp tenderness) or polymyalgia rheumatica (PMR) symptoms or elevation of erythrocyte sedimentation rate (ESR) ≥ 30 mm/h attributable to GCA that required treatment.