Results: Of 149 patients assigned to TCZ in part 1, 81 (54%) were in sustained remission on entering part 2. Of these 81 patients, 37 (46%) experienced at least one flare in part 2, including 17 with new-onset GCA and 20 with relapsing GCA at baseline. Median time to flare was 26.6 weeks. In patients with new-onset GCA, flares included cranial (53%) more often than PMR symptoms (18%). Cranial and PMR symptoms were balanced (both 60%) at the time of flare in patients with relapsing GCA. Visual manifestations occurred in two patients (5%) (Table 1). ESR and CRP were elevated in 65% and 36% of patients, respectively, at the time of flare. Three (8%) flares occurred with elevated ESR without clinical symptoms.

Table 1. Clinical manifestations during flare in part 2

	Part 1 Treatment ^a		
	TCZ QW+Pred	TCZ Q2W+Pred	All TCZ
New-onset disease			
Patients, n	28	14	42
Patients with ≥1 flare, n (%) ^b	9 (32.1)	8 (57.1)	17 (40.5)
Patients with ESR ≥30mm/h during flare, n (%) ^b	6 (66.7)	6 (75.0)	12 (70.6)
Patients with CRP ≥10mg/L during flare, n (%) ^b	4 (44.4)	2 (25.0)	6 (35.3)
Patients with GCA signs or symptoms during flare, n (%) ^c	7 (77.8)	7 (87.5)	14 (82.4)
PMR symptoms	1 (11.1)	2 (25.0)	3 (17.6)
Cranial symptoms ^d	4 (44.4)	5 (62.5)	9 (52.9)
Amaurosis fugax	0	0	0
Blurred vision	1 (11.1)	0	1 (5.9)
Diplopia	0	0	0
Blindness	0	0	0
Ischemic optic neuropathy	0	0	0
Fever	1 (11.1)	0	1 (5.9)
Other ^e	3 (33.3)	2 (25.0)	5 (29.4)
Relapsing disease			
Patients, n	28	11	39
Patients with ≥1 flare, n (%) ^b	14 (50.0)	6 (54.5)	20 (51.3)
Patients with ESR ≥30mm/h during flare, n (%) ^b	8 (57.1)	4 (66.7)	12 (60.0)
Patients with CRP ≥10mg/L during flare, n (%) ^b	5 (35.7)	3 (50.0)	8 (40.0)
Patients with GCA signs or symptoms during flare, n (%) ^c	14 (100)	6 (100)	20 (100)
PMR symptoms	8 (57.1)	4 (66.7)	12 (60.0)
Cranial symptoms ^d	8 (57.1)	4 (66.7)	12 (60.0)
Amaurosis fugax	1 (7.1)	0	1 (5.0)
Blurred vision	0	0	0
Diplopia	0	0	0
Blindness	0	0	0
Ischemic optic neuropathy	1 (7.1)	0	1 (5.0)
Fever	0	0	0
Other ^e	6 (4.3)	1 (16.7)	7 (35.0)

^aPatients from part 1 TCZ+Pred groups who were in sustained remission at week 52 entered part 2 on no treatment.

^bPercentage based on N in disease-onset group.

^cPercentage based on number of flare patients in disease-onset group. Individual signs or symptoms are shown as number of patients with each symptom; patients could have ≥1 sign or symptom at the time of flare.

^dNew-onset localized headache, scalp tenderness, temporal artery tenderness or decreased pulsation, ischemia-related vision loss, or jaw pain claudication.

^eIncludes fatigue, malaise, subjective weakness, and night sweats.

Conclusion: Overall, 46% of GCA patients successfully treated with TCZ for 1 year experienced disease flare within the next 2 years. Flares in patients with new-onset disease occurred more often with cranial than PMR symptoms. Visual manifestations were rare, and no blindness occurred. ESR and CRP were normal in a sizable percentage of patients experiencing flare.

Disclosure of Interests: Sebastian Unizony Consultant of: Sanofi and Kiniksa Pharmaceuticals, Grant/research support from: Genentech, Inc., Shalini Mohan Shareholder of: Genentech, Inc., Employee of: Genentech, Inc., Jian Han Shareholder of: Genentech, Inc., Employee of: Genentech, Inc., John H. Stone Consultant of: Roche/Genentech and Sanofi

DOI: 10.1136/annrheumdis-2021-eular.2602

POS0809

A BIOMARKER PROFILE AIDING AN EARLY DIAGNOSIS OF GIANT CELL ARTERITIS

Y. Van Sleen¹, P. Therkildsen², A. Boots¹, B. Dalsgaard Nielsen², K. Van der Geest¹, P. Heeringa³, M. G. Huitema¹, M. D. Posthumus⁴, M. Sandovici¹, E. Toonen⁵, J. Zijlstra¹, E. M. Hauge², E. Brouwer¹. ¹University Medical Center Groningen, Rheumatology and Clinical Immunology, Groningen, Netherlands; ²Aarhus University Hospital, Rheumatology, Aarhus N,

Denmark; ³University Medical Center Groningen, Pathology and Clinical Immunology, Groningen, Netherlands; ⁴Martini Ziekenhuis Groningen, Rheumatology, Groningen, Netherlands; ⁵Hycult Biotech, R&D Department, Uden, Netherlands

Background: Diagnosing patients with giant cell arteritis (GCA) remains difficult. Due to its non-specific symptoms, it is challenging to identify GCA in patients presenting with polymyalgia rheumatica (PMR), which is a more common disease (1). In addition, commonly used acute-phase markers fail to discriminate between GCA patients and (infectious) mimicry patients.

Objectives: To investigate a selection of biomarkers for their utility in the accurate diagnosis of GCA in two cohorts.

Methods: Treatment-naïve GCA patients participated in the Aarhus GCA/PMR cohort (N=52) and the Groningen GPS cohort (N=48). Symptoms and biomarker levels were compared to patients presenting phenotypically as isolated PMR, disease controls and healthy controls (HCs). Diagnosis or exclusion of diagnosis of GCA was based on clinical assessment and in the majority of cases aided by imaging. Serum/plasma levels of 12 biomarkers were measured by ELISA or Luminex.

Results: In both the Aarhus and the GPS cohort, we found that weight loss, elevated erythrocyte sedimentation rate (ESR) and higher angiopoietin-2/-1 ratios but lower matrix metalloproteinase (MMP)-3 levels identify concomitant GCA in PMR patients (Figure 1). In addition, we confirmed (1) that elevated platelet counts are characteristic of GCA but not of GCA look-alikes, and that low MMP-3 and proteinase 3 (PR3) levels may help to discriminate GCA from other diseases (Figure 1). Multiple biomarkers of inflammation were found elevated in patient and disease control groups when compared to HCs.

Conclusion: This study, performed in two independent cohorts, consistently shows the potential of angiopoietin-2/-1 ratios and MMP-3 levels to identify GCA in patients presenting with PMR. These biomarkers may be used to select which PMR patients require further diagnostic workup. Platelet counts may be used to discriminate GCA from GCA look-alike patients.

Figure 1. Summary of the most important and consistent findings in both cohorts. A shows the four factors that perform best in discriminating GCA/PMR patients overlap from isolated PMR patients in both cohorts. B shows the four factors that perform best in discriminate GCA patients from GCA look-alike patients in both cohorts. Cut-off values for the biomarkers are calculated by the Youden index.

REFERENCES:

- [1] van der Geest, KSM, Sandovici M, Brouwer E, Mackie SL. Diagnostic accuracy of symptoms, physical signs, and laboratory tests for giant cell arteritis: A systematic review and meta-analysis. JAMA internal medicine. 2020.

Disclosure of Interests: Yannick van Sleen: None declared, Philip Therkildsen: None declared, Annemieke Boots: None declared, Berit Dalsgaard Nielsen: None declared, Kornelis van der Geest: None declared, Peter Heeringa: None declared, Minke G. Huitema: None declared, M.D. Posthumus: None declared, Maria Sandovici: None declared, Erik Toonen Employee of: Is an employee of Hycult Biotech, Jannik Zijlstra: None declared, Ellen-Margrethe Hauge: None declared, Elisabeth Brouwer: None declared

DOI: 10.1136/annrheumdis-2021-eular.2636

POS0810

PULMONARY FINDINGS ON HIGH-RESOLUTION COMPUTED TOMOGRAPHY IN TAKAYASU ARTERITIS

X. Kong¹, J. Zhang², J. Lin³, P. LV³, H. Chen¹, Z. Ji¹, X. Dai¹, X. Jin⁴, L. Ma¹, L. Jiang¹. ¹Zhongshan Hospital Fudan University, Department of Rheumatology, Shanghai, China; ²Zhongshan Hospital Fudan University, Department of Pulmonary Medicine, Shanghai, China; ³Zhongshan Hospital Fudan University, Department of Radiology, Shanghai, China; ⁴Zhongshan Hospital Fudan University, Department of Clinical Epidemiology Center, Shanghai, China