POS0955

EXTRACELLULAR MATRIX PROTEIN TURNOVER MARKERS ARE ASSOCIATED WITH axSpA – A COMPARISON WITH CONTROL SUBJECTS WITH OR WITHOUT PELVIC, BUTTOCK OR BACK PAIN

H. Port^{1,2}, S. Holm Nielsen^{1,3}, J. Blair¹, S. F. Madsen¹, A. C. Bay-Jensen¹, M. Karsdal¹, S. Seven^{4,5}, I. J. Sørensen^{4,5}, L. Morsel-Carlson⁶, M. Østergaard^{4,5}, S. Juhl Pedersen⁴. ¹Nordic Bioscience A/S, Immunoscience, Herlev, Copenhagen, Denmark; ²Panum - University of Copenhagen, School of Health and Medical Sciences, København, Denmark; ³Technical University of Denmark, Biomedicine and Biotechnology, Kgs. Lyngby, Denmark; ⁴Rigshospitalet, Center for Arthritis Research, Glostrup, Denmark; ⁵University of Copenhagen, Clinical Medicine, København, Denmark; ⁶Frederiksberg Hospital, Radiology, Frederiksberg, Denmark

Background: Axial spondyloarthritis (axSpA) is a common chronic inflammatory disease, associated with extracellular matrix (ECM) remodeling of the cartilage, bone and connective tissues. The primary symptom of axSpA is back pain, caused by inflammation. However, there is a medical need to truly identify patients with axSpA from other subjects with buttock or low back pain attributable to other reasons. Tissue-derived extracellular (ECM) markers quantified in serum may serve to differentiate axSpA patients from other diagnostic entities.

Objectives: To investigate circulating ECM turnover markers as potential biomarkers to differentiate patients with axSpA from a control group of subjects with buttock or pelvic pain attributed to other reasons, including postpartum women and healthy subjects.

Methods: Biomarkers of ECM degradation/chronic inflammation (C1M, C3M, C4M, C6M, CRPM, C10C and COL10NC) and ECM formation (PRO-C3 and PRO-C6) were measured in 204 participants from the MASH study [1] (Table 1). Biomarker levels were compared among patients with axSpA and control group, and two new variables (Type 3 and Type 6) were included, corresponding to the index of formation/degradation of type III and type VI collagen respectively. The biomarker data was log10 transformed for normalization when necessary and linear regression models with pairwise comparisons were performed. Clinical scores and the SPARCC MRI scores for sacroiliac joint (SIJ) inflammation were correlated with the biomarkers data (data not shown).

Table 1. Comparison of blood-tested biomarkers levels in MASH study. Data was adjusted for confounders age, gender, and body mass index (BMI).

Biomarkers (Mean (SD))	Patients with axSpA (n = 41)	Control groups (n=163)	p
C1M	84.3 (85.8)	36.2 (22.1)	<0.0001
C3M	15.6 (4.0)	13.9 (3.0)	0.011
C4M	34.9 (10.2)	27.9 (7.7)	<0.0001
C6M	20.5 (5.8)	17.4 (4.2)	<0.0001
CRPM	11.9 (2.9)	11.0 (5.9)	0.027
C10C	2567 (462)	2568 (560)	0.31
COL10NC	9.15 (5.81)	9.43 (8.27)	0.43
PRO_C3	10.2 (2.5)	11.3 (3.0)	0.0052
PRO_C6	6.94 (2.45)	6.86 (2.53)	0.93
Type 3 (PRO-C3/C3M)	0.70 (0.28)	0.86 (0.31)	0.0004
Type 6 (PRO-C6/C6M)	0.36 (0.16)	0.41 (0.14)	0.0024

Results: We found that patients with axSpA had significantly increased MMP-mediated degradation of type I (C1M), type III (C3M), type IV (C4M) and type VI (C6M) collagens ($p<0.0001$, $p=0.01$, $p<0.001$, $p<0.001$, respectively), and CRP-metabolite (CRPM, $p=0.027$), and significantly decreased formation of type III collagen ($p=0.052$) compared with control groups when adjusting for age, gender and body mass index (BMI). Significantly decreased rates of Type 3 and Type 6 variables were also observed in patients with axSpA ($p=0.004$, $p=0.002$, respectively). We further adjusted the analyses by the SPARCC SIJ inflammation scores since differences in these were found among the two groups [1], and similar differences in biomarker levels were found, indicating that the relationship between the level of biomarkers and the SPARCC SIJ inflammation scores was independent between the two groups.

Conclusion: Biomarkers of type I, III, IV and V collagen and a CRP-metabolite showed an altered turnover in patients with axSpA compared with the control group including subjects with or without buttock or pelvic pain attributable

to other reasons. Such biomarkers may be used in combination with MRI or independently to separate patients with axSpA from other back pain conditions.

REFERENCES:

- [1] Seven, S., Østergaard, M., Morsel-Carlson, L., Sørensen, I. J., Bonde, B., Thamsborg, G., ... & Pedersen, S. J. (2019). Magnetic Resonance Imaging of Lesions in the Sacroiliac Joints for Differentiation of Patients With Axial Spondyloarthritis From Control Subjects With or Without Pelvic or Buttock Pain: A Prospective, Cross-Sectional Study of 204 Participants. *Arthritis & Rheumatology*, 71 (12), 2034-2046.

Acknowledgements: The study was supported by Innovation Fund Denmark, Rigshospitalet and the Danish Rheumatism Association (grant no. R102-A2132-B98, R159-A5061 and R150-A4467-B98).

Disclosure of Interests: Helena Port Employee of: Industrial PhD student at Nordic Bioscience and University of Copenhagen, Signe Holm Nielsen Employee of: Full time employee at Nordic Bioscience, Joseph Blair Employee of: Full time employee at Nordic Bioscience, Sofie Falkenløve Madsen Employee of: Full time employee at Nordic Bioscience A/S, Anne-Christine Bay-Jensen Shareholder of: Nordic Bioscience A/S, Employee of: Full time employee at Nordic Bioscience A/S, Morten Karsdal Shareholder of: Nordic Bioscience A/S, Employee of: Full time employee at Nordic Bioscience A/S, Sengül Seven: None declared, Inge Juul Sørensen: None declared, Lone Morsel-Carlson: None declared, Mikkel Østergaard: None declared, Susanne Juhl Pedersen: None declared

DOI: 10.1136/annrheumdis-2021-eular.786

POS0956

MEDITERRANEAN DIET IN AXIAL SPONDYLOARTHRITIS: A NUTRITIONAL INTERVENTION STUDY

F. Ometto¹, D. M. Farber¹, M. Lorenzin¹, A. Ortolan¹, G. Dellamaria², M. Felicetti¹, M. Favero¹, R. Valentini², A. Doria¹, R. Ramonda¹. ¹University of Padova, Rheumatology Unit, Department of Medicine, DIMED, Padova, Italy; ²University of Padova, Dietetics and Clinical Nutrition Unit, Department of Medicine DIMED, Padova, Italy

Background: Mediterranean diet (MD) proved to be beneficial on disease activity in patients affected by rheumatoid arthritis. A few data are available in other rheumatic conditions.

Objectives: A prospective monocentric study was conducted to evaluate the effect of a nutritional intervention based on the MD and to investigate the impact of such modification on disease activity of axial spondyloarthritis (ax-SpA).

Methods: Patients affected by ax-SpA randomized to a nutritional intervention arm (N) or a control arm (C) and were assessed at baseline (T0) and at month 6 (T6). Patients in N underwent a nutritional evaluation and received suggestions for dietary modification at T0, and then every 2 months. Adherence to the MD was evaluated with the PREDIMED questionnaire (range 0-10) [1]. A multivariable regression analysis was conducted to identify independent predictors of PREDIMED and of ASDAS-CRP improvement at T6.

Results: Eligible patients (161) were randomized to either N (81) or C (80); 47 in N and 63 in C completed the study. No relevant change of anthropometric or laboratory measures was observed at T6 in either group. A $\geq 20\%$ PREDIMED improvement was more frequent in N (22/47, 46.8%) compared to C (13/63, 20.6%) ($p<0.01$). Also, a $\geq 20\%$ ASDAS-CRP improvement was more frequent in N (8/47, 57.1%) compared to C (39/63, 40.6%) ($p=0.020$); and especially in those who achieved a $\geq 20\%$ PREDIMED improvement (12/35, 34.3%) vs those who did not (2/73, 2.7%) ($p<0.01$). The nutritional intervention, a higher age, and a lower BMI increased the odds of achieving a $\geq 20\%$ PREDIMED improvement; the $\geq 20\%$ PREDIMED improvement was associated higher odds of achieving a $\geq 20\%$ ASDAS-CRP improvement; psoriasis was negatively associated to the ASDAS-CRP improvement (Figure 1).

Conclusion: A nutritional intervention is effective in improving adherence to the MD in ax-SpA. A higher adherence to the MD may have a beneficial impact on activity of ax-SpA. Patients with psoriasis may have a limited benefit from the dietary improvement.

REFERENCES:

- [1] Martínez-González, M.A. et al. A 14-item mediterranean diet assessment tool and obesity indexes among high-risk subjects: The PREDIMED trial. *PLoS One* 2012,7.

Table 1. Characteristics of the patients.

	All patients	Nutrition	Controls	All patients	Nutrition	Controls
	Baseline			Month 6		
Females, n, (%)	40 (36.4)	18 (38.3)	22 (34.9)			
Age, years, mean±SD	51.7±1.3	53.0±1.3	49.6±1.3			
HLA-B27 positivity, n (%)	58 (52.7)	22 (46.8)	37 (57.1)			
Psoriasis, n (%)	58 (50.7)	26 (55.3)	32 (50.8)			
Disease duration (years), mean±SD	15.3±9.7	15.7±10	15±9.5			
Duration of b/tsDMARD treatment overall, years, mean±SD	5±4.1	5.8±4.5	4.5±3.8			
NSAID, n (%)	76 (69.1)	30 (63.8)	46 (73)			
csDMARD, n (%)	14 (12.7)	5 (10.6)	9 (14.3)			
BMI, Kg/meters ² , mean±SD	26.5±5.4	26.5±4.3	26.6±6.1	26.4±5.3	26.3±4	26.6±6.1
ASDAS-CRP, mean±SD	2.1±1	2.1±0.9	2.1±1	2±1.1	1.8±0.9	2.1±1.2
BASDAI, mean±SD	37.6±23	37.4±23.2	37.7±22.9	39.3±24.1	37.3±23.6	41.2±24.6
BASFI, mean±SD	20.5±21.4	21.6±19.2	19.7±23	19.8±19.6	19.1±18.8	20.5±20.4
BASMI, mean±SD	1.6±2	1.9±2.2	1.4±1.7	1.8±2.1	1.9±2.4	1.7±1.9
Tender joint count, mean±SD	1.1±2.3	0.8±2.1	1.4±2.5	0.9±2	0.8±2	0.9±1.9
Swollen joint count, mean±SD	0.3±1.3	0.4±2	0.1±0.5	0.2±1.3	0.3±1.8	0.2±0.8
Leeds Enthesitis Index, mean±SD	0.2±0.8	0.3±1	0.2±0.6	1.1±2	1.3±2.2	1±1.8
CRP, mg/L, mean ±SD	3.4±6.2	3.2±3.9	3.7±7.4	3.5±5.6	2.6±3.1	4.1±6.9
LDL-c, mg/dl, mean ±SD	130.8±36.5	132.3±35	129.7±38.4	130.6±34.7	125.7±34.8	134.4±34.5
PREDIMED score, mean ±SD	6.7±1.8	7±2.1	6.6±1.6	7.6±2.1	8.6±1.9	6.8±2

b/tsDMARD biological/targeted synthetic DMARDs; csDMARD conventional DMARDs. PREDIMED questionnaire to assess adherence to the Mediterranean diet.

Disclosure of Interests: None declared.

DOI: 10.1136/annrheumdis-2021-eular.812

POS0957 THE PREVALENCE AND FACTORS RELATED TO SLEEP APNOEA IN ANKYLOSING SPONDYLITIS

H. Forsblad-D'elia¹, A. Wigander², C. Sahlin-Inggridsson², M. Geijer³, K. Franklin², A. Blomberg². ¹Institute of Medicine, Department of Rheumatology and Inflammation Research, Gothenburg, Sweden; ²Umeå University, Department of Public Health and Clinical Medicine, Umeå, Sweden; ³Institute of Clinical Sciences, Department of Radiology, Gothenburg, Sweden

Background: An increased prevalence of obstructive sleep apnoea (OSA) has been suggested in ankylosing spondylitis (AS), but few controlled studies have been performed.

Objectives: We thus aimed to study the prevalence of OSA in patients with AS compared to controls and to study if disease-related and non-disease-related factors were determinants of OSA in AS patients.

Methods: One hundred and fifty-five patients with AS were included in the Backbone study that investigates severity and comorbidities in AS. Controls were recruited from the Swedish CardioPulmonary bio-Image Study (SCAPIS). Participants were asked to be examined with a home sleep-monitoring device during one night's sleep to evaluate the presence of OSA. For each AS patient, 45-70 years, four controls were matched for sex, age, weight and height. OSA was defined as an apnoea-hypopnea-index ≥5 events/hour.

Results: In total, 63/155(40.6%) patients with AS were examined with a home sleep-monitoring device out of which 46 patients were 45-70 years and therefore matched (mean age 57.2±7.5years, 30(65.2%) men) with 179 controls (mean age 57.2±4.5years, 123(68.7%) men). Twenty-two out of 46(47.8%) patients with AS vs. 91/179(50.8%) controls had OSA, p=0.72. No differences measurements evaluating OSA were noted in AS vs. controls. In logistic regression analysis, based on all 63 examined AS-patients, several AS-related variables were associated with OSA but after adjusting for age and sex, only higher age and BMI, remained to be significant determinants of OSA, Table 1.

Table 1. Univariable and age- and sex-adjusted logistic regression analyses with obstructive sleep apnoea as dependent variable in 63 patients with ankylosing spondylitis.

Variables	Univariable logistic regression analyses, Odds Ratio (95%CI)	P	Age- and sex-adjusted logistic regression analyses, Odds Ratio (95%CI)	P
Sex, male	1.9(0.6-5.5)	0.25	1.5(0.4-4.8)	0.53
Age	1.1(1.0-1.2)	0.002	1.1(1.0-1.2)	0.002
BMI	1.4(1.1-1.7)	0.001	1.6(1.2- 2.2)	0.001
Duration of symptoms	1.1(1.0-1.1)	0.028	1.0(0.9-1.1)	0.79
BASMI	1.9(1.3-2.9)	0.002	1.5(0.9 -2.5)	0.87
BASFI	1.4(1.0-2.0)	0.038	1.3(0.9-2.0)	0.88
≥1 Syndesmophyte	3.9(1.3-12.2)	0.017	3.0(0.8-11.3)	0.10
mSASS	1.0(1.0-1.1)	0.047	1.0(0.98-1.05)	0.25
Metabolic syndrome	4.3(1.5-12.9)	0.008	1.4(0.3-6.6)	0.69
Epworth Sleep Scale	1.2(1.0-1.3)	0.023	1.2(1.0-1.4)	0.29

Conclusion: In this case-control study, patients with AS did not have a higher prevalence of OSA compared to controls. AS patients with OSA had higher BMI and were older compared to patients without OSA.

Disclosure of Interests: None declared.

DOI: 10.1136/annrheumdis-2021-eular.823

POS0958 RESPONSIVENESS OF CONVENTIONAL, SEMI-AUTOMATIC AND FULL-AUTOMATIC METHODS TO QUANTIFY MARROW BONE EDEMA LESIONS IN MRI OF AXIAL SPONDYLOARTHRITIS PATIENTS: A PILOT STUDY

C. Garrido-González¹, M. L. Ladehesa Pineda², J. L. Garrido-Castro³, Z. Tabor⁴, I. Kucybała⁵, W. Wojciechowski⁵, P. Zarco-Montejo⁶, R. Almodovar⁶, E. Collantes Estevez² on behalf of CASTRO. ¹Universidad de Córdoba, Facultad de Medicina, Enfermería y Fisioterapia, Córdoba, Spain; ²Hospital Universitario Reina Sofía, Rheumatology, Córdoba, Spain; ³Maimónides Biomedical Research Institute of Córdoba, Rheumatology, Córdoba, Spain; ⁴AGH University of Science and Technology, Department of Biocybernetics and Biomedical Engineering, Krakow, Poland; ⁵Jagiellonian University Medical College, Department of Radiology, Krakow, Poland; ⁶Fundación Hospital Alcorcón, Rheumatology, Madrid, Spain

Background: The presence or absence of marrow bone edema (MBE) in the sacroiliac joints (SIJ) is very important in the diagnosis of axial Spondyloarthritis (axSpA). The quantification of this lesion and its extension may be important to analyze responsiveness of the treatment. Several scoring systems have been proposed for MRI images of SIJ, some of them being observer dependent (Berlin, SPARCC). Others, works in a semi-automatically way, such as the s-SCAISS[1], which makes it possible to quantify the size of the lesion, based on the indication of the expert. Recently, methods like the KITS4R[2], based on a deep patch-based classification network, allow a fully automated detection and quantification of the MBE lesions.

Objectives: To analyze responsiveness of several scores (observer independent, semiautomatic and full automatic) to quantify MBE in SIJ of axSpA patients.

Methods: Two rheumatologists independently quantified SIJ images from axSpA patients by visual inspection methods (Berlin and SPARCC indexes) and a semi-automatic system (s-SCAISS) on a single semi-coronal MRI slide (STIR). Full semi-coronal MRI images (15 to 18 slices) were used for an automatic detection algorithm (KITS4R), where total MBE was calculated as sum of areas of MBE in each slice. Patients were assessed before TNF-α therapy (PRE) and 3 months later (POST). Spearman correlations was used to analyze relationship between variables, Wilcoxon signed-rank test for significant differences and Cohen's d for calculating the effect size of improvement. Figure 1 shows processing of the MRI images: A) Area of MBE selected by the SCAISS, when the rheumatologist "click" on each MBE; B) Lesions detected automatically by KITS4R; C) Automated deep segmentation of bones and subchondral regions (with split into quadrants with central axes of joints also shown. D) Superposed A, B and C images.