[1]. The 3D-DXA and DXA measurements were compared using Student's *t* test. Categorical variables were compared using Chi-squared test. All hypothesis tests with *p* value lower than 5% were considered significant.

Results: Forty-six patients and 46 age and sex-matched controls were evaluated. The mean time of disease's duration was 7.37 years, and the most frequent JIA subtype was Oligoarticular ANA positive. More than 21% of the patients had an active disease. OP was present in 5 patients and 1 control. Despite not finding significant differences in the prevalence of osteoporosis, JIA patients had lower aBMD at total hip by DXA and lower cortical sBMD by 3D-DXA than controls (Table 1).

Table 1. Comparison of DXA and 3D-DXA measurements between JIA and healthy controls

Variable	JIA patients (n=46)	Healthy controls (n=46)	<i>P</i> value
DXA			
Lumbar aBMD (g/cm ²), mean ± sd	1.097 ± 0.02	1.155 ± 0.02	0.042
Right femoral neck aBMD (g/cm ²), mean ± sd	0.930 ± 0.02	1.014 ± 0.01	0.041
Right total hip aBMD (g/cm ²), mean ± sd	0.932 ± 0.02	1.001 ± 0.01	0.008
Lumbar T-score, mean ± sd	-0.716 ± 0.16	-0.212 ± 0.16	0.037
Right femoral neck T-score, mean ± sd	-0.563 ± 0.17	0.087 ± 0.12	0.036
Right total hip T-score, mean ± sd	-0.721 ± 0.18	-0.12 ± 0.13	0.008
3D-DXA			
Trabecular vBMD (mg/cm ³), mean ± sd	180.44 ± 5.54	194.57 ± 5.11	0.064
Cortical sBMD (mg/cm ³), mean ± sd	147.23 ± 3.41	161.52 ± 2.89	0.002
Osteoporosis (WHO definition)			
Yes, n (%)	5 (10.87)	1 (2.17)	0.091
No, n (%)	41 (89.13)	45 (97.83)	

DXA dual-energy X-ray absorptiometry, *sd* standard deviation, *aBMD* areal bone mineral density, 3D-DXA three-dimensional dual-energy X-ray absorptiometry, *vBMD* volumetric bone mineral density, *sBMD* superficial cortical bone mineral density

Conclusion: JIA patients do not have more OP than controls in our cohort but there are differences in BMD in the different locations and greater involvement of the cortical bone in JIA.

REFERENCES:

[1] Brance ML, et al. Trabecular and cortical bone involvement in rheumatoid arthritis by DXA and DXA-based 3D modelling. *Osteoporos Int*, 2020 Sep 24.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.742

POS1297

HOW HLA B27 PREVALENCE IN THE POPULATION INFLUENCE ON FEATURES OF JUVENILE IDIOPATHIC ARTHRITIS: DATA FROM SAKHA REPUBLIC (YAKUTIA)

F. Vinokurova¹, L. Sorokina², V. Argunova³, P. Sleptsova³, T. Bourtseva^{4,5}, V. Chasnyk², M. Kostik². ¹Yakut Science Center of Complex Medical Problems, Laboratory of Immunological Researches, Yakutsk, Russian Federation; ²Saint-Petersburg State Pediatric Medical University, Department of Hospital Pediatrics, St. Petersburg, Russian Federation; ³Republic Hospital № 1 - National Center of Medicine, Pediatric Center, Department of Cardio-Rheumatology, Yakutsk, Russian Federation; ⁴North-Eastern Federal University (NEFU), Department of Pediatrics and Pediatric Surgery of Medical Institute NEFU, Yakutsk, Russian Federation; ⁵Yakut Science Center of Complex Medical Problems, Laboratory of Children Health Monitoring and Medical and Environmental Research, Yakutsk, Russian Federation

Background: Sakha Republic (Yakutia) - SR(Y) is a bid arctic region of Russia with high proportion of aboriginals - Yakutians (50%), intra-national marriages, increased level of inbreeding and high distribution of HLAB27 among aboriginals - 33%, according the epidemiological studies. The main type of arthritis is ankylosing spondylitis in adults and enthesitis-related arthritis (ERA) of juvenile idiopathic arthritis (JIA) in children. The pattern of arthritis distribution in adults and children in SR(Y) is differ from Caucasians and similar to Native Americans.

Objectives: Our study aimed to evaluate the features of JIA in aboriginals of SR(Y), associated with high prevalence of HLAB27 antigen.

Methods: In the retrospective study we included 144 Yakutians who were admitted in the rheumatology department of Yakutsk in 2007-2016 years and 753 JIA patients Caucasian origin in Saint-Petersburg in the same years. We evaluated routine clinical and laboratorial features. HLA B27 was evaluated according the clinical judgement of the attending physicians.

Results: ERA is a main JIA category in Yakutians. The main features the male predominance, higher onset age, high inflammatory activity, lower number of active joints, high incidence of hip and sacroiliac joints involvement, lower levels of psoriasis and uveitis. Yakutians have rare the involvement of cervical spine (6% vs 14%, *p*=0.0000001), TMJ (1% vs 6%, *p*=0.027), elbow (8% vs 16%, *p*=0.012), wrist (18% vs 28%, *p*=0.017), MCP (7% vs 21%, *p*=0.00005), PIP (8% vs 25%, *p*=0.00005). The treatment rates of methotrexate and biologics were similar between groups, but

in Yakutians the biologics were administered earlier, because methotrexate failed and often was ineffective. The cumulative probability to receive biologics was higher in Yakutians compare to Caucasians: HR=3.4 [2.6; 4.4], *p*=0.000001 (Figure 1). The main biologic in Yakutians was etanercept (49/70; 70%). Yakutians received corticosteroids and cyclosporine A rarely, due to low incidence of systemic onset JIA and oligoarthritis with uveitis. It was observed, that the HLA B27 as a risk factors had different significance in Yakutians and Caucasians. HLA B27 presence increased the risk of ERA OR=2.72 (1.3; 5.6) *p*=0.01 in Yakutians and OR=69.2 (29.5; 162.3), *p*=0.00001 for Caucasians; for biologic administration: 0.86 (0.4; 1.8), *p*=0.529 and 2.45 (1.5; 4.0), *p*=0.0003, respectively.

Figure 1. Cumulative probability to leave without biologics between Yakutian and Caucasian JIA patients.

Conclusion: High distribution of HLA B27 antigene in Yakutians, lead to different pattern of JIA categories distributions and patient's management.

This work was supported by the Project of the Ministry of Science and Higher Education of the Russian Federation (basic part of funding to M.K. Ammosov North-Eastern Federal University #FSRG-2020-0016) and by the RFBR grant #18-05-600035_Artika.

Table 1. Differences between Yakutian and Caucasian JIA patients.

JIA features	Yakutians, n=144 (%)	Caucasians, n=723 (%)	<i>p</i>
Gender, boys, n (%)	85 (59.0)	279 (38.6)	0.000006
Onset age, years	10.6 (6.0; 13.4)	6.0 (3.0; 10.3)	0.0000001
JIA categories, n (%)			
Oligoarthritis	36 (25.0)	188 (26.0)	0.0000001
Poly, RF (-)	21 (14.6)	248 (34.3)	
Poly, RF (+)	1 (0.7)	22 (3.0)	
Systemic onset	4 (2.8)	53 (7.3)	
ERA	76 (52.8)	171 (23.7)	
Psoriatic arthritis	6 (4.2)	41 (5.7)	
Active joints	4.0 (3.0; 6.0)	6.0 (3.0; 12.0)	0.0000001
Uveitis, n (%)	16 (11.1)	114/503 (22.7)	0.002
Psoriasis, n(%)	3/143 (2.1)	46/719 (6.4)	0.043
Hip involvement	50 (34.7)	146 (20.2)	0.0001
Sacroiliitis	46 (31.9)	69 (9.6)	0.0000001
HLA B27, n (%)	76/131 (58.0)	105/301 (34.8)	0.00001
ANA, n (%)	4/46 (8.7)	204/444 (46.0)	0.00001
RF, n (%)	3/141 (2.1)	22/403 (5.5)	0.104
Biologics, primary, n (%)	70/144 (48.6)	347 (48.0)	0.892
Time before first biologics, years	0.6 (0.3; 1.4)	2.6 (1.0; 5.4)	0.0000001

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.810

POS1298

COMPARISON OF FMF CLINICAL FEATURES BETWEEN TURKISH AND CRIMEAN TATAR CHILDREN

O. Zhogova¹, U. Kaya Akca², E. Susptsin^{3,4}, E. Sag², V. Nizhnik³, A. Tumakova³, S. Ivanoskiy¹, N. Lagunova¹, Y. Bilgine², S. Özen², M. Kostik^{3,5}. ¹Crimean Federal University named after V.I. Vernadsky, Pediatrics, Simferopol, Russian Federation; ²Hacettepe University, Department of Pediatric Rheumatology, Faculty of Medicine, Ankara, Turkey; ³Saint Petersburg State Pediatric Medical University, Hospital Pediatrics, Saint-Petersburg, Russian Federation; ⁴N.N. Petrov National Research Center of Oncology, Molecular Diagnostics, Saint-Petersburg, Russian Federation; ⁵Almazov National Medical Research Centre,